


Carboxylic acids and derivatives

A-Level Chemistry

Carboxylic acids




Citrus fruits taste sour because they contain citric acid, a carboxylic acid

Image: Ivar Leidus, CC BY-SA 4.0 (commons.wikimedia.org)

Making and reacting

- an **alkylbenzene** 烷基苯 (such as methylbenzene) is oxidised by hot alkaline KMnO_4 , then dilute acid, to give **benzoic acid** 苯甲酸. The whole side-chain becomes a $-\text{COOH}$ group.
- a carboxylic acid reacts with PCl_3 and heat, PCl_5 , or SOCl_2 to form an **acyl chloride** 酰氯.



Hot KMnO_4 oxidises a methyl side-chain to a COOH group


Acids that can be oxidised further

Two **carboxylic acids** 羧酸 are special because they can still be oxidised:


- methanoic acid (HCOOH) is oxidised by Fehling's or Tollens' reagent, or acidified KMnO_4 / $\text{K}_2\text{Cr}_2\text{O}_7$, to carbon dioxide and water.
- ethanedioic acid (HOOC-COOH) is oxidised by warm acidified KMnO_4 to carbon dioxide.

Relative acidities

The **acidity** 酸性 order is:



A carboxylic acid is the strongest because, when it loses H^+ , the negative charge is spread over **two** oxygen atoms, making the ion very stable. In a **phenol** 苯酚 the charge spreads only into the ring, and in an **alcohol** 醇 (giving RO^-) it is not spread at all.



Acidity rises from alcohol to phenol to carboxylic acid: the more the negative charge on the conjugate base is spread out, the more stable the ion and the stronger the acid

Chlorine atoms make a carboxylic acid **more** acidic. Chlorine is **electron-withdrawing** 吸电子: through the **inductive effect** 诱导效应 it pulls electron density away, helping to spread the negative charge and stabilise the ion. So more chlorine atoms (closer to the $-\text{COOH}$) give a stronger acid.

Esters

An alcohol (or phenol) reacts with an acyl chloride at room temperature to give an **ester** 酯 and HCl . Examples are ethyl ethanoate (from ethanol) and phenyl benzoate (from phenol).




Aspirin is an ester, made from salicylic acid

Image: Ragesoss, CC BY-SA 4.0 (commons.wikimedia.org)

Acyl chlorides

Acyl chlorides are made from carboxylic acids (with PCl_3 , PCl_5 or SOCl_2). They are very reactive. At room temperature they react with:


Reactant	Product (plus HCl)
water	the carboxylic acid
an alcohol	an ester
phenol	an ester
ammonia 氨	an amide 酰胺
a primary or secondary amine 胺	an amide



Acyl chlorides are very reactive: with water, an alcohol, phenol, ammonia or an amine they give the labelled product plus HCl

The addition–elimination mechanism


All these reactions follow an **addition–elimination** 加成消去 mechanism: a nucleophile first **adds** to the slightly positive carbonyl carbon, then HCl is **eliminated**.



Addition–elimination: the nucleophile adds to the $\delta+$ carbonyl carbon, then the $C=O$ reforms and Cl^- leaves as HCl

Ease of hydrolysis

Compare how easily three chlorides react with water (**hydrolysis** 水解):



An acyl chloride reacts violently with cold water; an alkyl chloride reacts slowly; an aryl

chloride (halogenoarene) does not react, because its C–Cl bond is strengthened by the ring.

Worked example. Ethanoyl chloride reacts violently with cold water, while ethyl ethanoate needs prolonged reflux with acid or alkali. Explain the difference. Both are attacked at the **carbonyl carbon**, so compare how open that carbon is to attack and how good the leaving group is. In ethanoyl chloride the chlorine is strongly **electronegative** and withdraws electrons, making the carbonyl carbon much more $\delta+$ and so far more open to a nucleophile; and Cl^- , the anion of a strong acid, is a **good leaving group**. In the ester the $-\text{OR}$ oxygen **donates** a lone pair into the carbonyl, reducing that $\delta+$ charge, and RO^- is a **poor** leaving group. So the acyl chloride hydrolyses far more easily. Argue from **both** the $\delta+$ on the carbon **and** the leaving group: either one alone is usually only half the marks.

Exam tips

- **Acyl chlorides** are very reactive —learn their products with water (HCl), alcohols (esters), ammonia (amides) and amines.
- Reactivity order of derivatives: **acyl chloride** > **ester** > **amide**.
- Acid strength: electron-withdrawing groups (e.g. Cl) **increase** it —explain via the carboxylate ion.