



AP[®] Physics C: Mechanics 2016 Scoring Guidelines

© 2016 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.
Visit the College Board on the Web: www.collegeboard.org.
AP Central is the official online home for the AP Program: apcentral.collegeboard.org.




AP[®] PHYSICS C: MECHANICS 2016 SCORING GUIDELINES

Question 1

15 points total


Distribution
of points

(a) 3 points



For correctly drawing and labeling the force of tension 1 point
For correctly drawing and labeling the force of friction 1 point
For correctly drawing and labeling both forces in the vertical direction 1 point
Note: A maximum of two points may be earned if there are any extraneous vectors.

(b)
i. 3 points



For a correct scale that uses more than half the grid 1 point
For correctly plotting the given data 1 point
For drawing a straight line consistent with the given data 1 point
Note: Full credit can be earned if the axes are switched.

**AP[®] PHYSICS C: MECHANICS
2016 SCORING GUIDELINES**

Question 1 (continued)

**Distribution
of points**

- (b)
ii. 2 points
- For correctly calculating slope using the best-fit straight line and not data points 1 point
- $$\text{slope} = \frac{(y_2 - y_1)}{(x_2 - x_1)} = \frac{(0.70 - 0.16)}{(0.60 - 0.35)} \text{ kg}^{-1} = 2.16 \text{ kg}^{-1}$$
- Note: Linear regression gives slope = 2.12 kg⁻¹.
For correctly calculating the mass of the cart using the slope 1 point
- $$m = \frac{1}{\text{slope}} = \frac{1}{(2.16 \text{ kg}^{-1})}$$
- Correct answer:
 $m = 0.463 \text{ kg}$ (Note: linear regression gives $m = 0.472 \text{ kg}$)
- iii. 1 point
- For an answer with correct units consistent with the x-intercept of the graph from (b) i. 1 point
- $$f = 0.272 \text{ N}$$
- (c)
i. 1 point
- Applying Newton's second law and substituting the values from part (b)
- $$\sum F = ma = F_a - f$$
- $$a = \frac{F_a - f}{m} = \frac{0.45 \text{ N} - 0.272 \text{ N}}{0.463 \text{ kg}}$$
- For an answer with correct units consistent with part (b), either from the graph or calculated using the mass and frictional force. 1 point
- $$a = 0.376 \text{ m/s}^2$$

**AP[®] PHYSICS C: MECHANICS
2016 SCORING GUIDELINES**

Question 1 (continued)

**Distribution
of points**

- (c)
ii. 3 points
- For using a correct equation to solve for the speed of the cart when the string breaks, with an acceleration consistent with part (c) i. 1 point
- $$v_2 = v_1 + at$$
- $$v_2 = 0 + (0.376 \text{ m/s}^2)(2.0 \text{ s})$$
- $$v_2 = 0.752 \text{ m/s}$$
- For recognizing that the acceleration of the cart after the string breaks is due to the frictional force determined in part (b) 1 point
- Use a correct equation to solve for the time for the cart to stop after the string breaks
- $$v_2 = v_1 + at$$
- $$0 = v_1 - \frac{f}{m}t$$
- For using the final velocity before the string breaks as the initial velocity for the cart stopping in the correct equation for time 1 point
- $$t = \frac{mv_1}{f}$$
- $$t = \frac{(0.463 \text{ kg})(0.752 \text{ m/s})}{(0.272 \text{ N})}$$
- Correct answer
 $t = 1.28 \text{ s}$
- Alternate solution* *Alternate points*
- For setting the magnitude of the impulse before the string breaks equal to the magnitude of the impulse after the string breaks 1 point
- $$F_1 t_1 = F_2 t_2$$
- For correctly using the proper force (e.g., the tension force minus the friction force) for F_1 1 point
- For correctly using the proper force (e.g., the friction force) for F_2 1 point
- Correct answer
 $t = 1.28 \text{ s}$
- (d)
i. 1 point
- For selecting "Equal to" 1 point
- ii. 1 point
- For selecting "Greater than" 1 point

**AP[®] PHYSICS C: MECHANICS
2016 SCORING GUIDELINES**

Question 2

15 points total

(a) 2 points



For correctly drawing and labeling vectors on the block of mass $2M$
 For correctly drawing and labeling vectors on the block of mass $3M$ using symbols
 for the vectors that are physically correct and different from those on the block
 of mass $2M$
 Note: A maximum of one point can be earned if there are any extraneous vectors.

1 point
 1 point

(b) 2 points

Using a proper expression for conservation of momentum

$$p_1 = p_2$$

For correctly substituting into the above equation

$$3Mv_0 = (3M + 2M)v_f$$

For a correct answer

$$v_f = \frac{3}{5}v_0$$

1 point
 1 point

(c) 1 point

Using a proper expression for kinetic energy of the two-block system

$$K = \frac{1}{2}mv^2$$

$$K = \frac{1}{2}(5M)\left(\frac{3}{5}v_0\right)^2$$

For an answer consistent with part (b)

$$K = \frac{9}{10}Mv_0^2$$

1 point

**Distribution
of points**

**AP[®] PHYSICS C: MECHANICS
2016 SCORING GUIDELINES**

Question 2 (continued)

**Distribution
of points**

(d) 4 points

For a correct expression of the conservation of energy

$$\Delta K_{\text{system}} + \Delta U_{\text{system}} = 0$$

$$K_0 = U_{\text{final}}$$

For attempting to integrate the spring force equation

$$\frac{9}{10}Mv_0^2 = -\int_{x_1}^{x_2} Bx^3 dx$$

$$\frac{9}{10}Mv_0^2 = \int_{x_1}^{x_2} Bx^3 dx$$

For using the correct limits of integration or an appropriate constant of integration

$$\frac{9}{10}Mv_0^2 = \int_0^D Bx^3 dx$$

$$\frac{9}{10}Mv_0^2 = \left[\frac{Bx^4}{4} \right]_0^D$$

For an answer consistent with the speed from (b) or the kinetic energy from part (c)

$$D = \sqrt[4]{\frac{18Mv_0^2}{5B}}$$

1 point

1 point

1 point

1 point

(e)

i. 2 points

For selecting the correct answer "Right" with a reasonable attempt at a justification

If the incorrect selection is made, no points are earned for the justification.

For an indication that at maximum compression the block of mass $2M$ has an acceleration to the right due to the forces acting on the block of mass $2M$ or an acceleration to the right due to the external spring force acting on the system of blocks

Example: At maximum compression the two-block system is instantaneously at rest. The only horizontal external force acting on the system is due to the spring. This force is directed to the right. The system and therefore the block of mass $2M$ is accelerated to the right, which implies that the net force acting on the block of mass $2M$ is also to the right.

1 point

1 point

**AP[®] PHYSICS C: MECHANICS
2016 SCORING GUIDELINES**

Question 2 (continued)

**Distribution
of points**

- (e) ii. 2 points
- The magnitude of the net force is greater on the block of mass $3M$.
If the incorrect selection is made, no points are earned for the justification.
For an indication that both blocks will have the same acceleration 1 point
For a correct justification for why the net force is greater on the block of mass $3M$ 1 point
Example:
Because the blocks stick together, both blocks must have the same acceleration.
Because the block of mass $3M$ has more mass, the net force on it must be greater than the net force on the block of mass $2M$.
- (f) 2 points
- For selecting the correct answer "No," with a reasonable attempt at a justification 1 point
If the incorrect selection is made, no points are earned for the justification.
If the correct answer is selected without any justification, the point is not earned for the selection.
For an indication that the spring does not apply a linear force and that simple harmonic motion is the resulting motion of a linear restoring force 1 point
$$\left(a = -\frac{k}{m}\Delta x\right)$$

Example:
Because the blocks are sticking together and are attached to the spring, the spring will apply a restoring force to the blocks. However, because the restoring force exerted by a nonlinear spring is not proportional to the blocks' displacement from equilibrium, the blocks do not exhibit simple harmonic motion.

**AP[®] PHYSICS C: MECHANICS
2016 SCORING GUIDELINES**

Question 3

15 points total

**Distribution
of points**

- (a) 3 points
- For an indication that the spring force equals the mass times the centripetal acceleration 1 point
$$F_s = F_c = ma_c$$

$$kx = mr\omega^2$$

For a correct substitution for x in the equation 1 point
For a correct substitution for r in the equation that can be used to solve for k 1 point
$$k\left(\frac{d}{2}\right) = m(2d)\omega^2$$

Correct answer:
$$k = 4m\omega^2$$
- (b) i. 1 point
- Substitute values for the block into the equation for rotational inertia
$$I = \sum mr^2 = m(2d)^2$$

For a correct answer or an answer consistent with the radius used in part (a) 1 point
$$I = 4md^2$$
- ii. 2 points
- For an indication that the rotational inertia of the system must include the platform, the rod, and the object 1 point
$$I = I_P + I_R + I_O$$

$$I = \frac{m_P R^2}{2} + \frac{m_R d^2}{3} + mr^2$$

For correctly substituting into the equation the rotational inertia of the platform, the rod, and the object consistent with (b) i. 1 point
$$I = \frac{5m(2d)^2}{2} + \frac{3m d^2}{3} + 4md^2$$

$$I = (10 + 1 + 4)md^2$$

Correct answer:
$$I = 15md^2$$
- (c) 1 point
- Using a correct expression of angular momentum
$$L = I\omega$$

For an answer consistent with the answer from part (b) ii. 1 point
$$L = 15md^2\omega$$

**AP[®] PHYSICS C: MECHANICS
2016 SCORING GUIDELINES**

Question 3 (continued)

**Distribution
of points**

(d) 3 points

For indicating that the spring force equals the mass times the centripetal acceleration

1 point

$$F_s = F_C = ma_C$$

$$kx = mr\omega^2$$

For a correct substitution for k , or a substitution consistent with part (a), in the equation

1 point

For a correct substitution for r in the equation that can be used to solve for x

1 point

$$(4m\omega^2)x = m(d+x)\omega^2$$

$$4x = d + x$$

$$3x = d$$

Correct answer:

$$x = d/3$$

(e) 2 points

For selecting "Decreasing"

1 point

If the wrong selection is made, no points are earned for the justification.

For a correct justification

1 point

Example: The rotational inertia I of the system decreases while the angular velocity ω stays the same. Because the angular momentum is $I\omega$, it decreases.

(f) 2 points

For selecting "Negative"

1 point


If the wrong selection is made, no points are earned for the justification.

For a correct justification

1 point

Example: Because the angular velocity is constant and the rotational inertia has decreased, the rotational kinetic energy has decreased. Therefore, work must be negative.

(g) 1 point



For an arrow starting on the box and pointing down and to the right

1 point