

Trigonometry

IGCSE Mathematics


This handout covers Topic 6, Trigonometry. Parts marked **(Extended)** are only tested on the Extended papers; everything else is for both levels. Give angle answers to one decimal place.

Pythagoras' theorem

Pythagoras' theorem 勾股定理 links the three sides of a **right-angled triangle** 直角三角形 (a **triangle** 三角形 with one 90° angle). If the longest side (the **hypotenuse** 斜边, opposite the right angle) is c , then

$$a^2 + b^2 = c^2.$$

Use it to find a missing side.



Pythagoras as areas: the squares on the two shorter sides ($9 + 16$) add up to the square on the hypotenuse (25).

Worked example. A right-angled triangle has a hypotenuse of 13 cm and one short side of 5 cm. Find the other short side.


$$b^2 = 13^2 - 5^2 = 169 - 25 = 144, \quad b = \sqrt{144} = 12 \text{ cm.}$$

Trigonometry in right-angled triangles

Label the sides from the angle you are using: the **opposite** 对边 (across from the angle), the **adjacent** 邻边 (next to the angle), and the hypotenuse. The three ratios are the **sine** 正弦, **cosine** 余弦 and **tangent ratio** 正切 (sin, cos, tan):

$$\sin \theta = \frac{\text{opp}}{\text{hyp}}, \quad \cos \theta = \frac{\text{adj}}{\text{hyp}}, \quad \tan \theta = \frac{\text{opp}}{\text{adj}}.$$

Remember them as **SOH-CAH-TOA**. To find an angle, use the inverse ($\sin^{-1}$, $\cos^{-1}$, $\tan^{-1}$).



Name the sides from the angle θ : the opposite is across from it, the adjacent next to it, and the hypotenuse opposite the right angle (SOH-CAH-TOA).

Worked example (find a side). In a right-angled triangle the hypotenuse is 10 cm and the angle is 30° . Find the opposite side.

$$\text{opp} = 10 \times \sin 30^\circ = 10 \times 0.5 = 5 \text{ cm}.$$

Worked example (find an angle). The opposite side is 4 cm and the adjacent side is 3 cm.

$$\tan \theta = \frac{4}{3}, \quad \theta = \tan^{-1}\left(\frac{4}{3}\right) = 53.1^\circ.$$

Angles of elevation and depression (Extended)




Looking up at a tower involves an angle of elevation.

Image: Another one of my pictures:

This photograph was taken by Medium69 (William Crochot) and released under the license stated below. You are free to use it for any purpose as long as you credit the author (William Crochot), the Source (Wikimedia Commons) and the license (CC-BY-SA 4.0) in close relation to the image.

Please do not upload an updated image here without consultation with the Author. The author would like to make corrections only at his own source RAW. This ensures that the changes are preserved. Please if you think that any changes should be required, please inform the author. Otherwise you can upload a new image with a new name. Please use one of the templates derivative or extract, CC BY-SA 4.0 (commons.wikimedia.org)

The **angle of elevation** 仰角 is the angle up from the horizontal to an object above you. The **angle of depression** 俯角 is the angle down from the horizontal to an object below you. The shortest distance from a point to a line is the **perpendicular** 垂直 distance.



The angle of elevation looks up from the horizontal; the angle of depression looks down.


Worked example. From a point 50 m from the foot of a tower, the angle of elevation to the top is 40° . Find the height of the tower.

$$\text{height} = 50 \times \tan 40^\circ = 50 \times 0.839 = 42.0 \text{ m.}$$


Exact trigonometric values (Extended)

You must know these exact values without a calculator.

x	0°	30°	45°	60°	90°
$\sin x$	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1
$\cos x$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0
$\tan x$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	—



45° triangle



30° – 60° triangle

These two special triangles are where the exact values come from —worth memorising.

Graphs and trigonometric equations (Extended)




A Ferris wheel: a point on the rim traces a sine curve as it turns.

Image: Bob Collowan, CC BY-SA 3.0 (commons.wikimedia.org)

For $0^\circ \leq x \leq 360^\circ$:

- $y = \sin x$ is a wave starting at 0, peaking at 90° , back to 0 at 180° , down to -1 at 270° .
- $y = \cos x$ is the same wave but starting at 1.
- $y = \tan x$ rises steeply and repeats every 180° .



$y = \sin x$ and $y = \cos x$ are smooth waves between -1 and 1 ; cosine is sine shifted left by 90° .

A **trigonometric equation** 三角方程 often has more than one answer in this range. Use the graph (or the symmetry of the wave) to find them all.

Worked example. Solve $\sin x = \frac{\sqrt{3}}{2}$ for $0^\circ \leq x \leq 360^\circ$.

One answer is $x = 60^\circ$. The sine wave is also $\frac{\sqrt{3}}{2}$ at $180^\circ - 60^\circ = 120^\circ$. So $x = 60^\circ$ or 120° .

Worked example. Solve $2 \cos x + 1 = 0$ for $0^\circ \leq x \leq 360^\circ$.


$$\cos x = -\frac{1}{2} \Rightarrow x = 120^\circ \text{ or } 240^\circ.$$

The sine and cosine rules (Extended)

For **any** triangle (not just right-angled), with sides a, b, c opposite angles A, B, C :

$$\text{sine rule: } \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C},$$

$$\text{cosine rule: } a^2 = b^2 + c^2 - 2bc \cos A.$$



In any triangle, sides a, b, c lie opposite the angles A, B, C of the same letter.

Use the **sine rule** 正弦定理 when you have a side and its opposite angle. Use the **cosine rule** 余弦定理 when you have two sides and the angle between them, or all three sides. With the sine rule, watch for the **ambiguous case** 两解情况, where an angle could be acute or obtuse.

The area of any triangle is

$$\text{area} = \frac{1}{2}ab \sin C.$$

Worked example. A triangle has $b = 7$ cm, $c = 8$ cm and the angle between them $A = 40^\circ$. Find side a .

$$a^2 = 7^2 + 8^2 - 2(7)(8) \cos 40^\circ = 113 - 112 \times 0.766 = 27.2,$$

$$a = \sqrt{27.2} = 5.2 \text{ cm.}$$

Pythagoras and trigonometry in 3D (Extended)

In three dimensions, find a right-angled triangle inside the solid, then use Pythagoras or trigonometry on it. A common task is the angle between a line and a flat surface (a **plane** 平面).

Worked example. A box has a base 6 cm by 8 cm and height 5 cm. Find the angle between a space diagonal and the base.

First the base diagonal: $\sqrt{6^2 + 8^2} = \sqrt{100} = 10$ cm. This diagonal and the height form a right-angled triangle, so the angle θ with the base is

$$\tan \theta = \frac{5}{10} = 0.5, \quad \theta = \tan^{-1}(0.5) = 26.6^\circ.$$

Exam tips

- Use **Pythagoras** ($a^2 + b^2 = c^2$) when there is no angle; use **SOH-CAH-TOA** when an angle is involved.
- The **hypotenuse** is always opposite the right angle. Label the sides (opposite, adjacent, hypotenuse) relative to the angle you are using.
- To find an angle, use the inverse ($\sin^{-1}$, $\cos^{-1}$, $\tan^{-1}$), and check your calculator is set to **degrees**.
- For a triangle with no right angle, use the **sine rule** or the **cosine rule** —the cosine rule when you know two sides and the angle between them.