

6. The small invertebrate krill species *Thysanoessa inermis* is adapted to cold (4°C) seawater. Over the past ten years, there has been a gradual increase in the water temperature of the krill's habitat. A sustained increase in water temperature may ultimately affect the ability of the krill to survive.

One effect of higher temperatures is protein misfolding within cells. Krill have several *hsp* genes that code for heat-shock proteins (HSPs). These proteins help prevent protein misfolding or help to refold proteins to their normal shapes.

Scientists conducted experiments on *T. inermis* to detect changes in the expression of *hsp* genes when the krill were exposed to temperatures above 4°C. An experimental group of krill was maintained in tanks with 4°C seawater and then placed into tanks with 10°C seawater for approximately three hours. The krill were then given a six-hour recovery period in the 4°C seawater tanks. A control group of krill was moved from a tank of 4°C seawater to another tank of 4°C seawater for approximately three hours and then returned to the original tank. The scientists analyzed *hsp* gene expression by measuring the concentrations of three mRNAs (I, II, III) transcribed from certain *hsp* genes in both the heat-shocked krill (Figure 1) and the control krill. For the control krill, no transcription of the *hsp* genes was detected throughout the test period (data not shown).




Figure 1. Average concentration of three mRNAs (I, II, III) transcribed from *hsp* genes in krill heat shocked at 10°C. Error bars represent $\pm 2SE_{\bar{x}}$.

- (a) **Identify** the *hsp* mRNA that has the slowest rate of concentration increase in response to heat-shock treatment.
- (b) **Describe** the trend in the average concentration of mRNA I throughout the experiment.
- (c) The scientists hypothesized that the heat-shock protein (HSP) translated from mRNA I plays a greater role in refolding proteins than does the HSP translated from mRNA II. Use the data to **support** the hypothesis.
- (d) mRNAs I and II are transcribed from the same gene. **Explain** how a cell can produce two different mRNAs from the same gene.

Write your responses to this question only on the designated pages in the separate Free Response booklet.

STOP

END OF EXAM