

2.1 Testing for biological molecules

Name: _____ Class: _____ Date: _____

Total: 17 marks

KEY WORDS reducing sugar • iodine • hydrochloric acid • protein
• starch

Objective

Build the skills to answer exam questions on **testing for biological molecules** – carrying out and interpreting the food tests, the **semi-quantitative** Benedict's test, and the test for **non-reducing sugars**.

You must be able to:

- name the reagent, method and positive result for the reducing-sugar, starch, lipid and protein tests
- make the Benedict's test semi-quantitative and use it to estimate a concentration
- describe the acid-hydrolysis test for a non-reducing sugar

1 Key knowledge and worked examples


Study these first. Each one shows the **method** for a question type used later.

■ The four food tests

Learn the reagent and the colour change for each test.

test	finds	method	positive result
Benedict's	reducing sugar	add Benedict's solution, heat in a water bath	blue → green → yellow → orange → brick-red precipitate
iodine	starch	add orange-brown iodine solution	turns blue-black
emulsion	lipid	mix with ethanol, then pour into water	white cloudy emulsion
biuret	protein	add biuret solution (no heating)	blue → purple

Benedict's test for sugars



Benedict's test: more reducing sugar gives a colour further along, from blue to brick-red


■ Making Benedict's semi-quantitative

The normal test only says “yes/no”. To estimate **how much** sugar is present, first **standardise** the test on solutions of known **concentration**. Then estimate an unknown by either the **time to the first colour change** (more sugar → faster change) or by matching the final colour to your set of standards.

■ Testing for a non-reducing sugar

Sucrose gives a *negative* Benedict's test. To detect it:

1. Do a normal Benedict's test – it stays blue (no reducing sugar).
2. With a fresh sample, add dilute hydrochloric acid and heat – this **hydrolyses** the sugar into reducing sugars.
3. **Neutralise** the acid with sodium hydrogencarbonate.
4. Repeat the Benedict's test – a brick-red colour now shows a non-reducing sugar was present.



Hydrolyse the sugar with acid first, then the second Benedict's test turns red

2 Practice

2.1 State the reagent and the positive result for each test. [4]

test	reagent	positive result
starch		
protein		

2.2 A student wants to compare the reducing-sugar concentration of three drinks. Describe how the Benedict's test can be made **semi-quantitative** to do this. [3]

3 Exam-style questions

3.1 Which test and result correctly identifies a protein? [1]

- ☐ **A** biuret solution, stays blue
B biuret solution, turns purple
C Benedict's solution, turns brick-red
D iodine solution, turns blue-black

3.2 A solution gives a negative Benedict's test. After boiling it with hydrochloric acid, neutralising, and re-testing, it turns brick-red. The solution contained: [1]

- ☐ **A** a lipid
B a non-reducing sugar
C a protein
D starch

3.3 A student is given an unknown solution X and asked to find out whether it contains reducing sugar, starch and protein.

(a) Describe the tests the student should carry out and the result that would show each substance is present. [4]

(b) The student finds that X contains a non-reducing sugar but no reducing sugar. Describe how the student could have shown this. [3]

(c) State **one** variable the student should keep the same when comparing the colour from different samples. [1]
