


3 A cell in the human body is specialised for synthesis and secretion of lipids.

Which row shows the organelles required by the cell for synthesis **and** secretion of lipids?

	mitochondria	smooth endoplasmic reticulum	vesicles
A	✓	✓	✓
B	✓	x	x
C	x	✓	✓
D	x	✓	x

key
 ✓ = required
 x = **not** required

4 The photomicrograph shows part of a cell.



Which row is correct?

	this cell will contain 70S ribosomes	this could be part of a plant cell	the organelles shown could be used to make antibodies
A	✓	✓	✓
B	✓	x	x
C	x	✓	x
D	x	x	✓


key
 ✓ = correct
 x = **not** correct

2 Which structures will be present in a cell that causes cholera?

- 1 circular DNA
- 2 cytoplasmic DNA
- 3 70S ribosomes

A 1, 2 and 3 B 1 and 2 only C 1 and 3 only D 2 and 3 only

5 The electron micrograph shows a section through a cell.




What would be present in this type of cell?

	mitochondria	peptidoglycan cell wall	histone proteins
A	✓	✓	x
B	✓	x	✓
C	x	✓	x
D	x	x	✓

key
 ✓ = present
 x = **not** present

- 2 Cell structures can be isolated by centrifuging the cells in a sucrose density gradient. This breaks up the cells and separates the cell structures into fractions.


The diagram shows the positions of some of these fractions in a sucrose density gradient.



Which fractions contain cell structures that carry out protein synthesis?

- A 1, 2 and 3 B 1, 3 and 4 C 1 only D 2, 3 and 4

- 7 Which features shown in the diagram can be present in viruses?



- A 1, 2 and 3 B 1, 2 and 4 C 1, 3 and 4 D 2, 3 and 4

- 3 It is thought that some organelles in eukaryotic cells evolved from free-living prokaryotes.

Which organelles have features that suggest they evolved from free-living prokaryotic ancestors?

- A chloroplasts and mitochondria
B chloroplasts and Golgi bodies
C mitochondria and smooth endoplasmic reticulum
D Golgi bodies and smooth endoplasmic reticulum

- 4 A student completed a table showing the structures that are found in four different types of cells.

cell structure	cell Q	cell R	cell S	cell T
cell surface membrane	✓	✓	✓	✓
cellulose cell wall	✓	✓	x	✓
chloroplast	✓	x	x	x
mitochondrion	✓	✓	x	✓
nucleus	✓	✓	x	x

key
✓ = present
x = not present

How many of these types of cells can be found in plants?

- A 1 B 2 C 3 D 4

- Fig. 1.1 is a diagram of a section through a single villus and a crypt of Lieberkühn.



⑤

(a) A student incorrectly stated that an enterocyte has many **cilia** on its apical surface.

..... [2]

- [1]

- (i) Blood vessel **X** receives blood from an artery.

..... [1]

- [2]

Fig. 1.2

1 There are similarities and differences between the structure of a typical plant cell and a typical animal cell.

(a) The nucleus of plant cells and animal cells contains chromosomes. In interphase of the cell cycle, individual chromosomes are present but cannot be seen. The chromosome material is known as chromatin.

(i) Changes occur in interphase, which result in a difference between the chromatin in the G1 phase compared with the chromatin in the G2 phase.

State **and** explain the difference in chromatin in the G1 phase compared with chromatin in the G2 phase.

.....

.....

.....

.....

.....

..... [2]

(ii) Describe the features of a nucleus, **other than** containing chromatin.

You may use the space below the lines for a diagram.

.....

.....

.....

.....

.....

.....

.....

.....

[3]

(b) Starch, cellulose and pectins are polysaccharides found in plant cells but **not** in animal cells.

Pectins are complex polysaccharides that are found in the cell wall.

(i) Describe the structural features of starch that are **different** from the structural features of cellulose.

.....

.....

.....

.....

.....

.....

.....

.....

.....

..... [4]

(ii) Cell wall pectins can vary in different plant cell types and in different stages of cell development. Pectin molecules are released from cells as basic structures and then modified within the cell wall by adding side chains.

RG-I is a pectin molecule with a variable structure. The basic structure is a repeated disaccharide made from two different monosaccharides. RG-I has three different side chains that can be added in different positions.

Monoclonal antibodies (mAbs) are used to investigate the structure, location and role of cell wall pectins.

Suggest **and** explain why scientists need to use a number of different monoclonal antibodies when investigating a pectin such as RG-I.

.....

.....

.....

.....

.....

.....

.....

.....

.....

..... [3]

[Total: 12]