

- 16** A sample of healthy plant cells taken from the same tissue is placed in a beaker containing distilled water.


The cells change in size.

What would explain this change in size?

- A** Water will leave the cells by active transport.
- B** Water will enter the cells by osmosis.
- C** Solutes will enter the cells by active transport.
- D** Solutes will leave the cells by osmosis.

- 17** A student studying surface area to volume ratio and diffusion made a cuboid, S1, using agar stained blue with a pH indicator. The dimensions of S1 are shown in the diagram.

The student made a second agar cuboid, S2. Each dimension of S2, (the length, the width and the height), was half that of S1.



The student placed each cuboid in a test-tube and covered it in acid. The time taken for each cuboid to completely change colour was recorded. All variables other than the size of the cuboids were standardised.

Which row shows the surface area to volume ratio of S1 and the time taken for S1 to change colour completely in acid compared to the time taken for S2 to change colour completely?

	surface area to volume ratio of S1	time taken for S1 to change colour completely in acid compared to S2
A	0.72 : 1	S1 takes less time than S2
B	0.72 : 1	S1 takes more time than S2
C	1.4 : 1	S1 takes less time than S2
D	1.4 : 1	S1 takes more time than S2

- 18** Two test-tubes, labelled X and Y, were set up containing equal volumes of solution X or solution Y respectively.

A large number of type P cells and a large number of type Q cells were added into test-tube X and also into test-tube Y.

After a few minutes, samples of the solutions were taken and the cells were observed with a microscope.

- In solution X, all of cell type P had burst and cell type Q had **not** burst.
- In solution Y, **no** cells had burst.

Which row correctly identifies cell type Q and solutions X and Y?

	cell type Q	solution X	solution Y
A	red blood cells	5% NaCl solution	distilled water
B	goblet cells	5% glucose solution	distilled water
C	liver cells	distilled water	5% glucose solution
D	root hair cells	distilled water	5% NaCl solution

- 19** In an experiment, four different-sized cubes of colourless agar were placed into a blue dye. The time taken for each cube to turn blue was recorded.

Which row shows the surface area and volume of the cube that took the longest time to turn completely blue?

	surface area /cm ²	volume /cm ³
A	6.0	1
B	9.5	2
C	12.5	3
D	38.0	16


19 Which statements about the cell surface membrane are correct?

- 1 Channel proteins allow water soluble ions and molecules across the membrane.
- 2 Glucose can pass into the cell via carrier proteins.
- 3 Oxygen passes freely through the membrane as it is soluble in lipids.
- 4 Some glycoproteins act as antigens.

- A** 1, 2, 3 and 4
B 1, 3 and 4 only
C 1 and 2 only
D 2, 3 and 4 only

20 The diagram shows two cylinders of agar.

The agar cylinders are placed into a solution of dye at the same time. The time taken for the dye to diffuse into the centre of each cylinder is recorded.



Which statement explains why the dye reaches the centre of one agar cylinder faster than the other agar cylinder?

- A** The dye reaches the centre of X in a shorter time than the dye reaches the centre of Y because X has a larger surface area to volume ratio.
B The dye reaches the centre of X in a shorter time than the dye reaches the centre of Y because X has a larger surface area.
C The dye reaches the centre of Y in a shorter time than the dye reaches the centre of X because the dye has a shorter distance to diffuse to the centre.
D The dye reaches the centre of Y in a shorter time than the dye reaches the centre of X because Y has a smaller volume.

17 When red blood cells are put into distilled water they burst (haemolysis).

Which statements explain this haemolysis?

- 1 The water potential of the surrounding liquid is lower than the water potential of the contents of the red blood cell.
- 2 The cell surface membranes of red blood cells are **not** supported by cell walls.
- 3 More water moves into the red blood cells by osmosis than leaves the cells.
- 4 Water enters the red blood cells by osmosis but does **not** leave the cells.

A 1 and 3


B 1 and 4

C 2 and 3


D 2 and 4

18 The diagrams show the shape and size of two types of cell.

palisade mesophyll cell
surface area = $2200\ \mu\text{m}^2$



columnar epithelial cell
volume = $2250\ \mu\text{m}^3$



not to scale

Which statement about the palisade cell and epithelial cell shown in the diagrams is correct?

- A An increase in surface area reduces the distance for gases to reach the centre of the cell.
- B The surface area of the palisade mesophyll cell is $500\ \mu\text{m}^2$ greater than the columnar epithelial cell.
- C The surface area to volume ratio is greater in the columnar epithelial cell than the palisade mesophyll cell.
- D The volume of the palisade mesophyll cell is $2500\ \mu\text{m}^3$ greater than that of the columnar epithelial cell.

1 Many substances can move through cell surface membranes between the cytoplasm of animal cells and the extracellular environment.

(a) A student made a drawing to summarise the movement of substances across the cell surface membranes of mammalian red blood cells.

Fig. 1.1 shows the drawing made by the student:

- Each arrow indicates the movement of a substance through the membrane.
- The number of each of the 4 shapes represents the relative concentrations of each substance in the cytoplasm and in the blood plasma.




Fig. 1.1

The student carried out research and made a list of some of the substances found in red blood cells as shown in Fig. 1.2.




Fig. 1.2

Table 1.1 shows information about the 4 types of movement of substances across the cell surface membranes of red blood cells as shown in Fig. 1.1.

Complete Table 1.1 using the information in Fig. 1.1 and Fig. 1.2.

Table 1.1

letter from Fig. 1.1	type of movement	name of part of membrane involved	example of a substance that moves across the membrane (from Fig. 1.2)
A	simple diffusion	phospholipids	
B	facilitated diffusion		calcium ions
C	facilitated diffusion		
D			

[5]

(b) Some viruses infect plants through the surfaces of damaged leaves. These plant viruses can travel from one leaf cell to another without having to pass through any cell surface membranes.

Explain how some plant viruses can travel from one cell to another without passing through cell surface membranes.

[2]

[Total: 7]

1 *Beta vulgaris vulgaris*, as shown in Fig. 1.1, is a food crop with a swollen, edible root known as beetroot.




Fig. 1.1

A student determined the concentration of a sucrose solution that has the same water potential as the cells of the beetroot.

The student used distilled water and a stock solution of 2.0 mol dm⁻³ sucrose to prepare a range of sucrose solutions.

Each sucrose solution was prepared using proportional dilution and had a final volume of 30 cm³.

(a) Complete Table 1.1 to show how the concentrations were made.

Table 1.1

final concentration of sucrose solution /mol dm ⁻³	volume of distilled water /cm ³	volume of 2.0 mol dm ⁻³ stock sucrose solution /cm ³
2.0	0.0	30.0
.....		
.....		
.....		
.....		
0.0	30.0 (8)	0.0

The student was supplied with the sucrose solutions prepared in **1(a)** and standard laboratory equipment.

- prepared the sucrose solutions in **1(a)**
- would use the data collected to determine the concentration of a sucrose solution that has the same water potential as the cells of the beetroot.

Your method should be set out in a logical order and be detailed enough to allow another person to follow it.

This image shows a full page of white paper with horizontal dashed lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the paper.

(c) The student calculated the mean percentage change in mass of the beetroot for each concentration of sucrose solution.

(i) The student plotted these calculated values to obtain a line graph.

Complete Fig. 1.2 by:

- labelling the axes
- sketching the line that the student obtained.




Fig. 1.2

[2]

(ii) Explain how the student used the completed line graph in Fig. 1.2 to estimate the concentration of sucrose solution that has the same water potential as the cells of the beetroot.

.....
.....
..... [1]

(iii) The cells of the beetroot contain a red pigment which can leak out into the solution when the tissue is cut. The student noticed that each sucrose solution was coloured red at the end of the investigation.

Suggest **one** improvement the student could make to the method to reduce the amount of red pigment in each sucrose solution.

.....
.....
..... [1]

[Total: 13]

Question 2 starts on page 6.