

Starter Quiz

A-Level Chemistry

Name: _____ Score: _____

1. A functional group is:

- ☐ the carbon skeleton only
- ☐ the molecule's mass
- ☐ the reactive part of a molecule that decides its properties
- ☐ the number of atoms

2. Homolytic fission of a covalent bond produces:

- ☐ two ions
- ☐ no products
- ☐ two free radicals (one electron to each atom)
- ☐ one molecule

3. In an addition reaction:

- ☐ a molecule is split by water
- ☐ a group is swapped
- ☐ two molecules join to make one
- ☐ a small molecule is lost

4. An sp^3 carbon (four single bonds) is:

- ☐ tetrahedral, 109.5°
- ☐ planar, 120°
- ☐ linear, 180°
- ☐ bent, 104.5°

5. Isomers are compounds with:

- ☐ no functional group
- ☐ different molecular formulas
- ☐ the same molecular formula but a different arrangement of atoms
- ☐ the same arrangement of atoms

6. E/Z (cis-trans) isomerism arises because there is no rotation about a C=C double bond.

- ☐ True ☐ False

8. Breaking a bond so each atom keeps one electron is _____ fission.

9. A carbon atom forming four single bonds is trigonal planar with bond angles of 120° .

- ☐ True ☐ False

10. Compounds with the same molecular formula but different structures are structural _____.

1. **the reactive part of a molecule that decides its properties**

Molecules with the same functional group behave alike, because the group is the reactive part.

2. **two free radicals (one electron to each atom)**

Homolytic = even split, one electron each $\rightarrow$ two radicals; heterolytic = uneven $\rightarrow$ two ions.

3. **two molecules join to make one**

Addition combines two molecules into a single product (e.g. across a C=C bond).

4. **tetrahedral, 109.5°**

Four single bonds (sp^3) give a tetrahedral shape at 109.5°.

5. **the same molecular formula but a different arrangement of atoms**

Isomers share a molecular formula but differ in how the atoms are arranged.

6. **True**

The restricted rotation fixes the groups' positions.

7. **functional**

e.g. -OH, -COOH.

8. **homolytic**

It forms free radicals.

9. **False**

False. Four bonding pairs repel equally and point to the corners of a tetrahedron, giving bond angles of about 109.5°. Trigonal planar (120°) needs three electron pairs.

10. **isomers**

Chain, position and functional-group isomerism.