

Revision Sheet

A-Level Chemistry · Unit 28 Chemistry of transition elements

6 vocabulary items · 59 marks of exam questions (4 questions)

Part 1 · Vocabulary review answer key. Section A wants the exact English spelling.

Question	Answer	Marks
1	ligand	1
2	transition element	1
3	oxidation state	1
4	complex ion	1
5	monodentate	1
6	complex	1


7(a)	<i>Any two from:</i> <ul style="list-style-type: none">• they have variable / multiple oxidation states OWTTE• they behave as catalysts• they form complex ions / complexes• they form coloured compounds/ions																			
7(b)(i)	(is a molecule or ion formed by a central) metal atom / metal ion bonded / surrounded by (one or more) ligands																			
7(b)(ii)	<table><tr><th>reagent added</th><th>formula of species formed</th><th>colour / state of species formed</th><th>type of reaction</th></tr><tr><td>an excess of NH₃(aq)</td><td>[Co(NH₃)₆]²⁺</td><td>brown solution ALLOW <i>yellow-brown</i> solution</td><td>ligand exchange</td></tr><tr><td>an excess of concentrated HCl</td><td>[CoCl₄]²⁻</td><td>blue solution</td><td>ligand exchange</td></tr><tr><td>NaOH(aq)</td><td>Co(OH)₂ OR Co(OH)₂(H₂O)₄</td><td>blue ppt. ALLOW pink ppt</td><td>precipitation</td></tr></table> <p>formula and colour / state: any two [1], any four [2], all six [3] type of reaction: all three correct [1]</p>				reagent added	formula of species formed	colour / state of species formed	type of reaction	an excess of NH ₃ (aq)	[Co(NH ₃) ₆] ²⁺	brown solution ALLOW <i>yellow-brown</i> solution	ligand exchange	an excess of concentrated HCl	[CoCl ₄] ²⁻	blue solution	ligand exchange	NaOH(aq)	Co(OH) ₂ OR Co(OH) ₂ (H ₂ O) ₄	blue ppt. ALLOW pink ppt	precipitation
reagent added	formula of species formed	colour / state of species formed	type of reaction																	
an excess of NH ₃ (aq)	[Co(NH ₃) ₆] ²⁺	brown solution ALLOW <i>yellow-brown</i> solution	ligand exchange																	
an excess of concentrated HCl	[CoCl ₄] ²⁻	blue solution	ligand exchange																	
NaOH(aq)	Co(OH) ₂ OR Co(OH) ₂ (H ₂ O) ₄	blue ppt. ALLOW pink ppt	precipitation																	
7(c)(i)	M1 (a species) that donates / uses two lone pairs M2 to form a two dative / coordinate bond to a metal atom / metal ion																			

7(c)(ii)	<p>Any four from:</p> <div style="display: flex; flex-wrap: wrap;"> <div style="width: 50%; text-align: center;"> <p>isomer 1</p> </div> <div style="width: 50%; text-align: center;"> <p>isomer 2</p> </div> <div style="width: 50%; text-align: center;"> <p>isomer 3</p> </div> <div style="width: 50%; text-align: center;"> <p>isomer 4</p> </div> <div style="width: 50%; text-align: center;"> <p>isomer 5</p> </div> <div style="width: 50%; text-align: center;"> <p>isomer 6</p> </div> </div> <p>Any two [1], any three [2], all four [3]</p>
7(c)(iii)	<p>oxidation state of cobalt (+)3 AND type of stereoisomerism cis-trans OR geometric(al) OR optical</p>

8(a)(i)	donates one lone pair / forms one dative bond AND to metal atom / to metal ion [1]
8(a)(ii)	<p>Mo(CO)₆ 0 / no charge / neutral / zero Fe(CN)₆ 3–</p> <p>both formulae [1] both charges [1]</p>
8(a)(iii)	<p>either of these renditions, with a CN on each bond:</p> <div style="display: flex; justify-content: space-around; align-items: center;"> </div> <p>AND</p> <p>180° bond angle between two opposite bonds [1]</p>
8(b)(i)	<p>[Cu(H₂O)₆]²⁺ + 4NH₃ → [Cu(NH₃)₄(H₂O)₂]²⁺ + 4H₂O</p> <p>OR</p> <p>Cu²⁺ + 2H₂O + 4NH₃ → [Cu(NH₃)₄(H₂O)₂]²⁺ [1]</p>
8(b)(ii)	<p>M1 d-orbitals split (could be in diagram) [1]</p> <p>M2 electrons transition to higher orbital / electrons promoted or excited [1]</p> <p>M3 wavelength / frequency / light / photon / hf / hn absorbed [1]</p> <p>M4 colour seen / reflected / transmitted is complement of colour absorbed [1]</p>

8(c)(i)	<p>[Cu(NH₃)₄(H₂O)₂]²⁺ + 4HCl → CuCl₄²⁻ + 2H₂O + 4NH₄⁺</p> <p>OR</p> <p>[Cu(NH₃)₄(H₂O)₂]²⁺ + 4Cl⁻ (+ 4H⁺) → CuCl₄²⁻ + 2H₂O + 4NH₃ (+ 4H⁺)</p> <p>M1 formula and charge of CuCl₄²⁻ [1]</p> <p>M2 rest of equation [1]</p>
8(c)(ii)	ligand exchange / substitution / replacement / displacement [1]
8(c)(iii)	tetrahedral [1]
8(c)(iv)	yellow [1]
8(c)(v)	d -orbital splitting changes / ΔE changes [1]

9(a)	a d-block element that forms one or more stable ions with incomplete d-orbitals		
9(b)(i)		geometry	CN
	[Cu(H ₂ O) ₆] ²⁺	octahedral	6
	[CuCl ₄] ²⁻	tetrahedral	4
9(b)(ii)	<p>degenerate – of the same energy</p> <p>non-degenerate – not of the same energy / different energy</p>		
9(b)(iii)	different frequency / wavelength / photon absorbed		
9(c)(i)	forms 2 dative / coordinate bonds		
9(c)(ii)	<p>one isomer correct with correct 3D [1]</p> <p>three isomers different with correct 3D [1]</p>		
9(c)(iii)	<p>e.g. 1 and 2 optical [1]</p> <p>e.g. 1 or 2 and 3 cis-trans / geometric [1]</p>		
9(d)	<p>0.0230 moles ethanedioate in whole 100 cm³</p> <p>0.00575 moles ethanedioate in 25 cm³ sample [1]</p> <p>0.00230 moles of MnO₄⁻ needed [1]</p> <p>230 cm³ [1]</p>		
9(e)	<p>Cu²⁺ is smaller / has a smaller radius / than Ba²⁺ [1]</p> <p>Cu²⁺ polarises / distorts the anion more [1]</p>		

10(a)	M1 ΔH_{latt} and ΔH_{hyd} both become less exothermic / less negative M2 ΔH_{hyd} changes less / becomes less exothermic by a smaller extent OR ΔH_{latt} changes more / dominant factor / changes faster M3 ΔH_{sol} becomes more exothermic / more negative OR ΔH_{sol} becomes less endothermic / less positive				
10(b)(i)	M1 $K_{\text{sp}} = [\text{Be}^{2+}][\text{OH}^-]^2$ M2 $\text{mol}^3 \text{dm}^{-9}$ ecf				
10(b)(ii)	M1 solubility of $\text{Be}(\text{OH})_2 = 2.40 \times 10^{-6} / 43 = 5.58 \times 10^{-8} (\text{mol dm}^{-3})$ ecf (b)(i) M2 $K_{\text{sp}} = 4 \times (5.58 \times 10^{-8})^3 = 6.95 \times 10^{-22}$ ecf min 2sf				
10(c)(i)	as a molecule or ion formed by a central metal atom / metal ion dative bonded to / surrounded by one or more ligands				
10(c)(ii)	 tetrahedral and correct 3D structure				
10(d)(i)	d orbitals that are energetically accessible OR empty / vacant d orbitals AND form dative bonds / accept a lone pair from a ligand				
10(d)(ii)	complex ion	shape	coordination number	polarity	
	cis-[Pt(en)Cl ₂]	square planar	4	polar	
	[Ag(NH ₃) ₂] ⁺	linear	2	non-polar	
	[Fe(C ₂ O ₄) ₃] ³⁻	octahedral	6	non-polar	
three for one mark, six for two marks					
10(e)(i)	the equilibrium constant for the formation of the complex ion in a solvent (from its constituent ions or molecules)				
10(e)(ii)	M1 $K_{\text{stab}} = [\text{Ni}(\text{en})_3^{2+}] / [\text{Ni}(\text{H}_2\text{O})_6^{2+}][\text{en}]^3$ M1 units = $\text{mol}^{-3} \text{dm}^9$ ecf				
10(e)(iii)	the equilibria lie to the right / product side OR equilibrium 1 lies more to the right AND <i>en</i> complex is more stable / $[\text{Ni}(\text{en})_3]^{2+}$ is more stable				