

Solutions

Each answer shows the steps, then the **result**.

2.1

- the nitrogen lone pair can accept (donate to) an H^+

2.2

- excess ammonia, NH_3 , in ethanol; heat under pressure

3.1 C

- Ethylamine — the alkyl group makes the lone pair most available

3.2 A

- LiAlH_4 reduces the nitrile to a primary amine, ethylamine

3.3

- the ethyl group pushes electron density onto the nitrogen (positive inductive effect); this makes the lone pair more available to accept an H^+ , so ethylamine is a stronger base than ammonia

3.4

- in phenylamine the nitrogen lone pair overlaps with / is delocalised into the benzene ring; so it is less available to accept an H^+ ; whereas in ethylamine the lone pair is fully available (and the alkyl group pushes electrons onto N), so phenylamine is much weaker

3.5

(a)

- phenylamine < ammonia < ethylamine. A base donates the nitrogen lone pair, so the more available that pair, the stronger the base. In ethylamine the ethyl group **releases** electron density inductively towards the nitrogen, making the lone pair more available; in phenylamine the lone pair is delocalised into the benzene ring, so it is much less available

(b)

- $\text{CH}_3\text{CH}_2\text{Br} + 2\text{NH}_3 \rightarrow \text{CH}_3\text{CH}_2\text{NH}_2 + \text{NH}_4\text{Br}$; the mechanism is **nucleophilic substitution**; an excess of ammonia makes it more likely that ammonia — not the ethylamine already made — attacks the next bromoethane, which limits further substitution to secondary and tertiary amines

(c)

- Tin and concentrated hydrochloric acid, heated under reflux, then made alkaline with sodium hydroxide; $\text{C}_6\text{H}_5\text{NO}_2 + 6[\text{H}] \rightarrow \text{C}_6\text{H}_5\text{NH}_2 + 2\text{H}_2\text{O}$

(d)

- $\text{CH}_3\text{CH}_2\text{NH}_2 + \text{HCl} \rightarrow \text{CH}_3\text{CH}_2\text{NH}_3^+\text{Cl}^-$; **ethylammonium chloride**