

- 20** Equal masses of CaCO_3 , $\text{Ca}(\text{NO}_3)_2$, BaCO_3 and $\text{Ba}(\text{NO}_3)_2$ are thermally decomposed. The volume of gas produced in each experiment is measured under the same conditions.

Which compound will produce the greatest volume of gas?

- A** CaCO_3 **B** $\text{Ca}(\text{NO}_3)_2$ **C** BaCO_3 **D** $\text{Ba}(\text{NO}_3)_2$

- 19** When heated, magnesium nitrate decomposes.

Which equation for the thermal decomposition of magnesium nitrate is correct?

- A** $\text{Mg}(\text{NO}_3)_2 \rightarrow \text{MgO} + \text{NO}_2 + \text{NO} + \text{O}_2$
B $2\text{Mg}(\text{NO}_3)_2 \rightarrow 2\text{MgO} + 4\text{NO} + 3\text{O}_2$
C $2\text{Mg}(\text{NO}_3)_2 \rightarrow 2\text{MgO} + 4\text{NO}_2 + \text{O}_2$
D $3\text{Mg}(\text{NO}_3)_2 \rightarrow \text{Mg}_2\text{N}_3 + \text{MgO} + 3\text{NO} + 7\text{O}_2$

- 20** R is the aqueous solution of an ionic compound.

- A white precipitate is formed when R is added to $\text{Sr}(\text{NO}_3)_2(\text{aq})$.
- No visible reaction is seen when R is added to dilute $\text{HNO}_3(\text{aq})$.

What is the anion present in compound R?

- A** Cl^- **B** SO_4^{2-} **C** HCO_3^- **D** CO_3^{2-}

- 19** Magnesium nitrate, $\text{Mg}(\text{NO}_3)_2$, will decompose when heated to give a white solid and a mixture of gases. One of the gases released is oxygen.

29.7 g of anhydrous magnesium nitrate is heated until no further reaction takes place.

Which mass of oxygen is produced?

- A** 3.2 g **B** 6.4 g **C** 12.8 g **D** 19.2 g