

1 (a) Solutions of Group 2 hydrogencarbonates, $M(\text{HCO}_3)_2$, decompose on heating to give the corresponding metal carbonate, carbon dioxide and water.

(i) Write an equation for the decomposition of strontium hydrogencarbonate, $\text{Sr}(\text{HCO}_3)_2$.
..... [1]

(ii) The thermal stability of Group 2 carbonates increases down the group.

Explain this trend.

.....
.....
.....
..... [2]

(b) The hydroxides and fluorides of Group 2 elements show similar trends in solubility.

Describe the trend in the solubility of the fluorides of calcium, strontium and barium.

Explain your answer.

.....
least soluble most soluble
explanation
.....
.....
.....
.....
..... [4]

(c) (i) Define enthalpy change of hydration, ΔH_{hyd} .

.....
.....
..... [1]

(ii) State the main factors that affect the magnitude of enthalpy change of hydration.

Explain your answer.

.....
.....
.....
..... [2]

(d) Table 1.1 shows various energy changes.

Table 1.1

energy change	value / kJ mol^{-1}
lattice energy of MgF_2	–2957
enthalpy change of hydration, ΔH_{hyd} , of Mg^{2+}	–1926
enthalpy change of hydration, ΔH_{hyd} , of F^-	–505

Use data from Table 1.1 to calculate the enthalpy change of solution, ΔH_{sol} , for $\text{MgF}_2(\text{s})$.

It may be helpful to draw a labelled energy cycle. Show your working.

ΔH_{sol} of $\text{MgF}_2(\text{s}) = \dots\dots\dots \text{kJ mol}^{-1}$ [2]

- (e) Mercury(I) fluoride, Hg_2F_2 , is sparingly soluble in water.

The cation in Hg_2F_2 exists as the diatomic ion Hg_2^{2+} with a covalent Hg–Hg bond.

- (i) Write the expression for the solubility product, K_{sp} , of Hg_2F_2 . Include the units.

$$K_{\text{sp}} =$$

units

[2]

- (ii) The solubility of Hg_2F_2 is $9.20 \times 10^{-3} \text{ mol dm}^{-3}$ at 298 K.

Calculate the value of K_{sp} of Hg_2F_2 at 298 K.

$$K_{\text{sp}} = \dots\dots\dots [1]$$

[Total: 15]