

1(a)(i)	$\text{Sr}(\text{HCO}_3)_2 \rightarrow \text{SrCO}_3 + \text{CO}_2 + \text{H}_2\text{O}$
1(a)(ii)	M1: radius of cation / M^{2+} increases OR charge density of cation / M^{2+} decreases M2: less polarisation / less distortion of the anion / carbonate ion
1(b)	M1: least soluble $\text{CaF}_2 < \text{SrF}_2 < \text{BaF}_2$ most soluble M2: ΔH_{latt} and ΔH_{hyd} both become less exothermic / less negative M3: ΔH_{hyd} changes less / becomes less exothermic by a smaller extent OR ΔH_{latt} changes more / becomes less exothermic by a larger extent M4: ΔH_{sol} becomes more exothermic / more negative
1(c)(i)	enthalpy change when one mole of gaseous ions dissolve in water to form a solution
1(c)(ii)	M1: ionic radii AND ionic charge M2: (ionic) radii increase / charge density decreases AND ΔH_{hyd} decreases/less exothermic AND less attraction between water molecules and (gaseous) ions OR as ionic charge increases / charge density increases AND ΔH_{hyd} increases/more exothermic AND more attraction between water molecules and (gaseous) ions
1(d)	M1: use of $-2957 / -1926 / -505$ AND $2 \times (-505)$ M2: correct signs and evaluation ΔH_{sol} of $\text{MgF}_2(\text{s}) = -1926 + (2 \times -505) - (-2957) = (+)21 \text{ kJ mol}^{-1}$
1(e)(i)	M1: $K_{\text{sp}} = [\text{Hg}_2^{2+}] [\text{F}^-]^2$ M2: units = $\text{mol}^3 \text{ dm}^{-9}$
1(e)(ii)	$K_{\text{sp}} = 2^2 \times (9.20 \times 10^{-3})^3 = 3.11 \times 10^{-6} \text{ min 2sf}$