


6 (a) Compound **Z** is used in organic synthesis.



Complete Table 6.1 to show the number of sp , sp^2 and sp^3 hybridised carbon atoms present in one molecule of **Z**.

Table 6.1

type of hybridisation	sp	sp^2	sp^3
number of carbon atoms			

[1]

(b) **Z** can undergo different reactions, as shown in Fig. 6.1.




Fig. 6.1

(i) Name the **two** types of reaction occurring in reaction 7 in Fig. 6.1.

..... and [1]

(ii) Draw the structures of the organic products of reactions 7, 8 and 9 in Fig. 6.1. [4]

(1)

(c) Compound **Z** is dissolved in D_2O and analysed by carbon-13 NMR and proton (1H) NMR spectroscopy.

(i) Predict the number of peaks in the carbon-13 NMR spectrum of **Z**.

..... [1]

(ii) The proton (1H) NMR spectrum of **Z** in D_2O gives three peaks for the proton environments, labelled **a**, **b** and **c**, as shown on Fig. 6.2.




Fig. 6.2

Complete Table 6.2 for the proton (1H) NMR spectrum of **Z** in D_2O .

Table 6.2

proton environment	a	b	c
name of splitting pattern			
chemical shift range, δ /ppm			

[2]

Table 6.3

environment of proton	example	chemical shift range, δ /ppm
alkane	$-CH_3$, $-CH_2-$, $>CH-$	0.9–1.7
alkyl next to $C=O$	$CH_3-C=O$, $-CH_2-C=O$, $>CH-C=O$	2.2–3.0
alkyl next to nitrile	$-CH_2-CN$	2.0–3.0
alkyl next to electronegative atom	CH_3-O , $-CH_2-O$, $-CH_2-N$	3.2–4.0
attached to alkene	$=CHR$	4.5–6.0
alkyl amine	$R-NH-$	1.0–5.0
amide	$RCONHR$	5.0–12.0

(2)

[Total: 9]