

Week 45

A-Level Chemistry

Homework bundle for this week

本周作业合集

exercises.lol

Contents 目录

Topic 30 quiz	2
Exercise sheet 30.1	5
Past-paper questions 30.1	9

A-Level Chemistry

Name: _____ Score: _____

1. Benzene reacts mostly by electrophilic substitution because:
 - ☐ this keeps the stable delocalised ring, while addition would destroy it
 - ☐ the ring is unstable
 - ☐ it has no electrons
 - ☐ addition is impossible
2. A halogen reacts in the ring (not the side-chain) when there is:
 - ☐ a halogen-carrier catalyst (AlCl_3) and no UV light
 - ☐ UV light and no catalyst
 - ☐ only heat
 - ☐ water present
3. Nitration of benzene uses:
 - ☐ concentrated HNO_3 and concentrated H_2SO_4
 - ☐ bromine water
 - ☐ dilute NaOH
 - ☐ hydrogen and a catalyst
4. Under UV light with no catalyst, the halogen substitutes:
 - ☐ in the side-chain (free-radical substitution)
 - ☐ in the ring
 - ☐ nowhere
 - ☐ at the catalyst
5. A Friedel–Crafts reaction (with AlCl_3) is used to:
 - ☐ add an alkyl or acyl group to the ring
 - ☐ remove a hydrogen as a radical
 - ☐ oxidise the ring
 - ☐ hydrogenate the ring
6. Groups already on a benzene ring direct where the next group attaches.
 - ☐ True ☐ False

7. Benzene reacts mainly by electrophilic substitution, not addition, because that keeps its stable delocalised ring intact.
- ☐ True ☐ False
8. An -OH group already on the ring directs the next substituent to:
- ☐ positions 2 and 4
- ☐ position 3 only
- ☐ no particular position
- ☐ the side-chain
9. In electrophilic substitution, after the ring bonds to the electrophile:
- ☐ an H^+ is lost, restoring the stable ring
- ☐ the ring opens up
- ☐ another electrophile adds
- ☐ a radical forms
10. Arenes such as benzene typically react by:
- ☐ electrophilic substitution
- ☐ electrophilic addition
- ☐ nucleophilic substitution
- ☐ free-radical addition

A-Level Chemistry

1. **this keeps the stable delocalised ring, while addition would destroy it**

Substitution preserves the aromatic stabilisation; addition would break up the delocalised system.

2. **a halogen-carrier catalyst (AlCl_3) and no UV light**

The catalyst makes the electrophile for ring substitution; UV with no catalyst gives side-chain free-radical substitution.

3. **concentrated HNO_3 and concentrated H_2SO_4**

The conc. acid mixture generates the NO_2^+ electrophile, giving nitrobenzene.

4. **in the side-chain (free-radical substitution)**

UV light starts free-radical substitution in the alkyl side-chain.

5. **add an alkyl or acyl group to the ring**

Friedel-Crafts alkylation/acylation adds a carbon-based group to benzene.

6. **True**

This is the directing effect.

7. **True**

Addition would break the delocalised ring and lose its stability, so substitution (swapping an H) is favoured.

8. **positions 2 and 4**

$-\text{NH}_2$, $-\text{OH}$ and $-\text{R}$ are 2,4-directing; $-\text{NO}_2$, $-\text{COOH}$ and $-\text{COR}$ are 3-directing.

9. **an H^+ is lost, restoring the stable ring**

Losing H^+ from that carbon restores the delocalised ring, completing the substitution.

10. **electrophilic substitution**

Substitution keeps the stable ring intact.

30.1 Arenes

Name: _____ Class: _____ Date: _____

Total: 11 marks

KEY WORDS electrophilic substitution 亲电取代 • arene 芳烃 • nitration 硝化 • friedel-crafts acylation 傅克酰基化
• benzene 苯

Objective


Build the skills to answer exam questions on **arenes** 芳烃 — the reactions of benzene and methylbenzene, the **electrophilic substitution** 亲电取代 mechanism, side-chain vs ring substitution, and **directing effects** 定位效应.

You must be able to:

- give the reagents, conditions and products for benzene's reactions (halogenation, nitration, Friedel–Crafts, oxidation, hydrogenation)
- describe the electrophilic substitution mechanism (e.g. nitration)
- explain why benzene substitutes rather than adds
- predict where a second group goes from directing effects


1 Worked examples

■ Reactions and mechanism



Benzene keeps its stable ring, reacting mostly by electrophilic substitution (addition only with H_2)

- **nitration:** conc. HNO_3 + conc. H_2SO_4 , 25–60 °C → nitrobenzene.
- **Friedel–Crafts:** RCl (or $RCOCl$) + $AlCl_3$, heat → alkyl/acyl benzene.
- **side-chain oxidation:** hot alkaline $KMnO_4$ then acid → benzoic acid.



Electrophilic substitution: the ring attacks the electrophile, then loses H^+ to restore the stable ring

Why substitution, not addition? Substitution keeps the stable delocalised ring; addition would destroy it.

2 Practice

2.1 Give the reagents and conditions for the **nitration** of benzene. [2]

2.2 Name the electrophile formed in the nitration of benzene. [1]

3 Exam-style questions

3.1 Benzene reacts with CH_3COCl and AlCl_3 . This is an example of: [1]

- **A** Friedel–Crafts acylation
- **B** nitration
- **C** side-chain oxidation
- **D** hydrogenation

3.2 Methylbenzene is treated with Cl_2 in **UV light with no catalyst**. Substitution occurs at the: [1]

- **A** ring, at positions 2 and 4
- **B** ring, at position 3
- **C** side-chain ($-\text{CH}_3$)
- **D** both ring and side-chain equally

3.3 Describe the mechanism of the nitration of benzene. Include the formation of the electrophile and use curly arrows in words. [4]

3.4 Nitrobenzene already carries a $-\text{NO}_2$ group. Predict and explain the position a second nitro group will take. [2]

Solutions

2.1 concentrated nitric acid and concentrated sulfuric acid; 25–60 °C / warm.

2.2 NO_2^+ (the nitryl/nitronium cation).

3.1 A. $\text{RCOCl} + \text{AlCl}_3$ adds an acyl group — Friedel–Crafts acylation.

3.2 C. UV light with no catalyst gives free-radical substitution in the side-chain.

3.3 the acid mixture makes the electrophile NO_2^+ ; the delocalised ring electrons form a bond to NO_2^+ , giving a positively charged intermediate; the C–H bond breaks and H^+ is lost; this restores the delocalised ring, giving nitrobenzene.

3.4 $-\text{NO}_2$ directs the next group to **position 3**; because $-\text{NO}_2$ is an electron-withdrawing (deactivating) group that directs to the 3-position.

8 (a) In the electrophilic substitution of arenes, different substituents can direct to different ring positions.

(i) Describe the directing effect of the $-\text{NO}_2$ group. Explain your answer.

.....
..... [1]

(ii) The nitration of arenes uses a mixture of concentrated HNO_3 and concentrated H_2SO_4 to generate the NO_2^+ electrophile.

Write an equation for the formation of the NO_2^+ electrophile.

..... [1]

(b) Carbon-carbon bond formation is an important reaction in organic synthesis.

Fig. 8.1 shows the synthesis of compound **Q** from benzene in two reaction steps.




Fig. 8.1

(i) Draw the structure of compound **P** in the box in Fig. 8.1.

[1]

(ii) Suggest reagents and conditions for reactions 1 and 2 in Fig. 8.1.

reaction 1

reaction 2

[2]

(c) Separate samples of $\text{C}_6\text{H}_5\text{Br}$ and $\text{C}_6\text{H}_5\text{CH}_2\text{Br}$ are added to warm $\text{AgNO}_3(\text{aq})$.

State the expected observations, if any. Explain your answer.

$\text{C}_6\text{H}_5\text{Br}$ with $\text{AgNO}_3(\text{aq})$

$\text{C}_6\text{H}_5\text{CH}_2\text{Br}$ with $\text{AgNO}_3(\text{aq})$

explanation

.....

.....

.....

[3]

(d) Acyl bromides, RCOBr , react readily with H_2O .

The mechanism of this reaction is similar to that of the reaction of H_2O with acyl chlorides, RCOCl .

(i) Name the mechanism of this reaction.

..... [1]

(ii) Complete the mechanism in Fig. 8.2 for the reaction of RCOBr with H_2O .

Include all relevant lone pairs of electrons, curly arrows, charges and dipoles.

Draw the structure of the intermediate.




Fig. 8.2

[4]

[Total: 13]

9 Compound **X** is made from benzene by the route shown in Fig. 9.1.




Fig. 9.1

(a) Describe the bonding in benzene, C_6H_6 .

Your answer should include:

- the hybridisation of the six carbon atoms
- the types of bond between the carbon atoms
- the orbitals that overlap to produce the bonds between the carbon atoms
- the type of bond between the carbon atoms and the hydrogen atoms
- the orbitals that overlap to produce the bonds between the carbon atoms and the hydrogen atoms.

.....

.....

.....

.....

.....

.....

..... [3]

(b) Describe the reagents and conditions required for step 1 in Fig. 9.1.

.....

..... [1]

(c) In step 1 of Fig. 9.1 benzene reacts with $^+\text{CH}_3$.

Complete Fig. 9.2 to show the mechanism for this reaction, including:

- the movement of electron pairs using curly arrows
- the structure of the intermediate involved.




Fig. 9.2

[3]

(d) Describe the reagents and conditions required for step 2 of Fig. 9.1.

.....
..... [2]

(e) Identify the reagents required for step 3 of Fig. 9.1. Compound **W** is the product of this step.

..... [1]

(f) Name compound **W**.

..... [1]


(g) The reagents commonly used for step 4 will **not** reduce the $-\text{COOH}$ group.

Identify the reagents and conditions required for step 4 of Fig. 9.1.

..... [1]

(h) Benzene can also be used as a starting material to make compound **Y**.

compound **Y**



Describe how the route described in Fig. 9.1 (repeated below) can be changed to give compound **Y** instead of compound **X**.

Explain your answer.

.....
.....
.....
..... [2]




Fig. 9.1

[Total: 14]

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.02 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)

5 (a) Describe and explain the shape of benzene.

In your answer, include:

- the shape and bond angle in the ring
- the hybridisation of the carbon atoms
- how orbital overlap forms σ and π bonds between the carbon atoms in the ring.

.....

.....

.....

.....

.....

.....

.....

.....

..... [4]


(b) Fig. 5.1 shows two reactions of benzoic acid.

Fig. 5.1

(i) Suggest reagents and conditions for reaction 5 and for reaction 6 in Fig. 5.1.

reaction 5

reaction 6

[2]

(ii) State the type of reaction for reaction 5 in Fig. 5.1.

..... [1]

(c) In the electrophilic substitution of arenes, different substituents can direct to different ring positions.

(i) Describe the directing effect of the $-\text{CH}_2\text{CH}_3$ group.

Explain your answer.

.....
..... [1]

(ii) The alkylation of arenes uses a mixture of $\text{CH}_3\text{CH}_2\text{Br}$ and FeBr_3 to generate the CH_3CH_2^+ electrophile.

Write an equation for the formation of the CH_3CH_2^+ electrophile.

..... [1]

(iii) Complete the mechanism in Fig. 5.2.

Include all relevant curly arrows and charges. Draw the structure of the organic intermediate.




Fig. 5.2

[3]

(iv) Write an equation to show how FeBr_3 is regenerated after the reaction in Fig. 5.2.

..... [1]

[Total: 13]