

1 What is the electronic configuration for the particle ${}_{13}^{27}\text{Al}^{+}$?

- A $1s^2 2s^2 2p^6 3s^1 3p^1$
B $1s^2 2s^2 2p^6 3s^2$
C $1s^2 2s^2 2p^6 3s^2 3p^2$
D $1s^2 2s^2 2p^6 3s^2 3p^6 3d^7 4s^1$



7 When 1.0 mol of ethanoic acid, CH_3COOH , in aqueous solution is neutralised by an excess of aqueous sodium hydroxide, 55 kJ mol^{-1} of energy is released.

Which statement about this reaction is correct?

- A The reaction is exothermic because only bond breaking takes place.
B The reaction is exothermic because only bond forming takes place.
C The reaction is exothermic because more energy is given out in breaking bonds than is taken in to form bonds.
D The reaction is exothermic because more energy is given out in forming bonds than is taken in to break bonds.

9 Sulfite ions, SO_3^{2-} , react separately with zinc and with manganese dioxide.

a , b , c , d , w , x , y and z are all whole numbers.



Which numbers are correct for a , b , w and x ?


	a	b	w	x
A	1	2	2	4
B	2	2	2	4
C	1	2	4	2
D	2	2	4	2

11 1.00 g of nitrogen gas is stored in a 2.00 dm^3 vessel at 40.0°C .

What is the pressure in the vessel?

- A 5940 Pa B 11900 Pa C 46400 Pa D 92900 Pa

12 One molecule of haemoglobin, Hb, can bind with four molecules of oxygen according to the equation shown.



When the equilibrium concentration of O_2 is $7.6 \times 10^{-6} \text{ mol dm}^{-3}$, the equilibrium concentrations of Hb and $\text{Hb}(\text{O}_2)_4$ are equal.

What is the numerical value of K_c for this equilibrium?

- A 3.0×10^{20} B 1.3×10^5 C 7.6×10^{-6} D 3.3×10^{-21}

15 The reversible reaction between methanol and ethanoic acid liquids is catalysed by adding a small volume of concentrated sulfuric acid.

Two statements about this reaction are listed.

- 1 The sulfuric acid is a homogeneous catalyst.
2 The sulfuric acid lowers the activation energy of the reverse reaction.

Which statements are correct?

- A both 1 and 2
B 1 only
C 2 only
D neither 1 nor 2

16 The atomic radii and ionic radii for three elements in Period 3 are shown.

	atomic radius / nm	ionic radius / nm
element X	0.118	0.053
element Y	0.099	0.180
element Z	0.160	0.072

Using this data, which statement is correct?

- A Element X has lower electrical conductivity than element Y.
B Element Y has a higher melting point than element X.
C Element Y and element Z react to form an ionic compound.
D Element Z forms ionic compounds by gaining electrons.

19 The trends seen in Group 2 can be used to predict the properties of radium and its compounds.

Which statement is correct?

- A** Radium has the highest second ionisation energy of the elements in Group 2.
- B** Radium hydroxide is the least soluble of the hydroxides of the elements in Group 2.
- C** Radium carbonate has the lowest thermal stability of the carbonates of the elements in Group 2.
- D** Radium reacts faster with water than the other elements in Group 2.

40 A sample of magnesium contains the isotopes ^{24}Mg , ^{25}Mg and ^{26}Mg only.

The percentage abundance of ^{25}Mg and ^{26}Mg is the same.

The relative atomic mass of magnesium in the sample is 24.3.

What is the percentage abundance of ^{24}Mg ?

- A** 10% **B** 20% **C** 60% **D** 80%

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.02 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ ($4.18 \text{ J g}^{-1} \text{ K}^{-1}$)

The Periodic Table of Elements

Group																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																			
1	2	Key										13	14	15	16	17	18																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		
3	4	atomic number atomic symbol name relative atomic mass										5	6	7	8	9	2																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		
Li lithium 6.9	Be beryllium 9.0	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57–71 lanthanoids	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89–103 actinoids	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	524	525	526	527	528	529	530	531	532	533	534	535	536	537	538	539	540	541	542	543	544	545	546	547	548	549	550	551	552	553	554	555	556	557	558	559	560	561	562	563	564	565	566	567	568	569	570	571	572	573	574	575	576	577	578	579	580	581	582	583	584	585	586	587	588	589	590	591	592	593	594	595	596	597	598	599	600	601	602	603	604	605	606	607	608	609	610	611	612	613	614	615	616	617	618	619	620	621	622	623	624	625	626	627	628	629	630	631	632	633	634	635	636	637	638	639	640	641	642	643	644	645	646	647	648	649	650	651	652	653	654	655	656	657	658	659	660	661	662	663	664	665	666	667	668	669	670	671	672	673	674	675	676	677	678	679	680	681	682	683	684	685	686	687	688	689	690	691	692	693	694	695	696	697	698	699	700	701	702	703	704	705	706	707	708	709	710	711	712	713	714	715	716	717	718	719	720	721	722	723	724	725	726	727	728	729	730	731	732	733	734	735	736	737	738	739	740	741	742	743	744	745	746	747	748	749	750	751	752	753	754	755	756	757	758	759	760	761	762	763	764	765	766	767	768	769	770	771	772	773	774	775	776	777	778	779	780	781	782	783	784	785	786	787	788	789	790	791	792	793	794	795	796	797	798	799	800	801	802	803	804	805	806	807	808	809	810	811	812	813	814	815	816	817	818	819	820	821	822	823	824	825	826	827	828	829	830	831	832	833	834	835	836	837	838	839	840	841	842	843	844	845	846	847	848	849	850	851	852	853	854	855	856	857	858	859	860	861	862	863	864	865	866	867	868	869	870	871	872	873	874	875	876	877	878	879	880	881	882	883	884	885	886	887	888	889	890	891	892	893	894	895	896	897	898	899	900	901	902	903	904	905	906	907	908	909	910	911	912	913	914	915	916	917	918	919	920	921	922	923	924	925	926	927	928	929	930	931	932	933	934	935	936	937	938	939	940	941	942	943	944	945	946	947	948	949	950	951	952	953	954	955	956	957	958	959	960	961	962	963	964	965	966	967	968	969	970	971	972	973	974	975	976	977	978	979	980	981	982	983	984	985	986	987	988	989	990	991	992	993	994	995	996	997	998	999	1000

lanthanoids

57	La	lanthanum	138.9	58	Ce	cerium	140.1	59	Pr	praseodymium	140.9	60	Nd	neodymium	144.2	61	Pm	promethium	—	62	Sm	samarium	150.4	63	Eu	euroium	152.0	64	Gd	gadolinium	157.3	65	Tb	terbium	158.9	66	Dy	dysprosium	162.5	67	Ho	holmium	164.9	68	Er	erbium	167.3	69	Tm	thulium	168.9	70	Yb	ytterbium	173.1	71	Lu	lutetium	175.0
89	Ac	actinium	227	90	Th	thorium	232	91	Pa	protactinium	231	92	U	uranium	238	93	Np	neptunium	237	94	Pu	plutonium	244	95	Am	americium	243	96	Cm	curium	247	97	Bk	berkelium	247	98	Cf	californium	251	99	Es	einsteinium	252	100	Fm	fermium	257	101	Md	meitnerium	288	102	No	nobelium	289	103	Lr	lawrencium	260

actinoids

- 38** Butylamine can be produced by the reaction of butanenitrile with hydrogen in the presence of a suitable catalyst.

Which volume of hydrogen, measured at room conditions, is required to react completely with 0.500 g of butanenitrile?

- A** 145 cm³ **B** 174 cm³ **C** 289 cm³ **D** 348 cm³

- 3** What contains 9.03×10^{23} oxygen atoms?

- A** 0.25 mol aluminium oxide
B 0.75 mol sulfur dioxide
C 1.5 mol sulfur trioxide
D 3.0 mol water

- 5** Which statement explains why sodium and potassium have different melting points?

- A** The attraction between cations and delocalised electrons is stronger in sodium.
B The attraction between cations and anions is stronger in sodium.
C The attraction between atoms is stronger in sodium.
D The attraction between nuclei and shared electron pairs is stronger in sodium.

- 1** Chromium, Cr, and its compounds are widely used in many chemical reactions.

- (a)** Cr exists as four stable isotopes.

- (i)** The most common isotope of Cr is chromium-52.

Determine the number of protons, neutrons and electrons in an atom of chromium-52.

number of protons neutrons electrons [1]

- (ii)** Describe how an atom of chromium-54 differs from an atom of chromium-52. Refer to numbers of particles in your answer.

..... [1]

- (iii)** The relative isotopic masses of the isotopes of Cr can be determined using mass spectrometry.

State what other information is needed to calculate the relative atomic mass, A_r , of Cr.

..... [1]

- (iv)** Atoms of ^{52}Cr , ^{53}Cr and ^{54}Cr make up more than 95% of naturally occurring chromium atoms. The A_r of naturally occurring Cr is 51.996.

Suggest what these statements imply about the relative isotopic mass of the fourth stable isotope of chromium.

..... [1]

- (b)** The shorthand electronic configuration of chromium is $[\text{Ar}] 3d^5 4s^1$.



- (i)** Complete the full electronic configuration of chromium.

..... $3d^5 4s^1$ [1]

- (ii)** Deduce the total number of unpaired electrons in an atom of chromium.

..... [1]

- (c) Acidified dichromate(VI) ions will convert methanal, CH_2O , to carbon dioxide. The movement of electrons to or from relevant species is shown in the following half-equations.



- (i) Identify the species that is reduced in half-equation 2. Explain your answer.

..... [1]

- (ii) The oxidation state of the carbon atom in methanal is 0.

Calculate the oxidation state of carbon in carbon dioxide.

..... [1]


- (iii) Construct the ionic equation for the reaction of dichromate(VI) ions with methanal in acidic conditions.

..... [2]

- (iv) Methanal is a liquid at -30°C but carbon dioxide is a gas at this temperature. Explain why.

.....
.....
.....
..... [2]

- (d) In acidic conditions, a dynamic equilibrium is established between $\text{CrO}_4^{2-}(\text{aq})$ and $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$.



- (i) State what is meant by dynamic equilibrium.

.....
..... [1]

- (ii) Identify the condition necessary to establish dynamic equilibrium.

..... [1]

- (iii) $\text{CrO}_4^{2-}(\text{aq})$ ions are yellow and $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$ ions are orange.

State what is observed when the following changes are made to an equilibrium mixture of acidified $\text{CrO}_4^{2-}(\text{aq})$ and $\text{Cr}_2\text{O}_7^{2-}(\text{aq})$ ions.

Explain your answers.

- The equilibrium mixture is warmed gently.

observation

explanation

.....
.....

- Dilute $\text{HCl}(\text{aq})$ is added to the equilibrium mixture.

observation

explanation

.....
.....

[4]

- (e) Chromium(IV) fluoride, CrF_4 , is a covalent molecule that shows similar chemical properties to SiCl_4 .

Suggest the type of reaction that occurs when CrF_4 is placed in water.

Construct a relevant equation for this reaction.

type of reaction

equation

[2]

2 Aluminium oxide, Al_2O_3 , and phosphorus(V) oxide, P_4O_{10} , are both used as reagents and catalysts.

- (a) The melting point of Al_2O_3 is 2072 °C. The melting point of P_4O_{10} is 340 °C.

Explain the difference in the melting points of these two compounds.

.....

 [3]

- (b) A 5.00dm³ sealed flask contains 0.400 mol of CO(g) and 0.800 mol of H₂(g) and an Al_2O_3 catalyst. The flask is heated to a temperature of 290 °C and allowed to reach equilibrium. Equation 1 shows the reaction.



The equilibrium constant, K_c , of equation 1 is given.

$$K_c = \frac{[CH_3OH]}{[CO][H_2]^2}$$

- (i) State the units of K_c .

..... [1]

- (ii) The equilibrium mixture contains 0.280 mol of CH₃OH(g).

Calculate the value of K_c .

Give your answer to **three** significant figures.

value of K_c = [3]

- (iii) State and explain the effect, if any, on the value of K_c when the overall pressure in the sealed flask is increased.

.....
 [1]

- (c) P_4O_{10} catalyses the reversible reaction of CO with H₂ to form CH₃OH.



P_4O_{10} then acts as a dehydrating agent, causing CH₃OH to form CH₃OCH₃.

- (i) Explain how the presence of a catalyst affects a chemical reaction.

.....

 [1]

- (ii) Construct an equation for the dehydration reaction of CH₃OH to form CH₃OCH₃.

..... [1]

- (iii) Write an equation to show the reaction of P_4O_{10} with an excess of water.

..... [1]

[Total: 11]

4 Compounds **P** and **Q** are structural isomers.




Fig. 4.1

(a) (i) Define structural isomerism.

.....
 [2]

(ii) **P** shows geometrical isomerism.

Draw the geometrical isomer of **P**. Explain why the two isomers are **not** identical.



explanation

 [2]

(iii) Both **P** and **Q** react with aqueous bromine.

Name the mechanism of this reaction.

..... [1]

(iv) **Q** is oxidised by hot concentrated acidified $\text{KMnO}_4(\text{aq})$, forming two different organic products.

Construct an equation for this reaction. Use [O] to represent an atom of oxygen from the oxidising agent.

$(\text{CH}_3)_2\text{C}=\text{CHCH}_3$ [2]

(v) **Q** reacts with $\text{HBr}(\text{g})$ to produce two structural isomers, **R** and **S**, as shown in Fig. 4.2.





Fig. 4.2

State and explain why isomer **S** is the major product of the reaction.

.....
 [2]

(vi) The mass spectrum of **R** shows peaks at $m/e = 150$ and $m/e = 152$.

Suggest structures for the ions responsible for these peaks.



[2]

(b) Fig. 4.3 shows a synthesis starting from **T**, a different isomer of **R** and **S**.




Fig. 4.3

(i) Identify the reagent and conditions for reaction 1.

..... [1]

(ii) Reaction 2 is a hydrolysis reaction.

Construct an equation for reaction 2.

$(\text{C}_2\text{H}_5)_2\text{CHCN}$ [1]

(iii) **V** reacts with propan-2-ol in the presence of a catalytic amount of H_2SO_4 to form organic compound **W**.

Complete Table 4.1 to give details of this reaction.

Table 4.1

	reaction of V with propan-2-ol
type of reaction	
functional group formed	
molecular formula of organic product W	

[3]

[Total: 16]

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.02 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ ($4.18 \text{ J g}^{-1} \text{ K}^{-1}$)

The Periodic Table of Elements

Group																							
		1 H hydrogen 1.0														13	14	15	16	17	18		
		Key																					
		atomic number atomic symbol name relative atomic mass																					
3	4																	5	6	7	8	9	10
Li lithium 6.9	Be beryllium 9.0																	B boron 10.8	C carbon 12.0	N nitrogen 14.0	O oxygen 16.0	F fluorine 19.0	Ne neon 20.2
11	12																	Al aluminum 27.0	Si silicon 28.1	P phosphorus 31.0	S sulfur 32.1	Cl chlorine 35.5	Ar argon 39.9
Na sodium 23.0	Mg magnesium 24.3																	13	14	15	16	17	18
19	20																	31	32	33	34	35	36
K potassium 39.1	Ca calcium 40.1																	Ga gallium 69.7	Ge germanium 72.6	As arsenic 74.9	Se selenium 79.0	Br bromine 79.9	Kr krypton 83.8
37	38																	49	50	51	52	53	54
Rb rubidium 85.5	Sr strontium 87.6																	In indium 114.8	Sn tin 118.7	Sb antimony 121.8	Te tellurium 127.6	I iodine 126.9	Xe xenon 131.3
55	56																	81	82	83	84	85	86
Cs caesium 132.9	Ba barium 137.3																	Tl thallium 204.4	Pb lead 207.2	Bi bismuth 209.0	Po polonium —	At astatine —	Rn radon —
87	88																	113	114	115	116	117	118
Fr francium —	Ra radium —																	Nh nihonium —	Fl flerovium —	Mc moscovium —	Lv livermorium —	Ts tennessine —	Og oganeson —

lanthanoids

actinoids

57	58	59	60	61	62	63	64	65	66	67	68	69	70	71
La lanthanum 138.9	Ce cerium 140.1	Pr praseodymium 140.9	Nd neodymium 144.2	Pm promethium —	Sm samarium 150.4	Eu europium 152.0	Gd gadolinium 157.3	Tb terbium 158.9	Dy dysprosium 162.5	Ho holmium 164.9	Er erbium 167.3	Tm thulium 168.9	Yb ytterbium 173.1	Lu lutetium 175.0
89	90	91	92	93	94	95	96	97	98	99	100	101	102	103
Ac actinium —	Th thorium 232.0	Pa protactinium 231.0	U uranium 238.0	Np neptunium —	Pu plutonium —	Am americium —	Cm curium —	Bk berkelium —	Cf californium —	Es einsteinium —	Fm fermium —	Md mendelevium —	No nobelium —	Lr lawrencium —

2 Hydrated zinc sulfate has the formula $\text{ZnSO}_4 \cdot y\text{H}_2\text{O}$ where **y** is an integer.

Hydrated zinc sulfate decomposes when heated, losing only its water of crystallisation and becoming anhydrous.

You will determine the value of **y** in $\text{ZnSO}_4 \cdot y\text{H}_2\text{O}$ by heating the hydrated salt until it becomes anhydrous.

FA 4 is hydrated zinc sulfate, $\text{ZnSO}_4 \cdot y\text{H}_2\text{O}$.

(a) Method

- Weigh the crucible with its lid. Record the mass in the space for Results.
- Add between 3.20 g and 3.40 g of **FA 4** to the crucible.
- Weigh the crucible with its lid and **FA 4**. Record the mass.
- Place the crucible on the pipeclay triangle. Gently heat the crucible and contents for approximately 2 minutes with the lid on.
- Remove the lid. Heat the crucible and contents strongly for approximately 4 minutes.
- Replace the lid and leave the crucible and residue to cool for at least 5 minutes.

You may wish to begin work on Question 3 while the crucible is cooling.

- Weigh the crucible with its lid and its contents. Record the mass.
- Remove the lid. Heat the crucible strongly for approximately 3 minutes.
- Replace the lid and leave the crucible and residue to cool for at least 5 minutes.
- Weigh the crucible with its lid and its contents. Record the mass.
- Calculate the mass of **FA 4** used and the mass of residue obtained. Record the masses.

Results

I	
II	
III	
IV	
V	

(b) Calculations

- (i) Calculate the amount, in mol, of anhydrous zinc sulfate residue formed in the decomposition of **FA 4**.

amount of ZnSO_4 = mol

Calculate the amount, in mol, of water of crystallisation lost.

amount of H_2O = mol
[2]

- (ii) Calculate the value of **y** in the formula $\text{ZnSO}_4 \cdot y\text{H}_2\text{O}$.

Show your working.

y = [2]

- (c) A student suggests using this thermal decomposition method to investigate the number of moles of water of crystallisation in hydrated ethanedioic acid. The teacher says that this method is unsuitable.

Suggest why this method is unsuitable.

.....
..... [1]

[Total: 10]

Qualitative analysis

For each test you should record all your observations in the spaces provided.

Examples of observations include:

- colour changes seen
- the formation of any precipitate and its solubility (where appropriate) in an excess of the reagent added
- the formation of any gas and its identification (where appropriate) by a suitable test.

You should record clearly at what stage in a test an observation is made.

Where no change is observed, you should write 'no change'.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

If any solution is warmed, a boiling tube must be used. If a solid is heated, a hard-glass test-tube must be used.

Rinse and reuse test-tubes and boiling tubes where possible.

No additional tests should be attempted.

2 Effusion is the process in which a gas escapes through a small hole.

A student investigates the relationship between rate of effusion and relative molar mass of a gas using the apparatus shown in Fig. 2.1.




Fig. 2.1

The following method is used:

- step 1 Turn the tap and remove any gas from the syringe through the side-arm tube, by pushing in the plunger.
- step 2 Add 50 cm³ of the gas being tested to the syringe through the side-arm tube.
- step 3 Remove the gas from the syringe, through the side-arm tube, by pushing in the plunger.
- step 4 Add 70 cm³ of the gas being tested to the syringe through the side-arm tube.
- step 5 Turn the tap to connect the syringe to the tube with the aluminium foil and small hole.
- step 6 Allow the syringe plunger to fall and start a timer when the volume of gas in the syringe reaches 60 cm³.
- step 7 Stop the timer when the volume of gas in the syringe reaches 10 cm³. Record the time taken.
- step 8 Repeat steps 1 to 7 with different gases.

- (a) Suggest why the student adds 50 cm³ of the gas being tested to the syringe in step 2 and then removes this gas in step 3.

.....
..... [1]

- (b) The student's results are shown in Table 2.1.

Table 2.1

gas	hydrogen, H ₂	helium, He	neon, Ne	argon, Ar	krypton, Kr
relative molar mass, <i>M</i>	2.0	4.0	20.2	39.9	83.8
$\sqrt{\frac{1}{M}}$					
time taken /s	10.8	15.3	34.5	39.7	70.4
rate of effusion /cm ³ s ⁻¹					

$$\text{rate of effusion} = \frac{\text{volume of gas}}{\text{time taken}}$$

- (i) Complete Table 2.1.

Give the values for $\sqrt{\frac{1}{M}}$ to **three** significant figures.

Give the values for rate of effusion to **two** decimal places.

[2]

- (ii) Identify the dependent variable in this experiment.

..... [1]

- (iii) Identify a variable, other than temperature, that is controlled when carrying out this experiment.

..... [1]

- (c) Plot a graph on the grid in Fig. 2.2 to show the relationship between rate of effusion and $\sqrt{\frac{1}{M}}$.
Use a cross (x) to plot each data point. Draw a suitable line of best fit.




Fig. 2.2

21

[2]

- (d) Circle **one** point on the graph in Fig. 2.2 which you consider to be most anomalous.

Suggest **one** reason for this anomaly. Assume there is no error in $\sqrt{\frac{1}{M}}$.

.....
..... [1]

- (e) Graham's law of effusion can be expressed as:

the rate of effusion of a gas is proportional to $\sqrt{\frac{1}{M}}$.

State whether or not the student's results support Graham's law of effusion.

Explain your answer, using the graph in Fig. 2.2.

.....
..... [1]

- (f) Suggest how the position of the plotted points relative to the line of best fit in Fig. 2.2 is related to the reliability of the results.

.....
..... [1]

- (g) The student then repeats this method to determine the value of M of a sample of natural gas.

The time recorded in step 7 is 31.6 s.

- (i) Use the graph in Fig. 2.2 and the student's result to calculate the value of M for this sample.

$M =$ [2]

- (ii) Natural gas is a mixture of mainly methane, CH_4 , with small amounts of other gases.

Suggest what your calculated value of the M of natural gas in (g)(i) tells you about the other gases in the mixture.

.....
..... [1]

22

(h) The experiment described in (g) is repeated at a higher temperature.

Suggest how the rate of effusion for this sample of natural gas would change, if at all.

Explain your answer.

effect on the rate of effusion

explanation

[1]

[Total: 14]

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.02 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)

The Periodic Table of Elements

Group																		
1	2	Key														17	18	
		atomic number atomic symbol name relative atomic mass																
3 Li lithium 6.9	4 Be beryllium 9.0															1 H hydrogen 1.0		
11 Na sodium 23.0	12 Mg magnesium 24.3	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	
19 K potassium 39.1	20 Ca calcium 40.1	21 Sc scandium 45.0	22 Ti titanium 47.9	23 V vanadium 50.9	24 Cr chromium 52.0	25 Mn manganese 54.9	26 Fe iron 55.8	27 Co cobalt 58.9	28 Ni nickel 58.7	29 Cu copper 63.5	30 Zn zinc 65.4	31 Ga gallium 69.7	32 Ge germanium 72.6	33 As arsenic 74.9	34 Se selenium 79.0	35 Br bromine 79.9	36 Kr krypton 83.6	
37 Rb rubidium 85.5	38 Sr strontium 87.6	39 Y yttrium 88.9	40 Zr zirconium 91.2	41 Nb niobium 92.9	42 Mo molybdenum 95.9	43 Tc technetium —	44 Ru ruthenium 101.1	45 Rh rhodium 102.9	46 Pd palladium 106.4	47 Ag silver 107.9	48 Cd cadmium 112.4	49 In indium 114.8	50 Sn tin 118.7	51 Sb antimony 121.8	52 Te tellurium 127.6	53 I iodine 126.9	54 Xe xenon 131.3	
55 Cs caesium 132.9	56 Ba barium 137.3	57–71 lanthanoids		72 Hf hafnium 178.5	73 Ta tantalum 180.9	74 W tungsten 183.8	75 Re rhenium 186.2	76 Os osmium 190.2	77 Ir iridium 192.2	78 Pt platinum 195.1	79 Au gold 197.0	80 Hg mercury 200.6	81 Tl thallium 204.4	82 Pb lead 207.2	83 Bi bismuth 209.0	84 Po polonium —	85 At astatine —	86 Rn radon —
87 Fr francium —	88 Ra radium —	89–103 actinoids		104 Rf rutherfordium —	105 Db dubnium —	106 Sg seaborgium —	107 Bh bohrium —	108 Hs hassium —	109 Mt meitnerium —	110 Ds darmstadtium —	111 Rg roentgenium —	112 Cn copernicium —	113 Nh nihonium —	114 Fl flerovium —	115 Mc moscovium —	116 Lv livermorium —	117 Ts tennessine —	118 Og oganesson —
lanthanoids																		
57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.2	61 Pm promethium —	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0				
actinoids																		
89 Ac actinium —	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium —	94 Pu plutonium —	95 Am americium —	96 Cm curium —	97 Bk berkelium —	98 Cf californium —	99 Es einsteinium —	100 Fm fermium —	101 Md mendelevium —	102 No nobelium —	103 Lr lawrencium —				