

35 Which compound is a product of the hydrolysis of $\text{CH}_3\text{CO}_2\text{CH}_2\text{C}_2\text{H}_5$ using aqueous sodium hydroxide?

- A** $\text{CH}_3\text{CO}_2^-\text{Na}^+$
B $\text{CH}_3\text{CO}_2\text{H}$
C $\text{C}_2\text{H}_5\text{CH}_2\text{O}^-\text{Na}^+$
D $\text{C}_2\text{H}_5\text{CH}_2\text{CO}_2^-\text{Na}^+$

A-Level Chemistry: **w25 14**

36 The structural formula of compound P is $\text{CH}_3\text{CH}_2\text{COOCH}_2\text{CH}_3$.

P can be formed by reacting together two organic compounds in the presence of a suitable catalyst.

Which pair of compounds could react together to produce P?

- A** ethanoic acid and propanoic acid
B ethanoic acid and propan-1-ol
C ethanol and propanoic acid
D ethanol and propan-1-ol

A-Level Chemistry: **w25 33**

3 Half-fill the 250 cm^3 beaker with water and place it on a tripod and gauze. Heat the water until boiling then switch off your Bunsen burner. This will be your hot water bath for use in **(b)(i)**. Start **(a)(i)** while the water is heating.

(a) **FA 4** is an aqueous solution containing three ions. One ion is **not** listed in the Qualitative analysis notes. **FA 5** is an aqueous solution containing two ions. The cation in **FA 5** is listed in the Qualitative analysis notes. The anions in both **FA 4** and **FA 5** contain sulfur.

- (i)** Carry out the following tests using a 1 cm depth of either **FA 4** or **FA 5** in a test-tube. Record your observations in Table 3.1.

Table 3.1

<i>test</i>	<i>observations</i>	
	FA 4	FA 5
Test 1 Add aqueous sodium hydroxide.		
Test 2 Add aqueous ammonia.		
Test 3 Add aqueous sodium carbonate.		
Test 4 Add aqueous barium chloride or aqueous barium nitrate.		
Test 5 Add acidified aqueous potassium manganate(VII).		

[5]

- (ii)** From your observations in **(a)(i)**, deduce the identities of the ions present in **FA 4** and **FA 5**. Give the formula of each ion in Table 3.2. If you are unable to identify an ion write 'unknown'.

Table 3.2

<i>ions present</i>	FA 4	FA 5
cations		
anions		

[3]

- (b) (i) Both **FA 6** and **FA 7** are one of propan-1-ol, propan-2-ol, methanoic acid or ethanoic acid. You will carry out tests to investigate the identities of **FA 6** and **FA 7**. For each test use a 1 cm depth of **FA 6** or **FA 7** in a test-tube. Record your observations in Table 3.3.

Table 3.3

test	observations	
	FA 6	FA 7
Test 1 Add a few drops of acidified aqueous potassium manganate(VII), then		
place the test-tube in the hot water bath.		
Test 2 Add a 2 cm depth of aqueous iodine followed by drops of aqueous sodium hydroxide until the colour just disappears.		
If no reaction is visible, place the test-tube in the hot water bath.		
Test 3 Add aqueous sodium carbonate.		

[3]

- (ii) From your observations in (b)(i) deduce the identities of **FA 6** and **FA 7**. Give reasons for your answers.

FA 6 is

reasons

.....

FA 7 is

reasons

.....

[2]

- (c) A student carrying out a similar set of tests identifies two unknown compounds to be ethanol and propanoic acid.

Write an equation for the reaction between these two compounds.

..... [1]

[Total: 14]

Qualitative analysis notes**1 Reactions of cations**

cation	reaction with	
	NaOH(aq)	NH ₃ (aq)
aluminium, Al ³⁺ (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH ₄ ⁺ (aq)	no ppt. ammonia produced on warming	–
barium, Ba ²⁺ (aq)	faint white ppt. is observed unless [Ba ²⁺ (aq)] is very low	no ppt.
calcium, Ca ²⁺ (aq)	white ppt. unless [Ca ²⁺ (aq)] is very low	no ppt.
chromium(III), Cr ³⁺ (aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	pale blue ppt. soluble in excess giving dark blue solution
iron(II), Fe ²⁺ (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe ³⁺ (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg ²⁺ (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn ²⁺ (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn ²⁺ (aq)	white ppt. soluble in excess	white ppt. soluble in excess

2 Reactions of anions

anion	reaction
carbonate, CO ₃ ²⁻	CO ₂ liberated by dilute acids
chloride, Cl ⁻ (aq)	gives white ppt. with Ag ⁺ (aq) (soluble in NH ₃ (aq))
bromide, Br ⁻ (aq)	gives cream/off-white ppt. with Ag ⁺ (aq) (partially soluble in NH ₃ (aq))
iodide, I ⁻ (aq)	gives pale yellow ppt. with Ag ⁺ (aq) (insoluble in NH ₃ (aq))
nitrate, NO ₃ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and Al foil
nitrite, NO ₂ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and Al foil; decolourises acidified aqueous KMnO ₄
sulfate, SO ₄ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (insoluble in excess dilute strong acids); gives white ppt. with high [Ca ²⁺ (aq)]
sulfite, SO ₃ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (soluble in excess dilute strong acids); decolourises acidified aqueous KMnO ₄
thiosulfate, S ₂ O ₃ ²⁻ (aq)	gives off-white/pale yellow ppt. slowly with H ⁺

3 Tests for gases

gas	test and test result
ammonia, NH ₃	turns damp red litmus paper blue
carbon dioxide, CO ₂	gives a white ppt. with limewater
hydrogen, H ₂	'pops' with a lighted splint
oxygen, O ₂	relights a glowing splint