



- 3 (a) Solid manganese(IV) oxide,  $\text{MnO}_2$ , catalyses the decomposition of hydrogen peroxide.



State the type of catalysis for this reaction. Explain your answer.

.....  
..... [1]

- (b) Hydrogen peroxide reacts with iodide ions in acidic conditions as shown.



The initial rate of this reaction is investigated with different concentrations of  $\text{H}_2\text{O}_2$ ,  $\text{I}^-$  and  $\text{H}^+$ . The results obtained are shown in Table 3.1.

Table 3.1

| experiment | $[\text{H}_2\text{O}_2]/\text{mol dm}^{-3}$ | $[\text{I}^-]/\text{mol dm}^{-3}$ | $[\text{H}^+]/\text{mol dm}^{-3}$ | initial rate/ $\text{mol dm}^{-3} \text{s}^{-1}$ |
|------------|---------------------------------------------|-----------------------------------|-----------------------------------|--------------------------------------------------|
| 1          | 0.0450                                      | 0.0300                            | 0.0125                            | $2.42 \times 10^{-3}$                            |
| 2          | 0.0225                                      | 0.0600                            | 0.0125                            | $2.42 \times 10^{-3}$                            |
| 3          | 0.0225                                      | 0.120                             | 0.0125                            | $4.84 \times 10^{-3}$                            |
| 4          | 0.0450                                      | 0.120                             | 0.0500                            | $9.68 \times 10^{-3}$                            |

- (i) Use the information in Table 3.1 to deduce the rate equation for this reaction.

Explain your reasoning.

.....  
.....  
.....  
.....  
.....  
.....  
.....  
..... [4]

- (ii) Use your rate equation from (b)(i) and the data from Experiment 1 to calculate the rate constant,  $k$ , for this reaction. Include the units of  $k$ .

$k = \frac{1}{\tau_1}$  ..... units ..... [2]

- (c) The rate of the thermal decomposition of azomethane,  $\text{CH}_3\text{N}=\text{NCH}_3$ , is investigated.




Fig. 3.1 shows the results obtained. The reaction is first order with respect to  $\text{CH}_3\text{N}=\text{NCH}_3$ .




Fig. 3.1

- (i) Use Fig. 3.1 to calculate **two** half-lives,  $t_{1/2}$ , to show that the reaction is first order.

.....  
.....  
.....  
..... [2]

- (ii) Use your answer to (c)(i) to calculate the rate constant,  $k$ , for the decomposition of azomethane.


$k = \dots\dots\dots \text{s}^{-1}$  [1]

- (d) Describe the effect of increasing temperature on the rate constant and on the rate of a reaction.

.....  
.....  
..... [1]

[Total: 11]

- 2 Ethanal,  $\text{CH}_3\text{CHO}$ , reacts with nitrogen dioxide,  $\text{NO}_2$ . The products of the first step of this reaction are a  $\text{CH}_3\dot{\text{C}}=\text{O}$  radical and a molecule of nitrous acid,  $\text{HNO}_2$ .





- (a) (i) Use **two** words to complete the sentence.

This reaction involves ..... of the single covalent bond between a hydrogen atom and a carbon atom in  $\text{CH}_3\text{CHO}$ .

[1]

- (ii) The hydrogen atom mentioned in (a)(i) forms a covalent bond with one of the oxygen atoms of an  $\text{NO}_2$  molecule. An  $\text{NO}_2$  molecule has a single, unpaired electron on the nitrogen atom. All electrons are paired in an  $\text{HNO}_2$  molecule.

Draw dot-and-cross diagrams of  $\text{NO}_2$  and  $\text{HNO}_2$  in the boxes. Show outer shell electrons only.



[1]

- (iii) Use VSEPR theory to predict the bond angle at the nitrogen atom in an  $\text{HNO}_2$  molecule.

bond angle = ..... [1]

- (b) The rate equation for the reaction between  $\text{CH}_3\text{CHO}$  and  $\text{NO}_2$  is shown.

$$\text{rate} = k[\text{CH}_3\text{CHO}][\text{NO}_2]$$

Under certain conditions, when the concentrations of both  $\text{CH}_3\text{CHO}$  and  $\text{NO}_2$  are  $0.200 \text{ mol dm}^{-3}$ , the rate of the reaction is  $1.53 \times 10^{-4} \text{ mol dm}^{-3} \text{ s}^{-1}$ .

Calculate the value of the rate constant,  $k$ , under these conditions. Give the units of  $k$ .

$k = \dots\dots\dots$  units ..... [2]

- (c) The reaction mixture described in (b) is monitored over a period of time.

Predict whether the graph of  $[\text{NO}_2]$  against time shows a constant half-life.

Explain your answer.

prediction .....

explanation .....

[1]

- (d)  $\text{NO}_2$  also reacts with ozone,  $\text{O}_3$ .

The rate equation is shown.

$$\text{rate} = k_1[\text{NO}_2]$$

Under certain conditions, the value of  $k_1$  is  $0.0848 \text{ s}^{-1}$ .

The reaction has a constant half-life under these conditions.

Calculate the half-life in seconds.

half-life = ..... s [1]

- (e)  $\text{NO}_2$  is present in the exhaust gases of cars. It can react with carbon monoxide,  $\text{CO}$ , on the surface of a heterogeneous catalyst in the car's catalytic converter.

Describe the mode of action of this heterogeneous catalyst.

.....  
.....  
.....  
.....  
.....  
.....  
..... [2]

[Total: 9]

4 (a) Nitrogen monoxide, NO, reacts with hydrogen, as shown in reaction 3.



(i) The rate equation for reaction 3 is shown.

$$\text{rate} = k[\text{H}_2][\text{NO}]^2$$

Complete Table 4.1.

**Table 4.1**

|                                                      |  |
|------------------------------------------------------|--|
| the order of reaction with respect to $[\text{H}_2]$ |  |
| the order of reaction with respect to $[\text{NO}]$  |  |
| the overall order of the reaction                    |  |

[1]

(ii) Predict how the initial rate for reaction 3 changes when the concentration of NO is halved.

..... [1]

(iii) Predict how the initial rate for reaction 3 changes when the concentrations of NO and  $\text{H}_2$  are both increased **three** times.

..... [1]

(iv) Suggest why reaction 3 is unlikely to proceed by a mechanism involving only a single step.

..... [1]

(v) Suggest equations for the **three** steps of the reaction mechanism for reaction 3.

Each step involves a reaction between **two** molecules.

step 1 .....  $\rightarrow$  .....

step 2 ..... + .....  $\rightarrow \text{N}_2\text{O} +$  .....

step 3  $\text{N}_2\text{O} +$  .....  $\rightarrow$  ..... + .....

[2]

(vi) Suggest the role of  $\text{N}_2\text{O}$  in this mechanism.

Explain your reasoning.

..... [1]

(b) Iodine,  $\text{I}_2$ , reacts with thiosulfate ions,  $\text{S}_2\text{O}_3^{2-}$ , as shown in reaction 4.



Reaction 4 is carried out in the presence of a large excess of  $\text{I}_2$ . Under these conditions, the reaction is first order with respect to  $[\text{S}_2\text{O}_3^{2-}]$  and zero order with respect to  $[\text{I}_2]$ .

The half-life,  $t_{1/2}$ , for reaction 4 is 720 s under certain conditions.

Calculate the value of the rate constant,  $k$ , for reaction 4. Include the units of  $k$ .

$k =$  ..... units ..... [1]

(c) The reaction between iodide ions,  $\text{I}^-(\text{aq})$ , and peroxydisulfate ions,  $\text{S}_2\text{O}_8^{2-}(\text{aq})$ , is catalysed by  $\text{Co}^{3+}(\text{aq})$ . The mechanism is similar to the mechanism of this reaction when  $\text{Fe}^{3+}(\text{aq})$  is used as the catalyst.

(i) State the type of catalysis that occurs in this reaction.

Explain your reasoning.

..... [1]

(ii) Write **two** equations to show how  $\text{Co}^{3+}(\text{aq})$  catalyses this reaction.

equation 1 .....

equation 2 ..... [2]

(iii) Suggest why this reaction is slow in the absence of  $\text{Co}^{3+}(\text{aq})$ .

..... [1]

[Total: 12]