

- 7 (a) State the relative acidities of bromoethanoic acid, BrCH_2COOH , chloroethanoic acid, ClCH_2COOH , ethanoic acid, CH_3COOH and ethanol, $\text{CH}_3\text{CH}_2\text{OH}$.

Explain your answer.

.....

most acidic least acidic

.....

.....

.....

.....

.....

.....

[4]

- (b) Fig. 7.1 shows the reaction of methylbenzene and ethanedioic acid with KMnO_4 .

Predict the major carbon-containing product for each of these reactions.





Fig. 7.1


[2]

(c) Polyamide **X** can be synthesised from ethanedioic acid and benzene-1,4-diamine.



(i) Draw the repeat unit of polyamide **X** in the box.

The new functional group formed should be shown displayed.



[2]

(ii) Benzene-1,4-diamine can be formed by reduction of 1,4-dinitrobenzene.

Complete the equation for this reduction.

[H] represents one atom of hydrogen from a reducing agent.



[1]

(d) Fig. 7.2 shows the two-step synthesis of the azo compound **W**.




Fig. 7.2

- (i) Suggest structures for compounds **V** and **W** and draw them in the boxes in Fig. 7.2. [2]
- (ii) Give the reagents and conditions for step 1.

..... [1]

[Total: 12]

9 Compound **X** is made from benzene by the route shown in Fig. 9.1.




Fig. 9.1

(a) Describe the bonding in benzene, C_6H_6 .

Your answer should include:

- the hybridisation of the six carbon atoms
- the types of bond between the carbon atoms
- the orbitals that overlap to produce the bonds between the carbon atoms
- the type of bond between the carbon atoms and the hydrogen atoms
- the orbitals that overlap to produce the bonds between the carbon atoms and the hydrogen atoms.

.....

.....

.....

.....

.....

.....

..... [3]

(b) Describe the reagents and conditions required for step 1 in Fig. 9.1.

.....

..... [1]

(c) In step 1 of Fig. 9.1 benzene reacts with $^+\text{CH}_3$.

Complete Fig. 9.2 to show the mechanism for this reaction, including:

- the movement of electron pairs using curly arrows
- the structure of the intermediate involved.




Fig. 9.2

[3]

(d) Describe the reagents and conditions required for step 2 of Fig. 9.1.

.....
..... [2]

(e) Identify the reagents required for step 3 of Fig. 9.1. Compound **W** is the product of this step.

..... [1]

(f) Name compound **W**.

..... [1]


(g) The reagents commonly used for step 4 will **not** reduce the $-\text{COOH}$ group.

Identify the reagents and conditions required for step 4 of Fig. 9.1.

..... [1]

(h) Benzene can also be used as a starting material to make compound **Y**.

compound **Y**



Describe how the route described in Fig. 9.1 (repeated below) can be changed to give compound **Y** instead of compound **X**.

Explain your answer.

.....
.....
.....
..... [2]




Fig. 9.1

[Total: 14]

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.02 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 $\text{J g}^{-1} \text{ K}^{-1}$)

