

14. Hydrocarbons

Starter Quiz · 课前小测

A-Level Chemistry

Name: _____ Score: _____

1. Alkanes are:

- ☐ saturated hydrocarbons with general formula C_nH_{2n+2}
- ☐ unsaturated hydrocarbons with a C=C bond
- ☐ compounds containing oxygen
- ☐ ionic compounds

2. Alkanes react with chlorine by free-radical substitution in the presence of:

- ☐ ultraviolet light
- ☐ a nucleophile
- ☐ water
- ☐ an acid catalyst

3. Alkenes are reactive because they have:

- ☐ a C=C double bond (their functional group)
- ☐ only single bonds
- ☐ an -OH group
- ☐ no functional group

4. In addition polymerisation:

- ☐ many alkene molecules join into one long chain, with no other product
- ☐ a small molecule like water is lost
- ☐ a polymer is split apart
- ☐ two different monomers alternate

5. Cracking is used to:

- ☐ break long-chain alkanes into shorter, more useful molecules
- ☐ join short alkanes into long ones
- ☐ add hydrogen to alkenes
- ☐ burn alkanes

6. Alkanes are saturated hydrocarbons containing only single bonds.

- ☐ True ☐ False

7. Alkanes react with halogens in UV light by free-radical _____.

8. Many ethene molecules join to form the polymer poly(_____).

9. An alkene decolourises bromine water, while an alkane leaves it orange — that is the standard test for a C=C double bond.

- ☐ True ☐ False

10. The initiation step is:

- ☐ UV light splitting the halogen into two radicals
- ☐ two radicals joining
- ☐ a radical reacting with the alkane
- ☐ the product forming

1. saturated hydrocarbons with general formula C_nH_{2n+2}

Alkanes have only single C–C bonds (saturated) and follow C_nH_{2n+2} .

2. ultraviolet light

UV light starts the reaction by splitting the halogen into radicals.

3. a C=C double bond (their functional group)

The electron-rich C=C double bond is the reactive functional group of alkenes (C_nH_{2n}).

4. many alkene molecules join into one long chain, with no other product

Alkenes open their double bonds and join into a polymer such as poly(ethene); nothing else is formed.

5. break long-chain alkanes into shorter, more useful molecules

Cracking turns heavy crude-oil fractions into smaller, more useful alkanes and alkenes.

6. True

General formula C_nH_{2n+2} .

7. substitution

Initiation, propagation and termination steps.

8. ethene

Addition polymerisation of the alkene.

9. True

The π electrons of the C=C attack the bromine; alkanes have no double bond, so nothing happens.

10. UV light splitting the halogen into two radicals

Initiation: $Cl_2 \rightarrow 2Cl \cdot$, the homolytic split that starts the chain.