


Chemical Bonding


A-Level Chemistry ■ Topic 3

A-Level Chemistry



What we will cover

- 1 Electronegativity
- 2 Ionic & metallic bonding
- 3 Covalent & coordinate bonds
- 4 Sigma & pi bonds
- 5 Shapes of molecules
- 6 Intermolecular forces



molecules in 3-D

1

Bonding types

Electronegativity

Electronegativity 电负性 is the power of an atom to attract bonding electrons towards itself. The difference in **Pauling electronegativity** 鲍林电负性 values predicts the bond type.

nuclear charge

more protons pull the pair more strongly

atomic radius

closer to the nucleus
⇒ stronger pull

shielding


inner shells
weaken the pull

It **rises** across a period and **falls** down a group. Fluorine is the most electronegative element. A large difference ⇒ ionic; a small difference ⇒ covalent.

Ionic bonding


Ionic bonding 离子键 – positive **cations** 阳离子 and negative **anions** 阴离子 – is the electrostatic attraction between oppositely charged **ions** 离子. A metal gives electrons to a non-metal. The ions pack into a giant **lattice** 晶格 held in every direction.

NaCl: sodium transfers its outer electron to chlorine $\rightarrow \text{Na}^+$ and a full-octet Cl^- .



Metallic bonding

Metallic bonding 金属键 is the attraction between positive metal ions and a sea of **delocalised electrons** 离域电子. The free electrons explain why metals conduct and are strong.



positive

metal ions


sea of delocalised

electrons (free to move)

2

Covalent bonding

Covalent and coordinate bonds



A **covalent bond** 共价键 is a shared pair of electrons.


In a **coordinate bond** 配位键 (dative) **both** electrons come from one atom – e.g. N's lone pair makes NH_4^+ .

Sigma and pi bonds


Covalent bonds form when **orbitals** 轨道 overlap:

- a **sigma bond** σ 键 – head-on **overlap** 重叠
- a **pi bond** π 键 – sideways p-orbital overlap

Double = $1\sigma + 1\pi$; triple = $1\sigma + 2\pi$. **Hybridisation** 杂化: sp , sp^2 , sp^3 .



σ bond:
head-on overlap




π bond: sideways overlap
of p orbitals


Bond energy and bond length

Bond energy 键能 energy to break **one mole** of a covalent bond in the **gas** state.


Bond length 键长 distance between the **centres** of the two bonded atoms.



347 kJ mol⁻¹



612 kJ mol⁻¹




838 kJ mol⁻¹

more shared pairs: a **shorter, stronger** bond

Shapes of molecules

VSEPR theory 价层电子对互斥理论:
electron pairs push apart as far as possible.

A **lone pair** 孤对电子 repels harder than a bonding pair, squeezing the **bond angle** 键角 ($\sim 2.5^\circ$ each) – so NH_3 is 107° , H_2O 104.5° .




linear
 CO_2



trigonal planar
 BF_3



tetrahedral
 CH_4




pyramidal
 NH_3



bent
 H_2O



trig. bipyramidal
 PF_5 , 120° & 90°



octahedral
 SF_6 , 90°

The shapes, and the angles that come with them

Linear	CO ₂ – two bonding regions, 180°.
Trigonal planar	BF ₃ – three regions, 120°, all in one plane.
Tetrahedral	CH ₄ – four regions, 109.5°. Lone pairs push harder: NH ₃ is pyramidal at 107°, H ₂ O bent at 104.5°.
Trigonal bipyramidal	PF ₅ – five regions, 120° in the plane and 90° to the axis.
Octahedral	SF ₆ – six regions, all 90°.

Count the regions of electron density around the central atom, then subtract for lone pairs.
The last two shapes need an atom that can **expand the octet**.


3

Intermolecular forces

Intermolecular forces

A bond is **polar** 极性 if the atoms differ in electronegativity (a **dipole** 偶极).

- **London forces** 伦敦色散力 – between all molecules; stronger with more electrons
- permanent dipole forces – between polar molecules



Van der Waals' forces, and when a molecule is polar

Van der Waals' forces

Van der Waals' forces 范德华力 is the general name for all intermolecular forces. There are two main types.

London dispersion

The instantaneous dipole-induced dipole force, or **London dispersion force** 伦敦色散力. Moving electrons make a brief **instantaneous dipole** 瞬时偶极, which induces a matching **induced dipole** 诱导偶极 next door. It acts between **all** molecules.

Permanent dipole

Between molecules that are always polar, because each carries a **permanent dipole** 永久偶极.


Dipole moment

If the bond dipoles in a molecule do **not** cancel, the whole molecule has a **dipole moment** 偶极矩 and is polar – which is why CO_2 is not, and H_2O is.

Shape decides polarity: identical bonds arranged symmetrically cancel, however polar each bond is on its own.

Bond polarity and dipoles

When two atoms with different **electronegativity** share a bond, the electrons sit closer to the more electronegative atom.




electron density pulled
towards **Cl**

dipole arrow

The bond has a **polarity** 极性: one end slightly negative ($\delta-$), the other slightly positive ($\delta+$).


Hydrogen bonding



Hydrogen bonding 氢键 forms when H is bonded to N, O or F and meets a lone pair on N, O or F.

It gives water a high **boiling point** 沸点, high **surface tension** 表面张力, and makes ice **less dense** (so it floats).

Dot-and-cross diagrams



H₂O: 2 bonding pairs
+ 2 lone pairs




N₂: triple bond
(3 shared pairs)

• electron from one atom
x electron from the other


Covalent dot-and-cross: each shared pair is one electron from each atom. Water has two bonding pairs and two lone pairs; nitrogen shares three pairs (a triple bond)

The electronegativity trend in the periodic table



Electronegativity rises across a period and falls down a group, so fluorine is the most electronegative element


Dot-and-cross: ionic bonding in NaCl



sodium gives its outer electron (×) to chlorine

Ionic bonding in NaCl: sodium transfers its single outer electron to chlorine, giving Na^+ and a full-octet Cl^-

Ionic bonding, continued (2)



A real crystal of rock salt (halite, NaCl); the cubic shapes mirror the giant ionic lattice inside

Exam tips

- For **shapes**, count bonding pairs and lone pairs, name the shape, then give the exact bond angle (e.g. NH_3 : pyramidal, 107°) – each lone pair lowers the angle by about 2.5° .
- A **dative (coordinate) bond** has both electrons from one atom (e.g. NH_4^+ , H_3O^+); draw the arrow from the lone pair.
- Name the intermolecular force precisely: **hydrogen bonding** needs H bonded to N, O or F; otherwise it is permanent-dipole or induced-dipole (van der Waals). Never call van der Waals forces “bonds”.
- Explain a physical property by stating **which forces are broken**, not just “strong bonds”.

4

Recap

Worked example – shape of a molecule

Predict the shape and bond angle of NH_3 .

- N has 5 outer electrons: 3 bond pairs + 1 **lone pair** 孤对电子 = 4 electron pairs.
- 4 pairs repel to a **tetrahedral** arrangement (109.5°).
- A lone pair repels **more** than a bond pair, so the angle closes.
- Name the shape from the **atoms only**: **pyramidal**, $\approx 107^\circ$.

Count electron **pairs** for the arrangement, then name the shape from the **atoms**. Each lone pair closes the angle by about 2.5° .

Topic 3 – key takeaways

1

Bond types

ionic, metallic, covalent (& coordinate)

2

Sigma & pi

single 1σ ; double $1\sigma 1\pi$; triple $1\sigma 2\pi$

3

Shapes

VSEPR; lone pairs squeeze the angle


4

Forces

London, dipole, hydrogen bonding

Check yourself

- 1 Which is the most electronegative element?
- 2 In a coordinate bond, where do both electrons come from?
- 3 How many sigma and pi bonds in a triple bond?
- 4 Why does ice float on water?



Answers 1. fluorine 2. one atom (a lone pair) 3. one σ and two π 4. hydrogen bonds hold an open structure (less dense)