

Equilibria

A-Level Chemistry ■ Topic 25

A-Level Chemistry



What we will cover

1 Conjugate acids & bases

2 pH & the constants

3 Calculating pH

4 Buffer solutions


5 Solubility & partition



pH paper estimates acidity

Conjugate acids and bases

conjugate pair: differ by one H^+



An acid loses H^+ to give its **conjugate base** 共轭碱; a base gains H^+ to give its **conjugate acid** 共轭酸.

A **conjugate pair** 共轭酸碱对 differs by just one H^+ – e.g. CH_3COOH and CH_3COO^- .

pH and the constants

$$\text{pH} = -\log[\text{H}^+]$$

$$K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$$

$$K_w = [\text{H}^+][\text{OH}^-]$$

A larger K_a 酸解离常数 (smaller $\text{p}K_a$) is a stronger acid; $K_w = 1.0 \times 10^{-14}$ at 298 K.


Calculating pH

strong acid – $[\text{H}^+] = \text{conc.}$; $\text{HCl } 0.050 \Rightarrow \text{pH} = 1.30$

strong base – $[\text{H}^+] = K_w/[\text{OH}^-]$; $\text{NaOH } 0.10 \Rightarrow \text{pH} = 13.0$

weak acid – $[\text{H}^+] = \sqrt{K_a[\text{HA}]}$; ethanoic $0.10 \Rightarrow \text{pH} = 2.87$

Buffer solutions



either way, the pH barely changes

A **buffer** 缓冲溶液 (a weak acid + its conjugate base) resists pH change:

- added H^+ removed by A^-
- added OH^- removed by HA

HCO_3^- keeps blood near pH 7.4.


Solubility product



The **solubility product** 溶度积 $K_{sp} = [Ag^+][Cl^-]$
(each ion to the power of its number).

The **common ion effect** 同离子效应: a salt is **less** soluble in a solution already containing one of its ions (Le Chatelier).

Partition coefficients



$$K_{pc} = \frac{\text{solvent 1 (non-polar)} \left[\text{solute in 1} \right]}{\text{solvent 2 (water)} \left[\text{solute in 2} \right]}$$

The **partition coefficient** 分配系数 is the ratio of a solute's concentrations in two immiscible solvents.

$$K_{pc} = \frac{[\text{solute in 1}]}{[\text{solute in 2}]}$$

It depends on the **polarity** 极性 of the solute & solvents.

Worked example – pH of a strong acid

Find the pH of $0.050 \text{ mol dm}^{-3}$ hydrochloric acid.

- HCl is a **strong** acid: it dissociates completely.
- So $[\text{H}^+] = 0.050 \text{ mol dm}^{-3}$ – the concentration itself.
- $\text{pH} = -\log_{10}[\text{H}^+] = -\log_{10}(0.050)$
- $= 1.30$.

Strong acid: $[\text{H}^+]$
is the concentration.

Weak acid: you must
go through K_a – never as-
sume full dissociation.

Topic 25 – key takeaways

1

Conjugate pairs

differ by one H^+

2

pH

$-\log[\text{H}^+]$; K_a , K_w for
weak/strong

3

Buffers

weak acid + its salt resist pH
change

4

Solubility

K_{sp} ; the common ion effect

Check yourself

- 1 Write the pH formula.
- 2 What does a buffer resist?
- 3 What two things make a buffer?
- 4 What does the common ion effect do to solubility?

Answers 1. $\text{pH} = -\log[\text{H}^+]$ 2. a change in pH 3. a weak acid & its conjugate base
4. lowers it