


Analytical Techniques

A-Level Chemistry ■ Topic 37

A-Level Chemistry



What we will cover

- 1 Thin-layer chromatography
- 2 Gas/liquid chromatography
- 3 NMR basics
- 4 Splitting & the $n + 1$ rule
- 5 A proton NMR spectrum



NMR uses a strong magnet

1

Chromatography

Thin-layer chromatography

The **baseline** 基线 is the starting line where the spots are placed; the **solvent front** 溶剂前沿 is the highest level the solvent reaches. The **mobile phase** 流动相 is the solvent that travels up the plate, the **stationary phase** 固定相 the coating it travels through. A spot of mixture travels up a plate. Components that adsorb more strongly to the stationary phase move less.


$$R_f = \frac{\text{distance of the spot}}{\text{distance of the solvent front}}$$

R_f is always between 0 and 1. Identify a component by matching R_f with a known standard on the same plate.

Gas/liquid chromatography

In **GLC** 气液色谱 a carrier gas sweeps the mixture over a liquid stationary phase.

Each component peaks at its own **retention time** 保留时间; the peak **area** gives its percentage in the mixture.



t_R identifies · peak area $\approx$ amount

2

NMR

NMR basics

carbon-13 NMR

each carbon **environment**
化学环境 gives one peak

proton NMR

chemical shift 化学位移 &
peak areas give the proton ratio

NMR 核磁共振 studies nuclei in a strong magnetic field; the number of peaks = the number of environments.

Splitting and the $n + 1$ rule

**singlet**

$$n = 0$$

**doublet**

$$n = 1$$

**triplet**

$$n = 2$$

**quartet**


$$n = 3$$

n equivalent neighbours split a peak into $n + 1$ lines

Splitting 裂分: n equivalent neighbouring protons split a peak into $n + 1$ lines.


- $0 \rightarrow$ **singlet** 单峰; $1 \rightarrow$ doublet
- $2 \rightarrow$ triplet; $3 \rightarrow$ quartet

A proton NMR spectrum



Ethanol: peak **areas** 3:2:1 count the H; a peak splits into $n+1$ by its **neighbouring** H.

A proton NMR spectrum, in detail



Ethanol: a CH_3 triplet, a CH_2 quartet & an OH singlet (areas 3:2:1).

Practical points

TMS 四甲基硅烷

the standard,
set at $\delta = 0$

**deuterated
solvent** 氘代溶剂


e.g. CDCl_3 : the solvent
gives **no** proton signal

shake with D_2O

O–H and N–H
peaks **disappear**


D_2O swaps the O–H/N–H hydrogen for deuterium, which gives no signal: the vanished peak **identifies** those protons.

A TLC plate under UV light



A real TLC plate under UV light: each glowing spot is a separated component of the mixture

Thin-layer chromatography, continued (2)



Thin-layer chromatography: the R_f value is how far the spot moved divided by how far the solvent front moved

Exam tips

- In ^{13}C NMR the number of peaks equals the number of carbon environments – use symmetry to count them.
- In ^1H NMR use chemical shift (data booklet), integration (ratio of H's) and splitting (the $n + 1$ rule): a triplet + quartet means an ethyl group.
- TMS is the reference ($\delta = 0$); a D_2O shake removes O-H and N-H peaks.
- Combine IR, mass spectrum and NMR to deduce a structure, stating which evidence gives which feature.

3

Recap

Worked example – reading an NMR spectrum

$\text{C}_3\text{H}_8\text{O}$ gives three ^1H peaks: 6H doublet, 1H septet, 1H singlet that vanishes with D_2O . Identify it.

- Three peaks $\Rightarrow$ three proton environments; the areas 6 : 1 : 1 give the H in each.
- The singlet that vanishes with D_2O is an O–H (it exchanges, and does not split).
- 6H as a doublet: two equivalent CH_3 next to one H ($n + 1 = 2$).
- 1H as a septet: that H has six neighbours ($n + 1 = 7$).
- $(\text{CH}_3)_2\text{CH} - \text{OH}$: propan-2-ol.

Peaks = environments; area = how many H; splitting = $n + 1$ neighbours. A peak lost with D_2O is O–H or N–H.

Topic 37 – key takeaways

1

TLC

$R_f = \text{spot distance} / \text{solvent distance}$

2

GLC

retention time; peak area = percentage

3

NMR

peaks = chemical environments

4

Splitting

n neighbours give $n + 1$ lines

Check yourself

- 1 What does the R_f value compare?
- 2 What does a GLC peak area tell you?
- 3 What does the number of NMR peaks tell you?
- 4 Three neighbouring protons split a peak into how many lines?



Answers 1. spot distance vs solvent distance 2. the percentage of that component
3. the number of carbon environments 4. four (a quartet)