


- 5 The bonding between two atoms of nitrogen in an N_2 molecule involves the hybridisation of atomic orbitals to form sp orbitals.


Which row is correct?

	formation of the σ bond between the nitrogen atoms in N_2	type of orbital which contains the lone pair of electrons on each nitrogen atom in N_2
A	an sp orbital from one atom overlaps with an sp orbital of the other atom	p
B	an sp orbital from one atom overlaps with an sp orbital of the other atom	sp
C	an s orbital from one atom overlaps with a p orbital of the other atom	p
D	an s orbital from one atom overlaps with a p orbital of the other atom	sp


- 25 The diagrams show skeletal formulas of some isomers of $\text{C}_6\text{H}_{12}\text{O}$.




1



2



3




4

Which statement is correct?


- A 1 and 3 are chain isomers of each other and positional isomers of each other.
- B 2 and 3 are functional group isomers of each other and both have a chiral centre.
- C 1 and 4 are functional group isomers of each other and both have a chiral centre.
- D 2 and 4 are positional isomers of each other and functional group isomers of each other.

26 The skeletal formulas of two compounds are shown.

progesterone




testosterone



Which statements about progesterone and testosterone are correct?

- A** They both contain a ketone group, and they both have geometrical isomers due to the C=C bond.
- B** They have the same molecular formula, and they both have geometrical isomers.
- C** They have the same number of chiral carbons, and they have the same molecular formula.
- D** They have the same number of chiral carbons, and they both contain a ketone group.

28 The diagram shows the skeletal formula of the hormone testosterone.



What is the molecular formula of testosterone?

- A** C₁₉H₂₈O₂
- B** C₁₇H₂₂O₂
- C** C₁₇H₂₄O₂
- D** C₁₉H₂₆O₂

4 Fig. 4.1 shows a possible synthesis of propene, C_3H_6 .




Fig. 4.1

(a) (i) Identify the type of reaction that occurs in reaction 1.

..... [1]

(ii) Suggest a suitable reagent for reaction 2.

..... [1]

(iii) Reaction 3 is an elimination reaction.

Write an equation for this reaction and identify the solvent and conditions used.

equation

solvent and conditions [2]

(iv) $C_2H_5CH_2OH$ can be directly converted to C_3H_6 .

Suggest the reagent and conditions for this conversion.

..... [1]

(b) Under suitable conditions, propene polymerises to form poly(propene).

Poly(propene) exhibits stereoisomerism.

(i) Identify the type of polymerisation that forms poly(propene) from propene.

..... [1]


(ii) Define stereoisomerism.

.....

..... [2]

(iii) Draw a section of poly(propene), showing **two** repeat units.

Use your diagram to identify the type of stereoisomerism shown by poly(propene). Explain your answer.



.....

.....

.....

.....

[3]



(iv) State **two** difficulties associated with the disposal of poly(propene).

1

2

[2]

(c) Under different conditions, two molecules of propene can combine to form compounds **X** and **Y**.



(i) Name **X**.

..... [1]

(ii) Fig. 4.2 shows the mass spectrum of either compound **X** or compound **Y**.




Fig. 4.2

Identify which of **X** and **Y** gives this mass spectrum.

Give **one** reason for your answer, referring to the fragmentation pattern.

.....
.....
..... [1]

(iii) The molecular ion peak in the spectrum in Fig. 4.2 has relative abundance 34.7.

Calculate the relative abundance of the $[M+1]^+$ peak in this spectrum.

relative abundance of $[M+1]^+$ peak = [1]

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.02 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)

The Periodic Table of Elements

Group																							
1	2											13	14	15	16	17	18						
		1 Hhydrogen1.0																2 Hehelium4.0					
		Key atomic number atomic symbol name relative atomic mass																					
3 Li lithium 6.9	4 Be beryllium 9.0																	5 B boron 10.8	6 C carbon 12.0	7 N nitrogen 14.0	8 O oxygen 16.0	9 F fluorine 19.0	10 Ne neon 20.2
11 Na sodium 23.0	12 Mg magnesium 24.3																	13 Al aluminium 27.0	14 Si silicon 28.1	15 P phosphorus 31.0	16 S sulfur 32.1	17 Cl chlorine 35.5	18 Ar argon 39.9
19 K potassium 39.1	20 Ca calcium 40.1	21 Sc scandium 45.0	22 Ti titanium 47.9	23 V vanadium 50.9	24 Cr chromium 52.0	25 Mn manganese 54.9	26 Fe iron 55.8	27 Co cobalt 58.9	28 Ni nickel 58.7	29 Cu copper 63.5	30 Zn zinc 65.4	31 Ga gallium 69.7		32 Ge germanium 72.6	33 As arsenic 74.9	34 Se selenium 79.0	35 Br bromine 79.9	36 Kr krypton 83.8					
37 Rb rubidium 85.5	38 Sr strontium 87.6	39 Y yttrium 88.9	40 Zr zirconium 91.2	41 Nb niobium 92.9	42 Mo molybdenum 95.9	43 Tc technetium —	44 Ru ruthenium 101.1	45 Rh rhodium 102.9	46 Pd palladium 106.4	47 Ag silver 107.9	48 Cd cadmium 112.4	49 In indium 114.8	50 Sn tin 118.7	51 Sb antimony 121.8	52 Te tellurium 127.6	53 I iodine 126.9	54 Xe xenon 131.3						
55 Cs caesium 132.9	56 Ba barium 137.3	57–71 lanthanoids		72 Hf hafnium 178.5	73 Ta tantalum 180.9	74 W tungsten 183.8	75 Re rhenium 186.2	76 Os osmium 190.2	77 Ir iridium 192.2	78 Pt platinum 195.1	79 Au gold 197.0	80 Hg mercury 200.6	81 Tl thallium 204.4	82 Pb lead 207.2	83 Bi bismuth 209.0	84 Po polonium —	85 At astatine —	86 Rn radon —					
87 Fr francium —	88 Ra radium —	89–103 actinoids		104 Rf rutherfordium —	105 Db dubnium —	106 Sg seaborgium —	107 Bh bohrium —	108 Hs hassium —	109 Mt meitnerium —	110 Ds darmstadtium —	111 Rg roentgenium —	112 Cn copernicium —	113 Nh nihonium —	114 Fl flerovium —	115 Mc moscovium —	116 Lv livermorium —	117 Ts tennessine —	118 Og oganesson —					

lanthanoids	57 La lanthanum 138.9	58 Ce cerium 140.1	59 Pr praseodymium 140.9	60 Nd neodymium 144.2	61 Pm promethium —	62 Sm samarium 150.4	63 Eu europium 152.0	64 Gd gadolinium 157.3	65 Tb terbium 158.9	66 Dy dysprosium 162.5	67 Ho holmium 164.9	68 Er erbium 167.3	69 Tm thulium 168.9	70 Yb ytterbium 173.1	71 Lu lutetium 175.0
	89 Ac actinium —	90 Th thorium 232.0	91 Pa protactinium 231.0	92 U uranium 238.0	93 Np neptunium —	94 Pu plutonium —	95 Am americium —	96 Cm curium —	97 Bk berkelium —	98 Cf californium —	99 Es einsteinium —	100 Fm fermium —	101 Md mendelevium —	102 No nobelium —	103 Lr lawrencium —

- 3 Propan-2-ol, $(\text{CH}_3)_2\text{CHOH}$, is sometimes added to fuel to help it burn.

Fig. 3.1 shows some reactions of propan-2-ol.




Fig. 3.1

- (a) (i) Draw the structure of organic compound **K**.

[1]

- (ii) State an observation you would make in reaction 2.

..... [1]

- (iii) State the type of reaction that is shown in reaction 3.

..... [1]

- (iv) Complete Fig. 3.2 to show the pi (π) bond in **L** that is formed from orbital overlap.




Fig. 3.2

[1]

- (b) Propan-2-ol reacts with sodium to produce $(\text{CH}_3)_2\text{CH-O}^-$ anions.

These anions react with 2-bromopropane to form compound **N**, as shown in Fig. 3.3.




Fig. 3.3

- (i) Write an equation for the reaction of propan-2-ol with sodium.

..... [1]

- (ii) The reaction of 2-bromopropane with $(\text{CH}_3)_2\text{CH-O}^-$ anions follows an $\text{S}_{\text{N}}1$ mechanism.

Complete Fig. 3.4 to show this mechanism. Include charges, dipoles, lone pairs of electrons and curly arrows, as appropriate.



Fig. 3.4

[3]

- (iii) Suggest how the rate of the $\text{S}_{\text{N}}1$ reaction would change, if at all, if 2-chloropropane were used instead of 2-bromopropane.

Explain your answer.

.....
.....
..... [2]

- (iv) **N** is also added to petrol to make it burn more smoothly.

Construct an equation for the complete combustion of **N**, $\text{C}_6\text{H}_{14}\text{O}$.

$\text{C}_6\text{H}_{14}\text{O}$ [1]

[Total: 11]

- 3** Half-fill the 250 cm³ beaker with water and place it on a tripod and gauze. Heat the water until boiling then switch off your Bunsen burner. This will be your hot water bath for use in **(b)(i)**. Start **(a)(i)** while the water is heating.
- (a)** **FA 4** is an aqueous solution containing three ions. One ion is **not** listed in the Qualitative analysis notes. **FA 5** is an aqueous solution containing two ions. The cation in **FA 5** is listed in the Qualitative analysis notes. The anions in both **FA 4** and **FA 5** contain sulfur.
- (i)** Carry out the following tests using a 1 cm depth of either **FA 4** or **FA 5** in a test-tube. Record your observations in Table 3.1.

Table 3.1

<i>test</i>	<i>observations</i>	
	FA 4	FA 5
Test 1 Add aqueous sodium hydroxide.		
Test 2 Add aqueous ammonia.		
Test 3 Add aqueous sodium carbonate.		
Test 4 Add aqueous barium chloride or aqueous barium nitrate.		
Test 5 Add acidified aqueous potassium manganate(VII).		

[5]

- (ii) From your observations in (a)(i), deduce the identities of the ions present in **FA 4** and **FA 5**. Give the formula of each ion in Table 3.2. If you are unable to identify an ion write 'unknown'.

Table 3.2

<i>ions present</i>	FA 4	FA 5
cations		
anions		

[3]

- (b) (i) Both **FA 6** and **FA 7** are one of propan-1-ol, propan-2-ol, methanoic acid or ethanoic acid. You will carry out tests to investigate the identities of **FA 6** and **FA 7**. For each test use a 1 cm depth of **FA 6** or **FA 7** in a test-tube. Record your observations in Table 3.3.

Table 3.3

<i>test</i>	<i>observations</i>	
	FA 6	FA 7
Test 1 Add a few drops of acidified aqueous potassium manganate(VII), then		
place the test-tube in the hot water bath.		
Test 2 Add a 2 cm depth of aqueous iodine followed by drops of aqueous sodium hydroxide until the colour just disappears.		
If no reaction is visible, place the test-tube in the hot water bath.		
Test 3 Add aqueous sodium carbonate.		

[3]

- (ii) From your observations in (b)(i) deduce the identities of **FA 6** and **FA 7**. Give reasons for your answers.

FA 6 is

reasons

.....

FA 7 is

reasons

.....

[2]

- (c) A student carrying out a similar set of tests identifies two unknown compounds to be ethanol and propanoic acid.

Write an equation for the reaction between these two compounds.

..... [1]

[Total: 14]

Qualitative analysis notes

1 Reactions of cations

cation	reaction with	
	NaOH(aq)	NH ₃ (aq)
aluminium, Al ³⁺ (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH ₄ ⁺ (aq)	no ppt. ammonia produced on warming	–
barium, Ba ²⁺ (aq)	faint white ppt. is observed unless [Ba ²⁺ (aq)] is very low	no ppt.
calcium, Ca ²⁺ (aq)	white ppt. unless [Ca ²⁺ (aq)] is very low	no ppt.
chromium(III), Cr ³⁺ (aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	pale blue ppt. soluble in excess giving dark blue solution
iron(II), Fe ²⁺ (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe ³⁺ (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg ²⁺ (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn ²⁺ (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn ²⁺ (aq)	white ppt. soluble in excess	white ppt. soluble in excess

2 Reactions of anions

anion	reaction
carbonate, CO ₃ ²⁻	CO ₂ liberated by dilute acids
chloride, Cl ⁻ (aq)	gives white ppt. with Ag ⁺ (aq) (soluble in NH ₃ (aq))
bromide, Br ⁻ (aq)	gives cream/off-white ppt. with Ag ⁺ (aq) (partially soluble in NH ₃ (aq))
iodide, I ⁻ (aq)	gives pale yellow ppt. with Ag ⁺ (aq) (insoluble in NH ₃ (aq))
nitrate, NO ₃ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and Al foil
nitrite, NO ₂ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and Al foil; decolourises acidified aqueous KMnO ₄
sulfate, SO ₄ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (insoluble in excess dilute strong acids); gives white ppt. with high [Ca ²⁺ (aq)]
sulfite, SO ₃ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (soluble in excess dilute strong acids); decolourises acidified aqueous KMnO ₄
thiosulfate, S ₂ O ₃ ²⁻ (aq)	gives off-white/pale yellow ppt. slowly with H ⁺

3 Tests for gases

gas	test and test result
ammonia, NH_3	turns damp red litmus paper blue
carbon dioxide, CO_2	gives a white ppt. with limewater
hydrogen, H_2	'pops' with a lighted splint
oxygen, O_2	relights a glowing splint