


34 An organometallic lithium compound, RLi , contains the nucleophile R^- .

$\text{CH}_3\text{CH}_2\text{Li}$ reacts with pentan-2-one. The mechanism is nucleophilic addition. The first step produces an anion which is then protonated to form the final product.


Which organic product is formed?



36 Compound Q reacts when heated with HCN in the presence of a catalyst to produce compound R.

Molecules of compound R each contain four carbon atoms.

What is compound Q?



38 Seven aldehydes and ketones

- contain only **one** oxygen atom per molecule
- have four or fewer carbon atoms in their structures.

Alkaline $\text{I}_2(\text{aq})$ is added to each of these carbonyl compounds separately.





How many of these carbonyl compounds produce a yellow precipitate with alkaline $\text{I}_2(\text{aq})$?

- A** 1 **B** 2 **C** 3 **D** 4

34 Three tests were performed on an unknown organic compound.

test reagent	test result
2,4-DNPH reagent	orange ppt
Tollens' reagent	no change
alkaline $I_2(aq)$	yellow ppt

What is the organic compound tested?




35 Butanone, $CH_3CH_2COCH_3$, and HCN mixed with a little KCN react together in a nucleophilic addition reaction.

Which description of the mechanism of this nucleophilic addition reaction is correct?

- A** The π bond pair of $C=O$ accepts H^+ from HCN and then CN^- acts as a nucleophile.
- B** A π bond pair from CN^- attacks the δ^+ C and HCN then donates H^+ to O.
- C** The lone pair on O accepts H^+ from HCN and then CN^- acts as a nucleophile.
- D** A lone pair from CN^- attacks the δ^+ C and then O^- accepts H^+ from HCN.

- 3 Propan-2-ol, $(\text{CH}_3)_2\text{CHOH}$, is sometimes added to fuel to help it burn.

Fig. 3.1 shows some reactions of propan-2-ol.



- (a) (i) Draw the structure of organic compound **K**.

[1]

- (ii) State an observation you would make in reaction 2.

..... [1]

- (iii) State the type of reaction that is shown in reaction 3.

..... [1]

- (iv) Complete Fig. 3.2 to show the pi (π) bond in **L** that is formed from orbital overlap.




Fig. 3.2

[1]

- (b) Propan-2-ol reacts with sodium to produce $(\text{CH}_3)_2\text{CH-O}^-$ anions.

These anions react with 2-bromopropane to form compound **N**, as shown in Fig. 3.3.




Fig. 3.3

- (i) Write an equation for the reaction of propan-2-ol with sodium.

..... [1]

- (ii) The reaction of 2-bromopropane with $(\text{CH}_3)_2\text{CH-O}^-$ anions follows an $\text{S}_{\text{N}}1$ mechanism.

Complete Fig. 3.4 to show this mechanism. Include charges, dipoles, lone pairs of electrons and curly arrows, as appropriate.



Fig. 3.4

[3]

- (iii) Suggest how the rate of the $\text{S}_{\text{N}}1$ reaction would change, if at all, if 2-chloropropane were used instead of 2-bromopropane.

Explain your answer.

.....
.....
..... [2]

- (iv) **N** is also added to petrol to make it burn more smoothly.

Construct an equation for the complete combustion of **N**, $\text{C}_6\text{H}_{14}\text{O}$.

$\text{C}_6\text{H}_{14}\text{O}$ [1]

[Total: 11]

9 (a) Explain why amides are much weaker bases than amines.

.....

 [2]

(b) Fig. 9.1 shows the preparation of 2-phenylethylamine, $\text{C}_6\text{H}_5\text{CH}_2\text{CH}_2\text{NH}_2$, by three different routes.




Fig. 9.1

(i) Suggest structures for compounds **M** and **N** and draw them in the boxes in Fig. 9.1. [2]

(ii) Give the reagents and conditions for reaction 1.

..... [1]

(c) Fig. 9.2 shows compound **H** which is a useful starting material in organic synthesis.




Fig. 9.2

H contains an alkene and an amine functional group.

Name the **other** functional group and give the classification of the amine group in **H**.

other functional group in **H**

classification of amine

[1]

(d) Ozonolysis involves the oxidative cleavage of a C=C bond in alkenes using ozone, O₃, as shown in Fig. 9.3.




Fig. 9.3

Fig. 9.4 shows the first step in this reaction which involves the formation of an ozonide intermediate.





Fig. 9.4


(i) On Fig. 9.4, draw **three** curly arrows to complete the mechanism of this step.

[2]

- (ii) **L** is formed from alkene **K**, C_8H_{14} , by a similar reaction to that shown in Fig. 9.3.



Suggest the structure of **K**.






[1]

[Total: 9]

Important values, constants and standards



molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.02 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 $\text{J g}^{-1} \text{ K}^{-1}$)

- 7 **P, Q, R, S, T, U, and V** are the seven structural isomers with molecular formula $C_5H_{10}O$ that have a carbonyl group.



- (a) Only one of these seven compounds has stereoisomers.

Draw three-dimensional diagrams of the **two** stereoisomers of this compound.



[2]

- (b) **P, Q, R, S, T, U, and V** are treated separately with alkaline $I_2(aq)$ and the product mixture is acidified.

- (i) Identify the **two** compounds that give a positive result with alkaline $I_2(aq)$.

..... and [1]




- (ii) Describe the observations when **one** of the compounds you have identified in (b)(i) is treated with alkaline $I_2(aq)$ and give the structural formulae of the **two** carbon-containing products of this reaction.

observations

two carbon-containing products

and

[2]



(c) The proton (^1H) NMR spectra of **P**, **Q**, **R**, **S**, **T**, **U**, and **V** are compared.

(i) Identify the only compound that gives a spectrum with two singlets and no other peaks.

..... [1]

Fig. 7.1 shows the spectrum obtained from one of the compounds.




Fig. 7.1

(ii) Identify the compound that gives this spectrum.

..... [1]

(iii) Name the splitting pattern of the peak at $\delta = 1.1$ in Fig. 7.1.

Give the reason for this splitting.

name

reason [1]

(iv) Identify the substance that gives the small peak at $\delta = 0$ in Fig. 7.1.

..... [1]

(d) The carbon-13 NMR spectra of **R**, **S**, **T** and **U** are compared.

Complete Table 7.1 to state the number of peaks in the spectrum of each compound.

Table 7.1

	compound	number of peaks
R	$(\text{CH}_3)_2\text{CHCH}_2\text{CHO}$	
S	$(\text{CH}_3)_3\text{CCHO}$	
T	$\text{CH}_3\text{CH}_2\text{CH}_2\text{COCH}_3$	
U	$\text{CH}_3\text{CH}_2\text{COCH}_2\text{CH}_3$	

[2]

[Total: 11]