

- 33** HOCH_2CHO is heated under reflux with an excess of acidified $\text{K}_2\text{Cr}_2\text{O}_7$ until there is no further reaction.

What is the final product of this reaction?

- A** HOOCCHO
B HOCH_2COOH
C HOCCOOH
D $\text{HOOCCH}_2\text{COOH}$

- 34** Three mixtures are heated under reflux.

Which mixtures will produce sodium propanoate as one product?

- 1 $\text{CH}_3\text{CH}_2\text{CHO} + \text{NaBH}_4(\text{aq})$
2 $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH} + \text{Na}(\text{s})$
3 $\text{CH}_3\text{CH}_2\text{CN} + \text{NaOH}(\text{aq})$

- A** 1, 2 and 3 **B** 1 and 2 only **C** 2 and 3 only **D** 3 only

- 35** Which compound is a product of the hydrolysis of $\text{CH}_3\text{CO}_2\text{CH}_2\text{C}_2\text{H}_5$ using aqueous sodium hydroxide?

- A** $\text{CH}_3\text{CO}_2^-\text{Na}^+$
B $\text{CH}_3\text{CO}_2\text{H}$
C $\text{C}_2\text{H}_5\text{CH}_2\text{O}^-\text{Na}^+$
D $\text{C}_2\text{H}_5\text{CH}_2\text{CO}_2^-\text{Na}^+$

- 37** The table shows the reagents and products of three reactions.

	reagents	products
1	ethanoic acid + Na_2CO_3	$\text{CH}_3\text{CO}_2\text{Na} + \text{H}_2\text{O} + \text{CO}_2$
2	ethanoic acid + Mg	$(\text{CH}_3\text{CO}_2)_2\text{Mg} + \text{H}_2\text{O}$
3	ethanoic acid + CH_3OH	$\text{CH}_3\text{CO}_2\text{CH}_3 + \text{H}_2\text{O}$

Which rows are correct?

- A** 1, 2 and 3 **B** 1 and 2 only **C** 1 and 3 only **D** 2 and 3 only

36 The structural formula of compound P is $\text{CH}_3\text{CH}_2\text{COOCH}_2\text{CH}_3$.

P can be formed by reacting together two organic compounds in the presence of a suitable catalyst.

Which pair of compounds could react together to produce P?

- A** ethanoic acid and propanoic acid
- B** ethanoic acid and propan-1-ol
- C** ethanol and propanoic acid
- D** ethanol and propan-1-ol

- 4 Fig. 4.1 shows how propane, C_3H_8 , can be converted to propanoic acid, CH_3CH_2COOH .





Fig. 4.1

- (a) Reaction 1 in Fig. 4.1 takes place in the presence of sunlight.



The reaction takes place via initiation, propagation and termination steps.

- (i) Name the mechanism shown by reaction 1.

..... [1]

- (ii) Complete the mechanism for reaction 1.

Construct equations to describe the steps of the mechanism.



propagation 1

propagation 2


termination $\rightarrow CH_3CH_2CH_2Cl$
[3]

- (iii) Reaction 1 is initiated by the bond fission of Cl_2 .

State the type of bond fission shown in the initiation step.

..... [1]

- (iv) Compound **Q** is a by-product of reaction 1.



Name **Q**.

..... [1]

- (v) The molecular formula of **Q** is $\text{C}_3\text{H}_6\text{Cl}_2$.

Identify the types of structural isomerism and stereoisomerism that a molecule with molecular formula $\text{C}_3\text{H}_6\text{Cl}_2$ can show.

type of structural isomerism

type of stereoisomerism

[2]

- (b) State the reagent and solvent required for reaction 2.

..... [1]

- (c) Reaction 3 takes place when $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ is heated under reflux with acidified potassium dichromate(VI) solution.

- (i) State the colour change that takes place in the reaction mixture.

..... [1]

- (ii) Construct an equation to represent reaction 3. Use [O] to represent an atom of oxygen from the oxidising agent.

..... [1]

(d) $\text{CH}_3\text{CH}_2\text{COOH}$ reacts with an unsaturated alcohol **R** to form unsaturated ester **S**.

(i) State the **type** of reaction that forms **S**.

..... [1]

(ii) The infrared spectrum of **S** is shown in Fig. 4.2.




Fig. 4.2

Three absorptions in the infrared spectrum in Fig. 4.2 confirm that **S** is an ester and is unsaturated.

- Write **1**, **2** or **3** on Fig. 4.2 against each of these three absorptions.
- Complete Table 4.1 to show which bond is responsible for each absorption that you have identified in Fig. 4.2.

Table 4.1

absorption	1	2	3
bond responsible			

[3]

Table 4.2

bond	functional groups containing the bond	characteristic infrared absorption range (in wavenumbers)/cm ⁻¹
C–O	hydroxy, ester	1040–1300
C=C	aromatic compound, alkene	1500–1680
C=O	amide carbonyl, carboxyl ester	1640–1690 1670–1740 1710–1750
C≡N	nitrile	2200–2250
C–H	alkane	2850–2950
N–H	amine, amide	3300–3500
O–H	carboxyl hydroxy	2500–3000 3200–3650

(iii) The mass spectrum of **S** shows the following peaks.

Table 4.3

peak	relative abundance
M ⁺	4.7
[M+1] ⁺	0.31

Use Table 4.3 to calculate the number of carbon atoms in **S**.

number of carbon atoms in **S** = [1]

[Total: 16]

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.02 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)

The Periodic Table of Elements

Group																			
1	2													13	14	15	16	17	18

- 3** Half-fill the 250 cm³ beaker with water and place it on a tripod and gauze. Heat the water until boiling then switch off your Bunsen burner. This will be your hot water bath for use in **(b)(i)**. Start **(a)(i)** while the water is heating.
- (a)** **FA 4** is an aqueous solution containing three ions. One ion is **not** listed in the Qualitative analysis notes. **FA 5** is an aqueous solution containing two ions. The cation in **FA 5** is listed in the Qualitative analysis notes. The anions in both **FA 4** and **FA 5** contain sulfur.
- (i)** Carry out the following tests using a 1 cm depth of either **FA 4** or **FA 5** in a test-tube. Record your observations in Table 3.1.

Table 3.1

<i>test</i>	<i>observations</i>	
	FA 4	FA 5
Test 1 Add aqueous sodium hydroxide.		
Test 2 Add aqueous ammonia.		
Test 3 Add aqueous sodium carbonate.		
Test 4 Add aqueous barium chloride or aqueous barium nitrate.		
Test 5 Add acidified aqueous potassium manganate(VII).		

[5]

- (ii) From your observations in (a)(i), deduce the identities of the ions present in **FA 4** and **FA 5**. Give the formula of each ion in Table 3.2. If you are unable to identify an ion write 'unknown'.

Table 3.2

<i>ions present</i>	FA 4	FA 5
cations		
anions		

[3]

- (b) (i) Both **FA 6** and **FA 7** are one of propan-1-ol, propan-2-ol, methanoic acid or ethanoic acid. You will carry out tests to investigate the identities of **FA 6** and **FA 7**. For each test use a 1 cm depth of **FA 6** or **FA 7** in a test-tube. Record your observations in Table 3.3.

Table 3.3

<i>test</i>	<i>observations</i>	
	FA 6	FA 7
Test 1 Add a few drops of acidified aqueous potassium manganate(VII), then		
place the test-tube in the hot water bath.		
Test 2 Add a 2 cm depth of aqueous iodine followed by drops of aqueous sodium hydroxide until the colour just disappears.		
If no reaction is visible, place the test-tube in the hot water bath.		
Test 3 Add aqueous sodium carbonate.		

[3]

- (ii) From your observations in (b)(i) deduce the identities of **FA 6** and **FA 7**. Give reasons for your answers.

FA 6 is

reasons

.....

FA 7 is

reasons

.....

[2]

- (c) A student carrying out a similar set of tests identifies two unknown compounds to be ethanol and propanoic acid.

Write an equation for the reaction between these two compounds.

..... [1]

[Total: 14]

Qualitative analysis notes

1 Reactions of cations

cation	reaction with	
	NaOH(aq)	NH ₃ (aq)
aluminium, Al ³⁺ (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH ₄ ⁺ (aq)	no ppt. ammonia produced on warming	–
barium, Ba ²⁺ (aq)	faint white ppt. is observed unless [Ba ²⁺ (aq)] is very low	no ppt.
calcium, Ca ²⁺ (aq)	white ppt. unless [Ca ²⁺ (aq)] is very low	no ppt.
chromium(III), Cr ³⁺ (aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu ²⁺ (aq)	pale blue ppt. insoluble in excess	pale blue ppt. soluble in excess giving dark blue solution
iron(II), Fe ²⁺ (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe ³⁺ (aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
magnesium, Mg ²⁺ (aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn ²⁺ (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn ²⁺ (aq)	white ppt. soluble in excess	white ppt. soluble in excess

2 Reactions of anions

anion	reaction
carbonate, CO ₃ ²⁻	CO ₂ liberated by dilute acids
chloride, Cl ⁻ (aq)	gives white ppt. with Ag ⁺ (aq) (soluble in NH ₃ (aq))
bromide, Br ⁻ (aq)	gives cream/off-white ppt. with Ag ⁺ (aq) (partially soluble in NH ₃ (aq))
iodide, I ⁻ (aq)	gives pale yellow ppt. with Ag ⁺ (aq) (insoluble in NH ₃ (aq))
nitrate, NO ₃ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and Al foil
nitrite, NO ₂ ⁻ (aq)	NH ₃ liberated on heating with OH ⁻ (aq) and Al foil; decolourises acidified aqueous KMnO ₄
sulfate, SO ₄ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (insoluble in excess dilute strong acids); gives white ppt. with high [Ca ²⁺ (aq)]
sulfite, SO ₃ ²⁻ (aq)	gives white ppt. with Ba ²⁺ (aq) (soluble in excess dilute strong acids); decolourises acidified aqueous KMnO ₄
thiosulfate, S ₂ O ₃ ²⁻ (aq)	gives off-white/pale yellow ppt. slowly with H ⁺

3 Tests for gases

gas	test and test result
ammonia, NH_3	turns damp red litmus paper blue
carbon dioxide, CO_2	gives a white ppt. with limewater
hydrogen, H_2	'pops' with a lighted splint
oxygen, O_2	relights a glowing splint