


3 What contains 9.03×10^{23} oxygen atoms?

- A** 0.25 mol aluminium oxide
- B** 0.75 mol sulfur dioxide
- C** 1.5 mol sulfur trioxide
- D** 3.0 mol water

4 Methane and steam react to produce hydrogen.





0.80 g of methane and 1.35 g of steam react. One of the reactants is used up.

Which volume of hydrogen, measured at room conditions, will be produced?

- A** 1.80 dm³ **B** 3.60 dm³ **C** 4.80 dm³ **D** 7.20 dm³

17 Three equations are listed. x , y and z are all whole numbers.



Which equations can be balanced if $x = 4$ and $z = 2$?

- A** 1 and 3 **B** 1 only **C** 2 only **D** 3 only

40 A sample of magnesium contains the isotopes ^{24}Mg , ^{25}Mg and ^{26}Mg only.

The percentage abundance of ^{25}Mg and ^{26}Mg is the same.

The relative atomic mass of magnesium in the sample is 24.3.

What is the percentage abundance of ^{24}Mg ?

- A** 10% **B** 20% **C** 60% **D** 80%

Important values, constants and standards

molar gas constant	$R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
Faraday constant	$F = 9.65 \times 10^4 \text{ C mol}^{-1}$
Avogadro constant	$L = 6.02 \times 10^{23} \text{ mol}^{-1}$
electronic charge	$e = -1.60 \times 10^{-19} \text{ C}$
molar volume of gas	$V_m = 22.4 \text{ dm}^3 \text{ mol}^{-1}$ at s.t.p. (101 kPa and 273 K) $V_m = 24.0 \text{ dm}^3 \text{ mol}^{-1}$ at room conditions
ionic product of water	$K_w = 1.00 \times 10^{-14} \text{ mol}^2 \text{ dm}^{-6}$ (at 298 K (25 °C))
specific heat capacity of water	$c = 4.18 \text{ kJ kg}^{-1} \text{ K}^{-1}$ (4.18 J g ⁻¹ K ⁻¹)

- 3 Cycloalkanes show similar chemical properties to alkanes but have the same empirical formula as alkenes.

(a) Define empirical formula.

.....
..... [1]

- (b) Cyclopentane, C_5H_{10} , has four cyclic structural isomers. One of these isomers is **C**, shown in Fig. 3.1.

Complete Fig. 3.1 to show two other cyclic structural isomers of C_5H_{10} .


cyclopentane 			C 
--	--	--	---

Fig. 3.1

[2]

- (c) Cyclopentane reacts with Cl_2 in the presence of ultraviolet light to form C_5H_9Cl .

(i) The reaction is initiated by the bond fission of Cl_2 .

State the type of bond fission shown in the initiation step.

..... [1]


(ii) Complete the equations to show the two propagation steps that follow the initiation step.

propagation 1 $C_5H_{10} + \dots \rightarrow C_5H_9\cdot + \dots$

propagation 2 $C_5H_9\cdot + \dots \rightarrow \dots$

[2]

(iii) The final step is shown.



Give the name for this step in the reaction.

..... [1]

(d) Fig. 3.2 shows a reaction cycle involving cyclopentane, cyclopentene and $\text{C}_5\text{H}_9\text{Cl}$.




Fig. 3.2

(i) Identify a suitable reagent for reaction 3.

..... [1]

(ii) Use the data in Fig. 3.2 and in Table 3.1 to calculate the enthalpy change of reaction 2, ΔH_2 .

Table 3.1

compound	enthalpy change of combustion, $\Delta H_c / \text{kJ mol}^{-1}$
	-3292
	-3115
H_2	-286

⑤ $\Delta H_2 = \dots\dots\dots \text{kJ mol}^{-1}$ [2]

(e) Cyclopentene, C_5H_8 , reacts with hot concentrated acidified $KMnO_4$ to form compound **W**, $C_5H_8O_4$.

(i) Draw the structure of **W**.

[1]

(ii) The infrared spectrum of **W** is shown in Fig. 3.3.




Fig. 3.3

Identify **two** absorptions in the infrared spectrum of **W** that would **not** be present in the infrared spectrum of cyclopentene.

- Write **1** or **2** on Fig. 3.3 against each of these two absorptions.
- Complete Table 3.2 to show which bond is responsible for each absorption that you have identified in Fig. 3.3.

Table 3.2

absorption	1	2
bond responsible		

[2]


Table 3.3

bond	functional groups containing the bond	characteristic infrared absorption range (in wavenumbers)/cm ⁻¹
C–O	hydroxy, ester	1040–1300
C=C	aromatic compound, alkene	1500–1680
C=O	amide carbonyl, carboxyl ester	1640–1690 1670–1740 1710–1750
C≡N	nitrile	2200–2250
C–H	alkane	2850–2950
N–H	amine, amide	3300–3500
O–H	carboxyl hydroxy	2500–3000 3200–3650

[Total: 13]

- 1 Hydrated ethanedioic acid is a diprotic acid with the formula $(\text{COOH})_2 \cdot x\text{H}_2\text{O}$ where x is an integer.

Ethanedioic acid reacts with manganate(VII) ions when heated.



You will determine the value of x in $(\text{COOH})_2 \cdot x\text{H}_2\text{O}$ by titrating a solution containing ethanedioic acid with manganate(VII) ions.

- **FA 1** is 6.20 g dm^{-3} aqueous ethanedioic acid, $(\text{COOH})_2 \cdot x\text{H}_2\text{O}$.
- **FA 2** is $0.0200 \text{ mol dm}^{-3}$ potassium manganate(VII), KMnO_4 .
- **FA 3** is 1.0 mol dm^{-3} sulfuric acid, H_2SO_4 .

(a) Method

- Fill the burette with **FA 2**.
- Pipette 25.0 cm^3 of **FA 1** into a conical flask.
- Use the measuring cylinder to add approximately 20 cm^3 of **FA 3** to the conical flask.
- Place the conical flask on a tripod and gauze and heat carefully until the temperature of the solution is approximately 70°C .
- Remove the flame.
- Carefully lift the hot conical flask and place it on the white tile under the burette.
- Add **FA 2** drop-wise for the first $2\text{--}3 \text{ cm}^3$. Any initial pink colouring may take several seconds to disappear.
- If the reaction mixture turns brown, reheat it to about 70°C . If the brown colour disappears, continue the titration. If the brown colour remains, discard the contents of the flask and begin a new titration.
- The end-point is reached when a permanent pale pink colour is formed.
- Perform a rough titration with **FA 2**. Record your burette readings in the space below.

The rough titre is cm^3 .

- Carry out as many titrations as you think necessary to obtain consistent results.
- Make sure any recorded results show the precision of your practical work.
- Record all your burette readings and the volume of **FA 2** added in each accurate titration.

Results

I	
II	
III	
IV	
V	
VI	
VII	

[7]

- (b) From your accurate titration results, calculate a suitable mean value to be used in your calculations.

Show clearly how you obtained this value.

25.0 cm³ of **FA 1** required cm³ of **FA 2**. [1]

(c) Calculations

(i) Give your answers to **(c)(ii)**, **(c)(iii)** and **(c)(iv)** to the appropriate number of significant figures. [1]

(ii) Calculate the amount, in mol, of manganate(VII) ions, MnO_4^- , in the volume of **FA 2** calculated in **(b)**.

amount of MnO_4^- = mol [1]

(iii) Calculate the amount, in mol, of ethanedioic acid that reacts with the manganate(VII) ions in **(c)(ii)**.

amount of $(\text{COOH})_2$ = mol

Hence calculate the concentration, in mol dm^{-3} , of ethanedioic acid in **FA 1**.

concentration of $(\text{COOH})_2$ = mol dm^{-3}
[2]

(iv) Calculate the relative molecular mass, M_r , of the ethanedioic acid in **FA 1**.

M_r = [1]

(v) Calculate the value of **x** in $(\text{COOH})_2 \cdot x\text{H}_2\text{O}$.

Show your working.

x = [1]

(d) Explain why it is necessary to add **FA 3** in each titration.

.....

..... [1]

[Total: 15]