2.2 Matrices (eigenvalues and eigenvectors)

Name: _____ Class: _____ Date: _____ **Total: 34 marks**

KEY WORDS characteristic equation • eigenvalue • eigenvector • matrices

Objective

Build the skills to answer exam questions on **matrices** – solving $A\mathbf{x} = \mathbf{b}$, the **characteristic equation**, **eigenvalues** and **eigenvectors**, and diagonalisation.

You must be able to:

- find eigenvalues from $\det(A - \lambda I) = 0$
- find an eigenvector for each eigenvalue
- write $A = QDQ^{-1}$ and use it (e.g. for A^n)

1 Key knowledge and worked examples

Eigenvalues & eigenvectors



Multiplying an eigenvector by A only stretches it (by λ); a general vector changes direction



Eigenvectors are the directions a matrix leaves unturned

$A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$: $\det(A - \lambda I) = (2-\lambda)^2 - 1 = 0 \Rightarrow \lambda = 1$ or 3. Eigenvector for $\lambda = 3$: $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$; for $\lambda = 1$: $\begin{pmatrix} 1 \\ -1 \end{pmatrix}$.

Eigenvalues, eigenvectors, and what diagonalisation buys you

An **eigenvector** of A is a non-zero $\mathbf{v}$ that A only stretches: $A\mathbf{v} = \lambda\mathbf{v}$, where λ is the **eigenvalue**. The direction survives the transformation; only the length changes.

The method, which is the same every time:

1. **Characteristic equation:** $\det(A - \lambda I) = 0$. For a 2×2 this is $\lambda^2 - (\operatorname{tr} A)\lambda + \det A = 0$, which is worth knowing as a check.
2. **Solve** for the eigenvalues.
3. **For each** λ, solve $(A - \lambda I)\mathbf{v} = \mathbf{0}$. The rows are dependent by construction, so you get a **family** – quote any convenient member, and if the rows are NOT dependent you have made an arithmetic slip.
4. **Diagonalise:** put the eigenvectors in the columns of P and the matching eigenvalues down the diagonal of D, in the **same order**. Then $A = PDP^{-1}$.

Why it is worth doing: $A^n = PD^nP^{-1}$, and D^n is just each diagonal entry raised to n. Computing A^{10} directly is ten matrix multiplications; this is one.

The Cayley–Hamilton theorem says a matrix satisfies its own characteristic equation. For a 2×2 with $\lambda^2 - t\lambda + d = 0$, that gives $A^2 = tA - dI$ – which reduces any power of A, and rearranges to $A^{-1} = \frac{1}{d}(tI - A)$.

Worked example. For $A = \begin{pmatrix} 4 & 1 \\ 2 & 3 \end{pmatrix}$, find the eigenvalues and one eigenvector for each.

1. $\operatorname{tr} A = 7$, $\det A = 10$, so $\lambda^2 - 7\lambda + 10 = 0$.
2. $(\lambda - 5)(\lambda - 2) = 0$, so $\lambda = 5$ or $\lambda = 2$.
3. For $\lambda = 5$: $(A - 5I)\mathbf{v} = \begin{pmatrix} -1 & 1 \\ 2 & -2 \end{pmatrix}\mathbf{v} = \mathbf{0}$ gives $v_1 = v_2$, so $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$.

4. For $\lambda = 2$: $\begin{pmatrix} 2 & 1 \\ 2 & 1 \end{pmatrix} \mathbf{v} = \mathbf{0}$ gives $2v_1 = -v_2$, so $\begin{pmatrix} 1 \\ -2 \end{pmatrix}$.

■ **Key knowledge: Using diagonalisation for powers**

Match each eigenvector column to the corresponding diagonal eigenvalue. Verify $AP = PD$ before computing a matrix power.

■ **Worked example: Using diagonalisation for powers**

For $A = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$, take $P = \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}$ and $D = \operatorname{diag}(3, 1)$. Since $P^{-1} = P/2$, $A^3 = PD^3P^{-1} = \begin{pmatrix} 14 & 13 \\ 13 & 14 \end{pmatrix}$. Direct multiplication of $A^2 = \begin{pmatrix} 5 & 4 \\ 4 & 5 \end{pmatrix}$ by A confirms it.

2 Practice

2.1 Write down the characteristic equation of a matrix A. [1]

2.2 Find the eigenvalues of $\begin{pmatrix} 4 & 0 \\ 0 & -2 \end{pmatrix}$. [1]

2.3 Write down the characteristic equation of $\begin{pmatrix} 3 & 2 \\ 1 & 4 \end{pmatrix}$ using its trace and determinant. [2]

2.4 State what P and D contain in the diagonalisation $A = PDP^{-1}$. [2]

2.5 Use the Cayley–Hamilton theorem to express A^2 in terms of A and I for a matrix with $\lambda^2 - 6\lambda + 8 = 0$. [2]

3 Exam-style questions

3.1 A vector $\mathbf{v}$ is an eigenvector of A with eigenvalue λ when: [1]

☐ **A** $A\mathbf{v} = \mathbf{0}$
B $A\mathbf{v} = \lambda\mathbf{v}$
C $\det A = \lambda$
D $A = \lambda I$

3.2 The matrix $A = \begin{pmatrix} 3 & 1 \\ 0 & 2 \end{pmatrix}$.

(a) Find the eigenvalues. [2]

(b) Find an eigenvector for each eigenvalue. [4]

3.3 The matrix $M = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$ has eigenvalues 1 and 3. Using $M = QDQ^{-1}$, find M^2 and verify by direct multiplication. [5]

3.4 The matrix $A = \begin{pmatrix} 5 & 2 \\ 2 & 2 \end{pmatrix}$.

(a) Show that the characteristic equation of A is $\lambda^2 - 7\lambda + 6 = 0$. [3]

(b) Find the eigenvalues of A. [2]

(c) Find an eigenvector corresponding to each eigenvalue. [4]

(d) Write down matrices P and D such that $A = PDP^{-1}$. [2]

(e) Explain why this makes A^n easy to compute, and state the form of A^n. [3]