

- 1 (a) Use standard results from the list of formulae (MF19) to find $\sum_{r=1}^n (8r^3 + 12r^2 + 4r + 3)$ in terms of n , simplifying your answer. [3]

①

- (b) Show that

$$\frac{1}{r^4} - \frac{1}{(r+1)^4} = \frac{4r^3 + 6r^2 + 4r + 1}{r^4(r+1)^4}$$

and hence use the method of differences to find $\sum_{r=1}^n \frac{4r^3 + 6r^2 + 4r + 1}{r^4(r+1)^4}$. [5]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.

- (c) Deduce the value of $\sum_{r=1}^{\infty} \frac{4r^3 + 6r^2 + 4r + 1}{r^4(r+1)^4}$. [1]

2

- 1 (a) Use standard results from the list of formulae (MF19) to find $\sum_{r=1}^n (r^5 - r)$ in terms of n , fully factorising your answer. [3]

[3]

- (b) Express $\frac{r+3}{r^3-r}$ in the form $\frac{A}{r-1} + \frac{B}{r} + \frac{C}{r+1}$, where A , B and C are constants to be determined, and hence use the method of differences to find $\sum_{r=2}^n \frac{r+3}{r^3-r}$. [5]

[5]

- (c) Deduce the value of $\sum_{r=2}^{\infty} \frac{r+3}{r^3-r}$. [1]

[1]

(b) Express $\frac{2}{(r+1)(r+2)(r+3)}$ in partial fractions and hence use the method of differences to find

$$\sum_{r=1}^n \frac{2}{(r+1)(r+2)(r+3)}. \quad [5]$$

[3]

(c) State the value of $\sum_{r=1}^{\infty} \frac{2}{(r+1)(r+2)(r+3)}$. [1]

1 (a) Use standard results from the list of formulae (MF19) to show that

$$\sum_{r=1}^n (2-3r)(5-3r) = an^3 + bn^2 + cn,$$

where a, b and c are integers to be determined.

[3]

(b) Use the method of differences to find $\sum_{r=1}^n \frac{1}{(2-3r)(5-3r)}$ in terms of n .

[4]

(c) Deduce the value of $\sum_{r=1}^{\infty} \frac{1}{(2-3r)(5-3r)}$.

[1]

4 Let $w_r = r(r+1)(r+2) \dots (r+9)$.

(a) Show that

$w_{r+1} - w_r = 10(r+1)(r+2) \dots (r+9).$ [2]

(b) Given that $u_r = (r+1)(r+2) \dots (r+9)$, find $\sum_{r=1}^n u_r$ in terms of n . [3]

(c) Given that $v_r = x^{w_{r+1}} - x^{w_r}$, find the set of values of x for which the infinite series

$v_1 + v_2 + v_3 + \dots$

is convergent and give the sum to infinity when this exists. [3]