


2

When the angular speed is increased to ω_2 , the angle between the upward vertical and the normal reaction of the surface on P becomes θ_2 , where $\tan \theta_2 = \frac{4}{3}$.

Find the ratio $\frac{\omega_1}{\omega_2}$.

[4]

This image shows a full page of white paper with horizontal dotted lines. The lines are evenly spaced and run across the width of the page, providing a guide for handwriting practice. There are no margins, text, or other markings on the page.



A particle P of mass m is attached to one end of a light inextensible string of length a . The other end of the string is attached to a fixed point O . Initially P is held with the string taut and making an angle of 60° with the upward vertical through O . The particle P is projected perpendicular to the string in a downwards direction with speed $\sqrt{17ag}$. It then starts to move along a circular path in a vertical plane with centre O (see diagram). At the lowest point of its path, vertically below O , the particle P collides with a stationary particle Q .

- (a)** Find, in terms of a and g , an expression for the speed of P immediately before the collision with Q . [2]

This image shows a full page of blank handwriting practice paper. It features multiple sets of horizontal lines. Each set consists of a solid top line, a dashed middle line, and a solid bottom line, providing a guide for letter height and placement. The paper is otherwise empty, with no text or markings other than the pre-printed lines.


As a result of the collision, P rebounds and moves back along a circular path with centre O . The string becomes slack when P reaches the point on the circle vertically above O .

(b) Find, in terms of a and g , an expression for the speed of P immediately after the collision with Q .
[3]

The mass of particle Q is km and the collision between P and Q is perfectly elastic.

(c) Find the value of k .
[3]


1



Find the tension in string AP in terms of m , g , l and ω .

[4]

Handwriting practice lines (dotted lines on a grid background).



A fixed smooth sphere with radius a and centre O rests on horizontal ground. A particle is projected horizontally from the highest point, A , of the sphere with speed u . The particle begins to move in a vertical circle along the surface of the sphere. The particle loses contact with the sphere at the point B , where the angle AOB is θ .

After leaving the surface of the sphere, the particle moves freely under gravity before striking the horizontal ground with speed $3u$ at an angle β to the horizontal (see diagram).

Find the value of β .

[8]

[illegible]

This image shows a full page of white paper with horizontal dashed lines, typical of primary school writing paper. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.




Name: Further Mathematics: **9231 Mathematics - Further November 2025 Question Paper** 34

Name: Further Mathematics: **9231 Mathematics - Further November 2025 Question Paper** 34

Name: Further Mathematics: **9231 Mathematics - Further November 2025 Question Paper** 34

Name: Further Mathematics: **9231 Mathematics - Further November 2025 Question Paper** 34



A fixed smooth spherical shell has centre O and radius a . A particle of mass m moves in complete vertical circles on the smooth inner surface of the shell, where the plane of the circular motion is vertical and passes through O . The particle has speed v when it is at point A , where OA makes an angle θ with the upward vertical through O , and $\cos \theta = \frac{1}{18}$ (see diagram).


(a) Show that $v \geq \frac{1}{3}\sqrt{26ag}$.

[5]

It is given that $v = \frac{1}{3}\sqrt{26ag}$.

- (b)** Find, in terms of m and g , an expression for the greatest possible value of the normal reaction between the shell and the particle. [3]

[illegible]



A rough horizontal disc, centre O , rotates with constant angular speed $\omega \text{ rad s}^{-1}$. A particle P of mass 1.6 kg lies on the disc at a distance 1.5 m from O , and is attached to a point A vertically above O by a light elastic string. The string has natural length 2 m , modulus of elasticity 32 N and makes an angle α with the vertical OA (see diagram). Particle P moves in a horizontal circle also at a constant angular speed $\omega \text{ rad s}^{-1}$. Particle P is on the point of slipping in the direction OP . The coefficient of friction between the particle and the disc is 0.5 .

- (a)** Given that the tension in the string is 8 N, show that $\sin \alpha = 0.6$.


[2]

[illegible]

(b)

[5]

11



A fixed hollow sphere has radius a and centre O . The points A , B and C lie on the inner surface of the sphere with OA and BC horizontal. A portion of the sphere has been removed by a horizontal cut through points B and C at a vertical distance ka above the centre of the sphere, where k is a positive constant and $k < 1$. The points O , A , B and C all lie in the same vertical plane. OB makes an angle θ with the upward vertical through O (see diagram).

A particle P of mass m is free to move on the smooth inner surface of the sphere. The particle P is projected vertically downwards from A with speed u and begins to move in a vertical circle.

- (a)** In the case where $u = \sqrt{\frac{6}{5}ga}$, the reaction on P at B is half the reaction on P at A .

Find the value of k .

[5]

Handwriting practice lines (dotted lines on a solid background) for the number 12.

Additional page


If you use the following page to complete the answer to any question, the question number must be clearly shown.

[illegible]

- 1

Given that $\tan \theta = \frac{4}{3}$,

[3]



A hollow cylinder of radius r is fixed with its axis horizontal. Points A , B and O are in the same vertical plane perpendicular to the axis of the cylinder, with A and B on the smooth inner surface and O on the axis. OA and OB make angles 90° and α respectively with the upward vertical through O , with A and B on opposite sides of the vertical. A particle of mass m is projected vertically downwards from point A with speed $\sqrt{\frac{3}{2}rg}$ and moves in a vertical circle inside the cylinder (see diagram). The particle loses contact with the cylinder at point B .

- (a)** Find the value of α .

[4]

[illegible]

[4]

17