

1.1 Quadratics

Name: _____ Class: _____ Date: _____

Total: 16 marks

Objective

Build the skills to answer exam questions on **quadratics** 二次式—completing the square, the **discriminant** 判别式, solving equations and inequalities, and simultaneous equations.

You must be able to:


- write $ax^2 + bx + c$ in completed-square form $a(x + p)^2 + q$
- use $b^2 - 4ac$ to decide the number of real roots
- solve quadratic equations and inequalities
- solve a linear-quadratic pair by substitution

1 Worked examples


■ Completing the square & the discriminant

Write $2x^2 - 12x + 5$ as $a(x + p)^2 + q$.

$$2x^2 - 12x + 5 = 2(x^2 - 6x) + 5 = 2((x - 3)^2 - 9) + 5 = 2(x - 3)^2 - 13.$$



The sign of $b^2 - 4ac$ decides how many times the parabola meets the x-axis



2 Practice

2.1 Express $x^2 + 8x + 3$ in the form $(x + p)^2 + q$. [2]

2.2 State the condition on the discriminant for a quadratic to have **no** real roots. [1]

3 Exam-style questions

3.1 The equation $2x^2 - 4x + c = 0$ has a repeated root. The value of c is: [1]

- A 0
- B 1
- C 2
- D 4

3.2 Express $3x^2 + 12x + 7$ in the form $a(x + b)^2 + c$ and hence write down the least value

of the expression. [4]

3.3 Find the set of values of k for which the line $y = kx - 2$ meets the curve $y = x^2 - 3x + 2$ at two distinct points. [4]

3.4 Solve the simultaneous equations $y = 2x - 1$ and $x^2 + y^2 = 2$. [4]

4 Go further

You are now ready for the real exam questions on this subtopic. Open the **1.1 Quadratics** past-paper sheet in the Library, or try these in **Practice mode**:

- 9709/11 N25 —Q1 (quadratics)
- 9709/11 N25 —Q4 (discriminant / curve meets line)
- 9709/12 N25 —Q1 (quadratics)

Solutions

2.1 $(x + 4)^2 - 13$.

2.2 $b^2 - 4ac < 0$.

3.1 C. Repeated root needs $b^2 - 4ac = 0$: $16 - 8c = 0 \Rightarrow c = 2$.

3.2 $3(x^2 + 4x) + 7 = 3((x + 2)^2 - 4) + 7 = 3(x + 2)^2 - 5$; least value = -5 (when $x = -2$).

3.3 set equal: $x^2 - 3x + 2 = kx - 2 \Rightarrow x^2 - (3 + k)x + 4 = 0$; two points need $b^2 - 4ac > 0$: $(3 + k)^2 - 16 > 0$; $(k + 7)(k - 1) > 0$; $k < -7$ or $k > 1$.

3.4 substitute: $x^2 + (2x - 1)^2 = 2 \Rightarrow 5x^2 - 4x - 1 = 0$; $(5x + 1)(x - 1) = 0 \Rightarrow x = -\frac{1}{5}$ or $x = 1$; $y = 2x - 1$ gives $(1, 1)$ and $(-\frac{1}{5}, -\frac{7}{5})$.