

# 4 Mechanics 1 – Part 2

Name: \_\_\_\_\_ Class: \_\_\_\_\_ Date: \_\_\_\_\_

## Vocabulary 词汇

Write each word three times. 把每个单词抄写三遍。

| English                         | 中文     | Write it 3 times (English) |       |       |
|---------------------------------|--------|----------------------------|-------|-------|
| Newton's laws of motion         | 牛顿运动定律 | _____                      | _____ | _____ |
| linear momentum                 | 动量     | _____                      | _____ | _____ |
| conservation of linear momentum | 动量守恒   | _____                      | _____ | _____ |
| work done                       | 功      | _____                      | _____ | _____ |
| kinetic energy                  | 动能     | _____                      | _____ | _____ |
| gravitational potential energy  | 重力势能   | _____                      | _____ | _____ |
| power                           | 功率     | _____                      | _____ | _____ |

## Recall


In **Part 1** you met forces, equilibrium, weight, friction, and the suvat equations. Part 2 adds **Newton's second law**, **momentum**, and **work, energy and power**.

## New ideas

### ■ 1 Newton's second law

**Newton's laws of motion** 牛顿运动定律 connect force and acceleration. The key one is: resultant force = mass  $\times$  acceleration,

$$F = ma.$$



$$F = ma$$

*Worked example.* A resultant force of 12 N acts on a 4 kg mass:  $a = \frac{F}{m} = \frac{12}{4} = 3 \text{ m s}^{-2}$ .


**Micro-check.** A 6 N resultant force acts on a 2 kg mass. Find the acceleration. [1]

---

## ■ 2 Momentum

The **linear momentum** 动量 of a body is mass  $\times$  velocity (a vector). In a collision with no outside force, the total momentum is unchanged —the **conservation of linear momentum** 动量守恒:

$$m_1 u_1 + m_2 u_2 = m_1 v_1 + m_2 v_2.$$



*Worked example.* A 2 kg body at  $3 \text{ m s}^{-1}$  hits a stationary 1 kg body and they stick:  $2(3) = 3v$ , so  $v = 2 \text{ m s}^{-1}$ .

**Micro-check.** Find the momentum of a 5 kg body moving at  $4 \text{ m s}^{-1}$ . [1]

---

## ■ 3 Work, energy and power

- **work done** 功 by a force is force  $\times$  distance moved:  $W = Fd$  (in joules);
- **kinetic energy** 动能  $= \frac{1}{2}mv^2$ ; **gravitational potential energy** 重力势能  $= mgh$ ;
- **power** 功率 is the rate of doing work; for a driving force,  $P = Fv$ .

*Worked example.* A car engine works at 12 kW while moving at  $20 \text{ m s}^{-1}$ : driving force  $F = \frac{P}{v} = \frac{12000}{20} = 600 \text{ N}$ .

**Micro-check.** Find the kinetic energy of a 2 kg body moving at  $3 \text{ m s}^{-1}$ . [1]

---

**Watch out:**  $F = ma$  uses the *resultant* force. In a collision with no outside force, *momentum* ( $mv$ ) is conserved:  $m_1u_1 + m_2u_2 = m_1v_1 + m_2v_2$ . Power for a driving force is  $P = Fv$ .

## Practice

---

**4.1** A resultant force of 20 N acts on a 5 kg mass. Find the acceleration. [2]

**4.2** Find the momentum of a 1500 kg car moving at  $12 \text{ m s}^{-1}$ . [2]

**4.3** A 3 kg body at  $4 \text{ m s}^{-1}$  hits a stationary 1 kg body and they stick together. Find their common speed. [2]

**4.4** Find the kinetic energy of a 4 kg body moving at  $5 \text{ m s}^{-1}$ . [2]

**4.5** A force pulls a load at  $8 \text{ m s}^{-1}$  with a driving force of 50 N. Find the power ( $P =$

$Fv$ ).

[2]

**4.6 Challenge.** A 2 kg trolley moving at  $5 \text{ m s}^{-1}$  collides with a stationary 3 kg trolley and they stick together. Find their common speed, and the kinetic energy lost in the collision.

[3]