

A-Level Physics

Name: _____ Score: _____

1. The mass-energy relation is:

☐ $E = m^2c$

☐ $E = \frac{1}{2}mc^2$

☐ $E = mc^2$

☐ $E = mc$

2. Energy is released when nuclei move:

☐ toward the iron peak (higher binding energy per nucleon)☐ nowhere – energy is never released☐ away from the iron peak☐ to lower nucleon number, always

3. Select **all** the true statements about radioactive decay.

☐ the rate is unaffected by temperature☐ it is random☐ you can predict exactly when one nucleus will decay☐ it is spontaneous (no trigger needed)

Tick every correct option.

4. A mass change of 1 u corresponds to about how many MeV?

_____ MeV

5. An iron-56 nucleus can release energy by either fusion or fission.

☐ True ☐ False

6. What is the evidence that radioactive decay is random?

☐ the half-life is always the same☐ the count rate fluctuates from one interval to the next☐ the activity falls over time☐ heating the source does not change the rate

7. In alpha decay the nucleon number falls by 4 and the proton number falls by _____.

8. Fusion needs very high temperatures to overcome the electrostatic repulsion between nuclei.

☐ True ☐ False

9. The activity of a radioactive source is:

☐ $A = \lambda N$ (decays per second, in Bq)

☐ $A = \frac{N}{\lambda}$

☐ $A = \lambda + N$

☐ $A = \frac{\lambda}{N}$

10. A nucleus has _____ mass than its separate protons and neutrons added together.

A-Level Physics

1. $E = mc^2$

Mass and energy are equivalent: $E = mc^2$, so a mass change Δm releases $c^2\Delta m$.

2. **toward the iron peak (higher binding energy per nucleon)**

Climbing the B/A curve toward iron means the products are more tightly bound, so energy is released.

3. **the rate is unaffected by temperature; it is random; it is spontaneous (no trigger needed)**

It is spontaneous and random, and the rate does not depend on conditions. You can only give the probability of a single decay.

4. **931 MeV**

1 u = 1.66×10^{-27} kg gives $\Delta E = c^2\Delta m \approx 931$ MeV.

5. **False**

It sits at the peak, so every move takes it DOWN the curve and would need energy put in. This is why fusion inside a star stops at iron.

6. **the count rate fluctuates from one interval to the next**

A falling activity shows decay happens; a constant half-life shows the probability is fixed; heating shows it is SPONTANEOUS. Only the fluctuation shows randomness.

7. **2**

An alpha particle is a helium-4 nucleus, 2 protons and 2 neutrons. Beta-minus leaves A unchanged and raises Z by 1, and the antineutrino must be written too.

8. **True**

The nuclei must get close enough for the strong force to act, despite repelling – hence the millions of kelvin in stars.

9. **$A = \lambda N$ (decays per second, in Bq)**

Activity = decay constant $\times$ number of undecayed nuclei, measured in becquerel.

10. **less**

That missing mass (the mass defect) was released as energy when the nucleus formed.