

A-Level Physics

Name: _____ Score: _____

1. A piezo-electric crystal:
 - ☐ conducts current in one direction only
 - ☐ only emits light
 - ☐ stores charge like a battery
 - ☐ changes shape with a p.d., and makes an e.m.f. when squeezed
2. X-rays are produced when fast electrons:
 - ☐ are heated gently
 - ☐ hit a metal target and slow down sharply
 - ☐ travel through glass
 - ☐ pass through a magnet
3. A PET tracer is:
 - ☐ an ultrasound gel
 - ☐ a strong magnet
 - ☐ a positron (β^+) emitter taken up by active tissue
 - ☐ an X-ray source
4. An echo returns after 2.0×10^{-4} s; the speed of sound is $1500 \frac{\text{m}}{\text{s}}$. How deep is the boundary?
_____ m
5. Most of the electrons' kinetic energy in an X-ray tube becomes heat, not X-rays.
 - ☐ True ☐ False
6. Why is a tracer's half-life chosen as a compromise? Select **all** that apply.
 - ☐ long enough that it has not decayed before reaching the tissue
 - ☐ long enough to stay in the body for later scans
 - ☐ short enough that the patient's dose after the scan is small
 - ☐ short enough that the activity is high during the scan

Tick every correct option.
7. An ultrasound pulse returns $60 \mu\text{s}$ after transmission, with a speed of sound of $1500 \frac{\text{m}}{\text{s}}$. How deep is the boundary, in cm?

8. The minimum X-ray wavelength from a tube at accelerating voltage V is:

☐ $hceV$

☐ $\frac{eV}{hc}$

☐ $\frac{h}{eV}$

☐ $\frac{hc}{eV}$

9. When the emitted positron meets an electron, they:

☐ form a stable atom

☐ do nothing

☐ bounce off each other

☐ annihilate into two photons

10. A higher ultrasound frequency has which effects? Select **all** that apply.

☐ stronger attenuation, so less penetration

☐ better resolution

☐ a larger reflection at every boundary

☐ a shorter wavelength

Tick every correct option.

A-Level Physics

1. **changes shape with a p.d., and makes an e.m.f. when squeezed**

Both effects let one crystal send ultrasound (apply p.d.) and detect it (read the e.m.f.).

2. **hit a metal target and slow down sharply**

The sudden deceleration on the target gives out X-ray photons (Bremsstrahlung).

3. **a positron (β^+) emitter taken up by active tissue**

It is a β^+ emitter on a molecule (like glucose) that active tissue absorbs more of.

4. **0.15 m**

$$d = \frac{ct}{2} = \frac{1500 \times 2.0 \times 10^{-4}}{2} = 0.15 \text{ m (the pulse travels there and back).}$$

5. **True**

Only a small fraction becomes X-rays – which is why the target must be cooled.

6. **long enough that it has not decayed before reaching the tissue; short enough that the patient's dose after the scan is small; short enough that the activity is high during the scan**

Fluorine-18 has 110 minutes; oxygen-15 has only 2 and must be made on the spot. Leaving activity in the patient afterwards is the thing to avoid, not to aim for.

7. **4.5**

$$d = \frac{ct}{2} = 4.5 \text{ cm. Halve it: the pulse travelled there AND back. Forgetting the 2 gives 9.0 cm.}$$

8. $\frac{hc}{eV}$

$$\text{From } hf_{\max} = eV \text{ and } c = f\lambda: \lambda_{\min} = \frac{hc}{eV}.$$

9. **annihilate into two photons**

Matter and antimatter annihilate: their mass turns into two photons.

10. **stronger attenuation, so less penetration; better resolution; a shorter wavelength**

Reflection is set by the impedance mismatch, not the frequency. The frequency choice is a resolution-against-penetration compromise, so a deep organ is scanned at a lower frequency.