A-LEVEL PHYSICS • EXERCISE SHEET

6.2 Elastic and plastic behaviour

Name: ____________ Class: ________ Date: __________ **Total: 14 marks**

KEY WORDS elastic potential energy 弹性势能 • thermal energy 热能 • elastic 弹性 • extension 伸长量 • limit of proportionality 比例极限

Objective

Build the skills to answer exam questions on **elastic and plastic behaviour** — deformation types and elastic potential energy.

You must be able to:

- use **elastic deformation**, **plastic deformation** and **elastic limit**
- find work done from the **area under a force–extension graph**
- use $E_\mathrm{P} = \frac{1}{2}Fx = \frac{1}{2}kx^2$

1 Worked examples

Study these first. Each one shows the method for a question type used later — follow the steps and you can do the Practice and Exam-style questions yourself.

Elastic and plastic

- **Elastic deformation:** the object returns to its original length when the load is removed.
- **Plastic deformation:** a **permanent** extension remains after the load is removed (past the **elastic limit**).

Energy stored = area under the graph

The **work done** stretching a material is the **area under** its force–extension graph. For a Hooke's-law material this is a triangle:





Elastic and plastic behaviour on a force-extension graph

$$E_\mathrm{P} = \frac{1}{2}Fx = \frac{1}{2}kx^2 = \frac{F^2}{2k}.$$

Worked example. $k = 50\ \mathrm{N\ m^{-1}}$, $x = 0.20$ m: $E_\mathrm{P} = \frac{1}{2}(50)(0.20)^2 = 1.0$ J.

Loading and unloading

Past the elastic limit, the **unloading** line is parallel to the first straight part but shifted right (a permanent extension). The area **between** the loading and unloading lines is energy turned into **thermal energy** in the material.

2 Practice

Now apply the methods above.

2.1 Define **elastic deformation** and **plastic deformation**. [2]

2.2 State what the **area under** a force–extension graph represents. [1]

2.3 A spring of spring constant 50 N m^{-1} is stretched by 0.20 m. Find the **elastic potential energy** stored. [2]

2.4 State what remains when a material stretched **past its elastic limit** has the load removed. [1]

3 Exam-style questions

3.1 The elastic potential energy in a spring of spring constant k stretched by x is: [1]

- **A** kx
- **B** $\frac{1}{2}kx$
- **C** $\frac{1}{2}kx^2$
- **D** kx^2

3.2 A force of 8.0 N stretches a wire by 2.0 mm, within its limit of proportionality. Calculate the **elastic potential energy** stored. [2]

3.3 Explain the difference between **elastic** and **plastic** deformation. [2]

3.4 A material is loaded **past** its elastic limit and then unloaded. Explain why the loading and unloading lines differ, and state what the **area between** them represents. [3]

Solutions

Each answer shows the steps, then the **result**.

2.1

- **elastic**: the object **returns** to its original length when the load is removed
- **plastic**: a **permanent** extension remains

2.2

- the **work done** on the material (elastic potential energy stored, within the elastic region)

2.3

- $E_{\mathrm{P}} = \frac{1}{2}kx^2$

$$E_{\mathrm{P}} = \tfrac{1}{2}(50\ \mathrm{N\ m^{-1}})(0.20\ \mathrm{m})^2 = \mathbf{1.0}\ \mathrm{J}$$

2.4

- a **permanent extension** (it does not return to its original length)

3.1 C

- elastic potential energy stored in a spring is $\frac{1}{2}kx^2$

3.2

- $E_{\mathrm{P}} = \frac{1}{2}Fx$ (because $F = kx$)

$$E_{\mathrm{P}} = \tfrac{1}{2}(8.0\ \mathrm{N})(2.0 \times 10^{-3}\ \mathrm{m}) = \mathbf{8.0 \times 10^{-3}}\ \mathrm{J}$$

3.3

- elastic deformation is **reversible** — the material returns to its original shape
- plastic deformation is **permanent** — it stays deformed after the load is removed

3.4

- past the elastic limit some deformation is permanent, so unloading returns to a **non-zero** (permanent) extension
- the unloading line is parallel to but shifted from the loading line
- the area between them is the **energy transferred to thermal energy** in the material