

23.2 Radioactive decay

Name: _____ Class: _____ Date: _____

Total: 32 marks

KEY WORDS activity 活度 • half-life 半衰期 • spontaneous 自发 • decay constant 衰变常数
• random 随机

Objective

Build the skills to answer exam questions on **radioactive decay**.

You must be able to:



- describe decay as **spontaneous** and **random**
- use **activity** $A = \lambda N$ and $\lambda = 0.693/t_{1/2}$
- use the exponential decay $x = x_0 e^{-\lambda t}$ and **half-life**

1 Worked examples

Study these first. Each one shows the method for a question type used later — follow the steps and you can do the Practice and Exam-style questions yourself.

■ Random and exponential

Decay is **spontaneous** (no trigger) and **random** (the count rate fluctuates). The number of undecayed nuclei falls **exponentially**, halving every **half-life**:



$$N = N_0 e^{-\lambda t} \quad T_{1/2} = \ln 2 / \lambda$$

The exponential radioactive decay curve

$$A = \lambda N, \quad \lambda = \frac{0.693}{t_{1/2}}, \quad N = N_0 e^{-\lambda t}.$$

- activity** A — decays per second (becquerel, Bq); **decay constant** λ — probability of decay per second.

The same exponential law applies to the **activity** and the **count rate**: $A = A_0 e^{-\lambda t}$.

■ Finding the time — take logs

For a **whole number** of half-lives, just halve repeatedly (after n half-lives a fraction $(\frac{1}{2})^n$ remains). For **any** time or fraction, take the natural log of $N = N_0 e^{-\lambda t}$:

$$\ln \frac{N}{N_0} = -\lambda t \quad \Rightarrow \quad t = -\frac{1}{\lambda} \ln \frac{N}{N_0}.$$

Worked example. A source has half-life 8.0 days, so $\lambda = \frac{0.693}{8.0} = 0.0866 \text{ day}^{-1}$.

The time for its activity to fall to 20% is $t = -\frac{1}{0.0866} \ln(0.20) = 18.6 \text{ days}$.

2 Practice

Now apply the methods above.

2.1 State **two** features of radioactive decay.

[2]

2.2 Define **activity** and the **decay constant**. [2]

2.3 Define **half-life**. [1]

2.4 Write the equation linking **activity**, **decay constant** and **number** of nuclei. [1]

3 Exam-style questions

3.1 The **half-life** is the time for: [1]

- | | |
|---|---|
| A | B |
| C | D |
- A** all the nuclei to decay
B half the (undecayed) nuclei to decay
C the activity to reach exactly zero
D the decay constant to halve

3.2 A source has a decay constant of 0.040 s^{-1} . Calculate its **half-life**. [2]

3.3 A sample has 2.0×10^{20} undecayed nuclei and a decay constant of $1.5 \times 10^{-9} \text{ s}^{-1}$. Calculate its **activity**. [2]

3.4 A radioactive isotope has a half-life of 6.0 hours. What **fraction** of the original nuclei remains after 18 hours? [2]

3.5 A radioactive source has a half-life of 8.0 days. Calculate the **time** for its activity to fall to 20% of its initial value. [4]

3.6 Oxygen-15 is used as a tracer in medical imaging. It decays by β^+ emission with a half-life of 122 s. A freshly prepared sample contains 4.8×10^{14} nuclei.

(a) **State** what is meant by a **tracer**. [2]

(b) The decay is ${}^{15}_{8}\text{O} \rightarrow {}^P_Q\text{N} + {}^R_S\text{Z}$. **State** the values of P , Q , R and S , and **name** the particle Z. [3]

(c) **Define** the **activity** of a radioactive sample. [1]

(d) **Determine** the decay constant of oxygen-15. Give a **unit** with your answer. [2]

(e) **Determine** the initial activity of the sample. [2]

(f) **Determine** the activity 10 minutes after preparation. [3]

(g) **Suggest** why a tracer with a half-life of a few **days** would be unsuitable for this use. [2]