

7.5 Polarisation

Name: _____ Class: _____ Date: _____

Total: 20 marks

KEY WORDS polarising filter 偏振片 • polarisation 偏振 • transverse 横波
• plane-polarised 平面偏振 • wave 波

Objective

Build the skills to answer exam questions on **polarisation** 偏振 and **Malus's law**.

You must be able to:

- explain that only **transverse** 横波 waves can be polarised
- use **Malus's law** $I = I_0 \cos^2 \theta$, and work **backwards** to find the angle
- sketch how the transmitted intensity varies as a filter is **rotated**
- link transmitted **intensity** to **amplitude** 振幅 using $I \propto A^2$

1 Worked examples


Study these first. Each one shows the method for a question type used later — follow the steps and you can do the Practice and Exam-style questions yourself.

■ Polarisation

Polarisation makes a transverse wave oscillate in **one plane** only. Only **transverse** waves can be polarised (their oscillation is perpendicular to the travel direction, so different planes exist). **Longitudinal** 纵波 waves such as sound **cannot** — so polarisation is a **test for a transverse wave**.

■ Malus's law

Plane-polarised light of intensity I_0 meets a filter whose **transmission axis** 透射轴 is at angle θ to the plane of polarisation. The transmitted intensity is:



$$I = I_0 \cos^2 \theta.$$


- $\theta = 0^\circ$: $I = I_0$ (all passes); $\theta = 90^\circ$: $I = 0$ (blocked); $\theta = 45^\circ$: $I = \frac{1}{2}I_0$; $\theta = 60^\circ$: $I = \frac{1}{4}I_0$.

Worked example (forward). $I_0 = 12 \text{ W m}^{-2}$, $\theta = 30^\circ$: $I = 12 \cos^2 30^\circ = 12 \times 0.75 = 9.0 \text{ W m}^{-2}$.

Worked example (backwards). For what θ is the intensity **halved**? Set $\cos^2 \theta = 0.5$, so $\cos \theta = \sqrt{0.5}$ and $\theta = 45^\circ$.

■ Rotating the filter

If the angle θ between the axis and the polarisation is changed (by **rotating** the filter), the transmitted intensity traces a $\cos^2$ curve — maximum I_0 at $0^\circ, 180^\circ, 360^\circ$ and zero at $90^\circ, 270^\circ$:



The "starts perpendicular" case. If the axis **begins perpendicular** to the polarisation and is then rotated by an angle ϕ , the angle in the formula is $\theta = 90^\circ - \phi$. So $I = I_0 \cos^2(90^\circ - \phi) = I_0 \sin^2 \phi$: the intensity **starts at zero**, rises to I_0 after $\phi = 90^\circ$, and so on.

■ Intensity and amplitude

Intensity is proportional to amplitude squared, $I \propto A^2$, so for a wave through a filter:

$$\frac{A}{A_0} = \sqrt{\frac{I}{I_0}}.$$

Worked example. If a filter transmits $I = 0.25 I_0$, the amplitude is $A = A_0 \sqrt{0.25} = 0.50 A_0$ — the amplitude falls to **half**, even though the intensity falls to a **quarter**.

■ Two filters in series

Apply Malus's law **once per filter**; the second angle is measured from the **first** filter's axis. Two **crossed** filters (90° apart) transmit **nothing**.

2 Practice

Now apply the methods above.

2.1 State what **polarisation** means. [1]

2.2 State why **sound** cannot be polarised. [1]

2.3 Plane-polarised light meets a filter whose axis is at 60° to its plane. State the **fraction** of intensity transmitted. [1]

2.4 A filter transmits **half** the incident intensity. State the angle θ between its axis and the plane of polarisation. [1]

2.5 A filter transmits $I = 0.25 I_0$. State the transmitted amplitude as a fraction of A_0 . [1]

3 Exam-style questions

3.1 Which of these **can** be polarised? [1]

- | | |
|---|---|
| A | B |
| C | D |
- A sound waves
B all electromagnetic waves
C longitudinal waves
D all of these

3.2 Plane-polarised light of intensity 12 W m^{-2} meets a filter whose axis is at 30° to the plane of polarisation. Calculate the transmitted intensity. [2]

3.3 Plane-polarised light passes through **two** filters. The first axis is at 30° to the plane of polarisation; the second is a further 30° beyond the first. The light reaching the first filter has intensity 8.0 W m^{-2} . Calculate the intensity emerging from the **second** filter. [3]

3.4 A photographer looks at a shop window through a polarising filter. The light reflected from the glass is plane-polarised, of intensity I_0 , and the filter's transmission axis starts **parallel** to the plane of polarisation. The filter is then rotated through 360° about the viewing direction.

(a) On the axes, sketch how the transmitted intensity I varies with the rotation angle ϕ . [2]

(b) Determine the smallest rotation ϕ that **halves** the transmitted intensity. [2]

(c) At that angle, find the transmitted **amplitude** as a fraction of the incident amplitude A_0 , and explain why the photographer rotates the filter to cut the glare. [3]

3.5 Explain why the ability to **polarise** a wave shows that the wave is **transverse**. [2]