

1(a)	Value of P in range 5.00 to 15.00 s, with unit seen somewhere
	Evidence of repeated measurement of P
1(b)	Six sets of readings of z (different values) and P with correct overall trend and without help scores 5 marks, five sets scores 4 marks etc. Correct trend is P decreases as z increases.
	Range: at least one reading of $z \leq 10$ g and at least one reading of $z \geq 70$ g
	Column headings: Each column heading must contain a quantity and a unit where appropriate. The presentation of quantity and unit must conform to accepted scientific convention e.g. $1/\sqrt{M} (\text{g}^{-1/2})$
	Consistency: All raw values of P must be given to the nearest 0.01 s or all to the nearest 0.1 s.
	Significant figures: All values of $1/\sqrt{M}$ must be given to 3 s.f. or 4 s.f.
	Calculation: $1/\sqrt{M}$ calculated correctly
1(c)(i)	Axes: Axes must be labelled with the correct quantities . Scales must be chosen so that the plotted points occupy at least half the graph grid in both the x and y directions. Scale markings are no more than 2 cm (one large square) apart. Sensible scales must be used, no awkward scales (e.g. 3:10 or fractions).
	Plotting of points: All observations in the table must be plotted on the grid. Diameter of plotted points must be $\leq$ half a small square. Points must be plotted to an accuracy of half a small square in both x and y directions.
	Quality: General trend of points must be negative. All points in the table must be plotted (at least 5) for this mark to be awarded. It must be possible to draw a straight line that is within $\pm 0.001 \text{ g}^{-1/2}$ on the $1/\sqrt{M}$ axis of all plotted points.
1(c)(ii)	Line of best fit: 'Best fit' is judged by the balance of all points on the grid (at least 5 points) about the candidate's line. There must be an even distribution of points either side of the line along the full length. Lines must not be kinked or thicker than half a square. Some candidates may choose to identify an anomalous point. If they identify one point as anomalous (e.g. by circling or labelling) then this point is to be disregarded when judging the line of best fit. There must be at least 5 points left after the anomalous point is disregarded.

1(c)(iii)	Gradient: Gradient sign on answer line consistent with graph drawn. The hypotenuse of the triangle used should be greater than half the length of the drawn line. Both read offs must be accurate to half a small square in both the x and y directions. Method of calculation must be correct, not $\Delta x / \Delta y$.
1(c)(iii)	y-intercept: Either Intercept read directly from the graph, with read off at $x = 0$, accurate to half a small square in y direction. Or Correct read off from a point on the line is substituted into $y = mx + c$ or an equivalent expression. Read-off accurate to half a small square in both x and y directions
1(d)	Value of a = candidate's gradient value and Value of b = candidate's y-intercept value. The values must not be written as fractions or given to only one significant figure.
	Units for a and b correct and consistent with readings (e.g. $\text{g}^{\frac{1}{2}}\text{s}^{-1}$ for a and s^{-1} for b)

4(a)(i)	temperature = $-273.15\text{ }^{\circ}\text{C}$
4(a)(ii)	temperature = 0 K
4(b)(i)	gas is ideal
4(b)(ii)	$pV = NkT$ $N = 270 / (8.0 \times 10^{-21})$ $= 3.4 \times 10^{22}$
4(b)(iii)	$n = (3.4 \times 10^{22}) / (6.02 \times 10^{23})$ $= 0.056\text{ mol}$
4(c)	$\frac{1}{2} m \langle c^2 \rangle = (3 / 2) kT$ $\frac{1}{2} \times m \times 1900^2 = 1.5 \times 8.0 \times 10^{-21}$ $(m = 6.65 \times 10^{-27}\text{ kg})$ $m = (6.65 \times 10^{-27}) / (1.66 \times 10^{-27})$ $= 4.0\text{ u}$

5(a)	work done per unit charge
	work (done) moving positive charge from infinity (to the point)
5(b)(i)	potential (due to proton) = $(1.60 \times 10^{-19}) / (4\pi \times 8.85 \times 10^{-12} \times 10 \times 10^{-12})$
	or
	potential (due to electron) = $(-1.60 \times 10^{-19}) / (4\pi \times 8.85 \times 10^{-12} \times 110 \times 10^{-12})$
	$V = [(1.60 \times 10^{-19}) / (4\pi \times 8.85 \times 10^{-12})] \times [(10^{-1} - 110^{-1}) \times 10^{12}] = 130 \text{ V}$
5(b)(ii)	$V = [(1.60 \times 10^{-19}) / (4\pi \times 8.85 \times 10^{-12})] \times [(30^{-1} - 90^{-1}) \times 10^{12}]$
	= (+) 32 V
5(b)(iii)	cross drawn midway between the electron and the proton
5(b)(iv)	line from (10, +130) to (110, -130)
	curve getting shallower until $x = 60 \text{ pm}$, crossing $V = 0$ at (60, 0) and then getting steeper after $x = 60 \text{ pm}$
	curve passing through (30, ± 32) and (90, ± 32)
6(a)	series charges: $Q_s = Q_1 = Q_2$
	series p.d.s: $V_s = V_1 + V_2$
	parallel charges: $Q_s = Q_1 + Q_2$
	parallel p.d.s: $V_s = V_1 = V_2$
6(b)(i)	$E = \frac{1}{2} CV^2$
	p.d. = $[(2 \times 19 \times 10^{-3}) / (470 \times 10^{-6})]^{1/2}$
	= 9.0 V
6(b)(ii)	$E = Q^2 / 2C$ or $C = Q / V$
	$Q = (19 \times 10^{-3} \times 2 \times 470 \times 10^{-6})^{1/2}$
	or
	$Q = 470 \times 10^{-6} \times 9.0$
	$Q = 4.2 \times 10^{-3} \text{ C}$
6(b)(iii)	total charge unchanged
	total capacitance = $(470 + 180) \times 10^{-6} \text{ (F)}$
	$E = Q^2 / 2C = (4.23 \times 10^{-3})^2 / (2 \times 650 \times 10^{-6}) \text{ (= 0.014 J)}$
	$E = 14 \text{ mJ}$

7(a)	(induced) e.m.f. is (directly) proportional to rate
	of change of (magnetic) flux (linkage)
7(b)(i)	flux = e.m.f. $\times$ time
	flux = 0.54×15 = 8.1 Wb
7(b)(ii)	$\phi = BA$
	area = $8.1 / (38 \times 10^{-6})$ = $2.1 \times 10^5 \text{ m}^2$
7(b)(iii)	area = speed $\times$ time $\times$ width
	$v = (2.1 \times 10^5) / (15 \times 68)$ = 210 m s^{-1}
7(b)(iv)	opposing force (due to current in wings) must be backwards
	from Fleming's left-hand rule, current (in wings) must be from Q to P
	current is from – to + inside an e.m.f. source so P is at higher potential

2(a)	work done on / by system
	thermal energy supplied to / removed from system
2(b)(i)	no thermal energy transferred to / from system (due to lack of time)
	work is done on the gas to compress it / to decrease its volume
	internal energy increases so temperature increases
2(b)(ii)	(during vaporisation) molecular separation increases
	(heating causes) potential energy of molecules to increase
	kinetic energy of molecules unchanged so temperature unchanged