

- 5 A girl throws a ball vertically upwards. It takes a time of 3.20 s to return to her hand.

Assume air resistance is negligible.

What is the initial speed with which the ball is thrown?

- A** 3.07 ms^{-1} **B** 7.85 ms^{-1} **C** 15.7 ms^{-1} **D** 31.4 ms^{-1}

A-Level Physics: **w21 12**

- 7 A stone is projected vertically upwards from the ground at an initial speed of 15 ms^{-1} . Air resistance is negligible.

What is the maximum height reached by the stone?

- A** 0.76 m **B** 11 m **C** 23 m **D** 110 m

A-Level Physics: **w24 12**


- 5 A stone falls vertically from rest. Air resistance is negligible.

What is the speed of the stone when it has fallen through a distance of 0.40 m?


- A** 2.0 ms^{-1} **B** 2.8 ms^{-1} **C** 3.9 ms^{-1} **D** 7.8 ms^{-1}

A-Level Physics: **s21 12**

- 6 Which graph shows the variation with time t of the velocity v of an object falling vertically downwards in a vacuum?



- 5 The diagram shows a laboratory experiment in which a feather falls from rest in a long evacuated vertical tube of length L .



The feather takes time T to fall from the top to the bottom of the tube.

How far does the feather fall from the top of the tube in time $0.50T$?


- A** $0.13L$ **B** $0.25L$ **C** $0.38L$ **D** $0.50L$

A-Level Physics: **s21 13**

- 7 A steel ball is dropped from rest from a height h above the ground. The ball hits the ground after a time t .

This is repeated for a number of different heights.

The graph shows the variation of h with t^2 for the ball.



The gradient of the graph is G .

Which expression gives the acceleration of the ball?

- A** $\frac{G}{2}$ **B** G **C** $2G$ **D** G^2

- 6 The water surface in a deep well is 78.0 m below the top of the well. A person at the top of the well drops a heavy stone down the well.


Air resistance is negligible. The speed of sound in the air is 330 m s^{-1} .

What is the time interval between the person dropping the stone and hearing it hitting the water?

- A 3.75 s B 3.99 s C 4.19 s D 4.22 s

- 6 A rock is launched vertically upwards from the surface of the Earth and an identical rock is launched vertically upwards from the surface of Mars. Each rock is launched with an initial velocity of 12 m s^{-1} .

Each rock then reaches its maximum height above the surface before returning back down to the surface. The velocity–time graph for each rock is shown. In both cases, air resistance is negligible.



What is the difference in the maximum heights of the two rocks?

- A 12 m B 15 m C 19 m D 24 m

- 6 On the Earth, an object takes time T_E to fall from rest through a vertical distance h .

On the Moon, the same object takes time T_M to fall from rest through the same vertical distance h .

The ratio $\frac{\text{acceleration of free fall on the Earth}}{\text{acceleration of free fall on the Moon}}$ is equal to 6.

Air resistance is negligible for the object on the Earth and on the Moon.

What is the ratio $\frac{T_E}{T_M}$?

- A $\frac{1}{6}$ B $\frac{1}{\sqrt{6}}$ C $\sqrt{6}$ D 6

- 3 A man standing on a wall throws a small ball vertically upwards with a velocity of 5.6 m s^{-1} . The ball leaves his hand when it is at a height of 3.1 m above the ground, as shown in Fig. 3.1.




Fig. 3.1 (not to scale)

Assume that air resistance is negligible.

- (a) Show that the ball reaches a maximum height above the ground of 4.7 m.

[2]

- (b) The man does not catch the ball as it falls.

Calculate the time taken for the ball to fall from its maximum height to the ground.

time taken = s [2]

- (c) The ball leaves the man's hand at time $t = 0$ and hits the ground at time $t = T$.

On Fig. 3.2, sketch a graph to show the variation of the velocity v of the ball with time t from $t = 0$ to $t = T$. Numerical values of v and t are not required. Assume that v is positive in the upward direction.




Fig. 3.2

[3]

- (d) State what is represented by the gradient of the graph in (c).

..... [1]

- (e) The man now throws a second ball with the same velocity and from the same height as the first ball. The mass of the second ball is greater than that of the first ball. Assume that air resistance is still negligible.

For the first and second balls, compare:

- (i) the magnitudes of their accelerations

..... [1]

- (ii) the speeds with which they hit the ground.

..... [1]

[Total: 10]