


Ideal Gases

A2 Physics ■ Topic 15

A-Level Physics



What we will cover

- 1 The mole 15.1
- 2 The equation of state 15.2
- 3 Kinetic theory 15.3
- 4 Deriving $pV = \frac{1}{3}Nm\langle c^2 \rangle$ 15.3.1
- 5 Recap



1

The mole

Sheet 15.1 – what you need to know

You must be able to:

- state what the **Avogadro constant** is and use $N = nN_A$
- find the mass of one molecule from the molar mass
- predict how p , n , $\langle c^2 \rangle$ and U change between samples

What the exam asks

“State what is meant by the Avogadro constant.” [1]

“Complete Table 3.1 by giving expressions, in terms of some or all of N , m , T , V and the constants.” [4]

“Sketch the variation with thermodynamic temperature of the r.m.s. speed of the molecules.” [2]

Practise on exercise sheet 15.1, then try the real exam questions on past-paper sheet 15.1.

The mole

Amount of substance 物质的量 n is measured in **moles** 摩尔. One mole holds

Avogadro constant 阿伏伽德罗常数 particles


– **atoms** 原子 for a **monatomic** 单原子 gas like helium, molecules for O_2 :

$$N_A = 6.02 \times 10^{23} \text{ mol}^{-1}$$

Number of particles $N = nN_A$. Molar mass in grams is the relative formula mass – 18 g of water is one mole.

Count particles with moles, not with 10^{23} on every line. n is the quantity the gas laws actually use.

One mole, counted



1 mole

$= 6.02 \times 10^{23}$
particles (N_A)

A mole is 6.02×10^{23} particles – the number chosen so that a mole of a substance weighs its relative molecular mass in grams.

One mole is 6.02×10^{23} particles – whatever the particle is. The molar mass in grams equals the relative formula mass, so 18 g of water is exactly one mole.

The Avogadro constant, and the mass of one molecule

“State what is meant by the Avogadro constant.” [1]

The number of particles in **one mole** of a substance:

$$N_A = 6.02 \times 10^{23} \text{ mol}^{-1}.$$

$$N = nN_A, \quad m = \frac{M}{N_A}$$

M : molar mass

Nitrogen has a molar mass of $0.028 \text{ kg mol}^{-1}$. Find the mass of one molecule.

One mole has mass M and holds N_A molecules:

$$\begin{aligned} m &= \frac{M}{N_A} = \frac{0.028 \text{ kg mol}^{-1}}{6.02 \times 10^{23} \text{ mol}^{-1}} \\ &= 4.7 \times 10^{-26} \text{ kg} \end{aligned}$$

Convert the molar mass to kg mol^{-1} first:
 28 g mol^{-1} is $0.028 \text{ kg mol}^{-1}$.

Change the sample: what happens to each quantity?

Sample X: N molecules of mass m in volume V at T . Sample Y: $2N$ molecules of mass $2m$ in $2V$, at the same T .

quantity	Y compared with X	because
pressure p	the same	$p = NkT/V$: N and V both double
amount n	doubled	$n = N/N_A$
$\langle c^2 \rangle$	halved	$\langle c^2 \rangle = 3kT/m$, from $\frac{1}{2}m\langle c^2 \rangle = \frac{3}{2}kT$
internal energy U	doubled	$U = \frac{3}{2}NkT$: twice the molecules, same mean KE

Write the formula with every symbol, then ask which symbols change. The mean KE depends on T **only**.

2

The equation of state

Sheet 15.2 – what you need to know

You must be able to:

- use $pV = nRT = NkT$, with T in kelvin
- define an **ideal gas**
- read N or n from the gradient of a pV – T graph
- sketch a cycle of changes on p – V axes

What the exam asks

“Identify the constant B .” [1]

“Show that the r.m.s. speed of molecules of this gas is approximately 500 m s^{-1} .” [3]

“Sketch the variation with volume of pressure for the gas as the gas undergoes the three changes.” [3]

Practise on exercise sheet 15.2, then try the real exam questions on past-paper sheet 15.2.

The equation of state


An **ideal gas** 理想气体 obeys $pV \propto T$ exactly:

$$pV = nRT \quad \text{or} \quad pV = NkT$$

- the **molar gas constant** 摩尔气体常数 $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$ – per **mole**
- the **Boltzmann constant** 玻尔兹曼常数 $k = 1.38 \times 10^{-23} \text{ J K}^{-1}$ – per **molecule**
- $N = nN_A$, so $k = R/N_A$

Match the form to your **amount**: moles $\rightarrow nRT$, molecules $\rightarrow NkT$. SI throughout – Pa, m^3 , and **kelvin**.

The equation of state, in detail



gas: p , V , T , n moles

$$pV = nRT$$

Every quantity in $pV = nRT$ is in SI units: pressure in Pa, volume in m^3 , and temperature in [kelvin](#) – never $^{\circ}\text{C}$.

Worked example – how many moles?

A 0.020 m^3 cylinder holds gas at 27°C and $2.0 \times 10^5 \text{ Pa}$. Find n . ($R = 8.31$)

First to kelvin: $T = 27 + 273 = 300 \text{ K}$.

$$n = \frac{pV}{RT} = \frac{(2.0 \times 10^5)(0.020)}{8.31 \times 300} \approx 1.6 \text{ mol}$$

The kelvin conversion is the **first** line, every time. 27 in place of 300 would inflate n by a factor of **11**.

A fixed mass changing state

For a fixed amount of gas:

$$\frac{p_1 V_1}{T_1} = \frac{p_2 V_2}{T_2}$$

Hold one quantity fixed and a named law drops out.

Boyle

T fixed: pV constant

Charles

p fixed: V/T constant


pressure law

V fixed: p/T constant

$pV \propto T$ holds **only** in kelvin – °C here is the classic lost mark.

Boyle's law


At constant temperature pV is constant, so p against $1/V$ is a **straight line through the origin** – the usual way an exam asks you to verify it.



Charles's law and absolute zero

At constant pressure the volume climbs **linearly** with temperature. Extrapolate the line back to **zero volume** and it meets $\approx -273^{\circ}\text{C}$.

Every gas points at the **same** intercept – which is what makes absolute zero a property of **temperature**, not of any one gas.



$V \propto T$ · Charles: extrapolates to absolute zero

Worked example – heating at constant pressure


**A fixed mass at 300 K fills 0.50 m^3 .
Heat it to 450 K at constant pressure.
Find the new volume.**

p is fixed, so V/T is constant:

$$V_2 = V_1 \frac{T_2}{T_1} = 0.50 \times \frac{450}{300} = 0.75 \text{ m}^3$$

Cancel what is **constant** before substituting – here p drops out and only the temperature **ratio** matters.

Reading a pV - T graph



A straight line **through the origin**: the gas is ideal.


The line passes through (300 K, 750 J). Find the number of molecules N and the amount n .

$$pV = NkT \Rightarrow N = \frac{\text{gradient}}{k} = \frac{750 \text{ J}/300 \text{ K}}{1.38 \times 10^{-23} \text{ J K}^{-1}} = 1.8 \times 10^{23}$$

$$n = \frac{N}{N_A} = \frac{1.8 \times 10^{23}}{6.02 \times 10^{23} \text{ mol}^{-1}} = 0.30 \text{ mol}$$

Ideal gas: obeys $pV \propto T$ at **all** pressures, volumes and temperatures.

A gas taken round a cycle



Fixed mass, so $\frac{pV}{T}$ is constant. Start at X:
 $2p, V, T$.

- $X \rightarrow Y$, constant V : p halves, so $T \rightarrow T/2$ – a **vertical** line
- $Y \rightarrow Z$, constant p : V doubles, so $T/2 \rightarrow T$ – a **horizontal** line
- $Z \rightarrow X$, constant T : pV stays $2pV$ – a **curve**, $p \propto 1/V$

Constant T is never a straight line on p - V axes: it is the curve $pV = \text{constant}$.

3

Kinetic theory

Sheet 15.3 – what you need to know

You must be able to:

- state the **assumptions** of the kinetic model
- use $pV = \frac{1}{3}Nm\langle c^2 \rangle$ and $p = \frac{1}{3}\rho\langle c^2 \rangle$
- link mean KE to temperature: $\frac{1}{2}m\langle c^2 \rangle = \frac{3}{2}kT$
- explain why $U \propto T$ for an ideal gas

What the exam asks

“Explain why the internal energy of an ideal gas is proportional to its thermodynamic temperature.”
[2]

“Suggest why a gas at the surface of a star is unlikely to be ideal.”
[2]

“Determine, to three significant figures, the temperature of the surface of the star.” [2]


Practise on exercise sheet 15.3, then try the real exam questions on past-paper sheet 15.3.

Kinetic theory: the assumptions

- 1 many identical molecules in **random motion** 无规则运动
- 2 their own volume is negligible
- 3 collisions are brief; straight lines between
- 4 no **intermolecular** 分子间 forces except in a collision
- 5 collisions are **elastic** 弹性碰撞
- 6 Newton's laws apply

Elastic is what keeps a gas from **cooling itself**. The model fails at **high** p (volume matters) and **low** T (forces matter).

Assumptions



1. many identical molecules, in continuous **random motion**
2. their own volume is **tiny** next to the container
3. straight lines between hits - intermolecular forces **ignored**
4. collisions are **elastic**: no kinetic energy is lost


poor at very high pressure (volume matters) or very low temperature (forces matter)

The key assumptions of the kinetic theory of an ideal gas

Root-mean-square speed

r.m.s. speed 均方根 $= \sqrt{\langle c^2 \rangle}$:
square each speed, average, then
square-root. It sits slightly **above** the
mean speed – squaring weights the
fast molecules more.

The gas needs $\langle c^2 \rangle$, not $\langle c \rangle$, because **energy** goes as c^2 . Averaging the speeds first gives the wrong answer.



read the name **backwards**: root of the **mean** of the **squares**

Why it matters under water




A diver's air is compressed: at 10 m the pressure is doubled, so a given mass of gas takes half the volume.

The spread of speeds

Molecules do not share one speed – they occupy a **distribution**. Raise the temperature and the curve **broadens** and shifts to faster speeds.

The area is fixed (the molecules still all exist), so a broader curve must also be a **lower** one.



higher T : peak $\rightarrow$ higher speeds, flatter curve

The r.m.s. speed from pressure and density

Divide $pV = \frac{1}{3}Nm\langle c^2 \rangle$ by V . The mass per unit volume Nm/V is the density ρ :

$$p = \frac{1}{3}\rho\langle c^2 \rangle$$

No N , no m , no T needed: pressure and density are enough.

Air at 1.0×10^5 Pa has a density of 1.2 kg m^{-3} . Show that the r.m.s. speed of its molecules is about 500 m s^{-1} .

$$\begin{aligned} c_{\text{rms}} &= \sqrt{\langle c^2 \rangle} = \sqrt{\frac{3p}{\rho}} \\ &= \sqrt{\frac{3 \times 1.0 \times 10^5 \text{ Pa}}{1.2 \text{ kg m}^{-3}}} = 500 \text{ m s}^{-1} \end{aligned}$$

Average translational KE

The **average translational kinetic energy** 平均平动动能, from two expressions for the same pV – the second using the **mean square** 均方 speed $\langle c^2 \rangle$:


$$pV = NkT, \quad pV = \frac{1}{3}Nm\langle c^2 \rangle$$

Set equal, cancel N , multiply by $\frac{3}{2}$:

$$\langle E_k \rangle = \frac{1}{2}m\langle c^2 \rangle = \frac{3}{2}kT$$


The average molecular KE depends on **temperature alone** – not the gas, not the pressure. Squeezing a gas at fixed T gives more wall hits, **not** faster molecules.

Average translational KE, in detail



Mean molecular kinetic energy depends on T alone. Squeezing a gas at fixed T gives more wall hits, not faster molecules.

Consequences



$\langle c^2 \rangle$ **doubles**

$$c_{\text{r.m.s.}} \times \sqrt{2}$$

same T



lighter $\Rightarrow$ faster on average

- **Mean KE** of a molecule $\propto T$ (absolute).
- Same T : same mean KE, so the **lighter** gas has the larger $\langle c^2 \rangle$.

$$E_k = \frac{3}{2} NkT = \frac{3}{2} nRT$$

total translational KE of N molecules

Worked example – r.m.s. speed of oxygen

Find $c_{\text{r.m.s.}}$ for O_2 at 300 K.

$$m = 5.3 \times 10^{-26} \text{ kg}, \quad k = 1.38 \times 10^{-23} \text{ J K}^{-1}$$

$$\text{From } \frac{1}{2} m \langle c^2 \rangle = \frac{3}{2} kT: \quad \langle c^2 \rangle = 3kT/m$$

$$c_{\text{r.m.s.}} = \sqrt{\frac{3(1.38 \times 10^{-23})(300)}{5.3 \times 10^{-26}}}$$

$$\approx 480 \text{ m s}^{-1}$$

Faster than a jet – yet the gas goes nowhere, because the **directions** are random. Lighter gas, same $T \Rightarrow$ **faster**: hydrogen outruns oxygen in the same room.

Worked example – a temperature from an r.m.s. speed

Helium atoms (mass $6.6 \times 10^{-27} \text{ kg}$) have an r.m.s. speed of 1500 m s^{-1} . Find the temperature of the gas.

Mean KE of one atom: $\frac{1}{2} m \langle c^2 \rangle = \frac{3}{2} kT$

$$T = \frac{m \langle c^2 \rangle}{3k} = \frac{6.6 \times 10^{-27} \text{ kg} \times (1500 \text{ m s}^{-1})^2}{3 \times 1.38 \times 10^{-23} \text{ J K}^{-1}} \\ = 360 \text{ K}$$


Square the r.m.s. speed to get $\langle c^2 \rangle$. Then $T \propto \langle c^2 \rangle$: doubling T multiplies c_{rms} by $\sqrt{2}$, not by 2.

Internal energy of an ideal gas

Ideal molecules are points: no intermolecular PE, no rotation or vibration in this model. So the **internal energy** 内能 is **all** kinetic:

$$U = \frac{3}{2}NkT = \frac{3}{2}nRT$$

$U \propto T$ – double the thermodynamic temperature and you double the internal energy.



Two explanations from the model

“Explain why the internal energy of an ideal gas is proportional to its thermodynamic temperature.” [2]

There are **no intermolecular forces**, so no potential energy: U is the total **kinetic** energy of the molecules.

Mean KE = $\frac{3}{2}kT$, so $U = \frac{3}{2}NkT \propto T$.

“A gas is heated at constant volume. Explain why its pressure rises.” [3]

The molecules move **faster**: each hits a wall with a **larger change of momentum**, and they hit the walls **more often**. So the force on the walls, and the pressure, increase.

Compressing at constant T is different: the molecules are not faster, they only hit the walls more often.

4

Deriving $pV = \frac{1}{3}Nm\langle c^2 \rangle$

Sheet 15.3.1 – what you need to know

You must be able to:

- derive $pV = \frac{1}{3}Nm\langle c^2 \rangle$ in five steps
- tie each **assumption** to the step it allows
- state the meaning of N , m and $\langle c^2 \rangle$
- show that $\frac{1}{2}m\langle c^2 \rangle = \frac{3}{2}kT$

What the exam asks


“State two of the basic assumptions of the kinetic theory of gases.” [2]

“State the meaning of the symbols N , m and $\langle c^2 \rangle$.” [3]

“Use the equation of state to show that $E_K = \frac{3}{2}kT$.” [2]

Practise on exercise sheet 15.3.1, then try the real exam questions on past-paper sheet 15.3.

Where the pressure comes from



One molecule, one wall, in a cube of side L :

- bounce: the **momentum** 动量 change is $\Delta p_x = -2mu_1$, so the wall takes an **impulse** 冲量 of $+2mu_1$
- next hit after $\Delta t = 2L/u_1$
- so $F_1 = \frac{\Delta p}{\Delta t} = \frac{mu_1^2}{L}$

Sum over N , then use $\langle u_x^2 \rangle = \frac{1}{3} \langle c^2 \rangle$:

$$pV = \frac{1}{3} Nm \langle c^2 \rangle$$

The $\frac{1}{3}$ is **symmetry**: no direction is special, so each axis carries a third of the motion.

Each assumption has a job in the derivation

assumption	the step it allows
collisions are elastic	the molecule rebounds at the same speed: $\Delta p = 2mu$
molecules' own volume is negligible	each one crosses the whole length L : $\Delta t = 2L/u$
no forces between collisions	straight lines at constant speed between walls
collision time negligible	the time between hits is all travel time
large number , random motion	average over N , and no direction is special

$$c^2 = u^2 + v^2 + w^2 \Rightarrow \langle u^2 \rangle = \frac{1}{3} \langle c^2 \rangle$$

Pythagoras in 3D, then average

“State the meaning of $\langle c^2 \rangle$ ”: the **mean square speed** – the mean of the squared speeds, not the square of the mean.

5

Recap

Exam tips

- Use $pV = nRT$ (n in mol) or $pV = NkT$ (N molecules), with temperature in **kelvin**.
- Learn the **kinetic-theory assumptions** (random motion, negligible molecular volume, elastic collisions, no intermolecular forces).
- Mean translational KE = $\frac{3}{2}kT$ – it depends **only on temperature**.

Topic 15 – key takeaways

1 The mole

$$N = nN_A, N_A = 6.02 \times 10^{23}$$

2 State

$$pV = nRT = NkT, \text{ always in kelvin}$$

3 Kinetic theory


$$pV = \frac{1}{3}Nm\langle c^2 \rangle \text{ from elastic bounces}$$

4 Energy

$$\langle E_k \rangle = \frac{3}{2}kT - \text{temperature only}$$

Check yourself

- 1 How many molecules in 2.0 mol?
- 2 Which form suits a count of molecules?
- 3 Why is there a $\frac{1}{3}$ in $pV = \frac{1}{3} Nm \langle c^2 \rangle$?
- 4 At the same T , which is faster: H_2 or O_2 ?



Answers 1. 1.2×10^{24} 2. $pV = NkT$ 3. symmetry – a third per axis 4. H_2