


Electric Fields

A2 Physics ■ Topic 18

A-Level Physics



What we will cover

- 1 Fields and field lines 18.1
- 2 Uniform fields 18.2
- 3 Coulomb's law 18.3
- 4 Field of a point charge 18.4
- 5 Electric potential 18.5
- 6 Recap



1

Fields and field lines

Sheet 18.1 – what you need to know

You must be able to:

- define an **electric field** and field strength
 $E = F/q$
- draw field lines for plates, point charges and pairs
- draw a charge's path between plates and after it leaves them

What the exam asks

“Define electric field.” [2]

“On Fig. 5.1, draw the path of the electron as it moves between and beyond the plates.” [2]

“Explain, with reference to the forces exerted by the two fields on the electron, why the path of the electron is undeviated.” [2]

Practise on exercise sheet 18.1, then try the real exam questions on past-paper sheet 18.1.

Electric field strength



An **electric field** 电场 is a region where a charge feels a force. Its **field strength** 电场强度 is the force per unit **positive** charge:

$$E = \frac{F}{q} \Rightarrow F = qE, \quad [E] = \text{N C}^{-1}$$



Defined on a **positive** test charge – so a **negative** charge feels a force **against** the field.

Field-line patterns


a) parallel plates (uniform field)



b) dipole (+ and -)



c) point charge (radial)



d) sphere above earthed plate

- they run from + to - (or off to infinity)
- they never cross
- closer lines = stronger field

Two crossing lines would give the force at that point **two** directions at once – impossible, which is why it never happens.

Four field patterns worth recognising

Point charge

Radial lines pointing *out* from a positive **point charge** 点电荷, and *in* towards a negative one.

A dipole

Two opposite charges – a **dipole** 偶极子 – give lines curving from + to –, densest in the gap between them.

Parallel plates


Two parallel charged plates a distance d apart with **potential difference** 电势差 V give a **uniform field** 匀强电场 of size $E = V/d$, apart from edge effects.

Several charges

For more than one charge, add the separate fields as a **vector sum** 矢量和 at each point.

A charge in a uniform field feels a constant force $F = qE$ and so a constant acceleration $a = qE/m$ – exactly like a mass in a uniform **gravitational field** 引力场.

Between the plates, and beyond them



Field lines between plates: **parallel**, **evenly spaced**, from + to -. The path is a parabola, **not** a circle: the force keeps **one direction**.

“Define electric field.” [2]

A region where a **charge** experiences a **force**.

“With a magnetic field added, explain why the electron is undeviated.” [2]

The electric force eE and the magnetic force Bev are **equal in size** and **opposite in direction**, so the resultant force is zero: $v = E/B$.

2

Uniform fields

Sheet 18.2 – what you need to know

You must be able to:

- use $E = V/d$ between parallel plates
- find the force and acceleration of a charge:
 $a = qE/m$
- explain the velocity selector, $u = V/Bd$

What the exam asks


“Use the symbols + and – to indicate which of plates X and Y is the positive plate.” [1]

“Determine the acceleration of the electron.” [2]

“Show that the speed u of ions that pass undeviated through the velocity selector is given by $u = V/Bd$.” [2]

Practise on exercise sheet 18.2, then try the real exam questions on past-paper sheet 18.2.

The uniform field between plates



Plates a distance d apart with p.d. V across them:

$$E = \frac{V}{d}$$

pointing from the **high**-potential plate to the low.

This is where $V \text{ m}^{-1}$ comes from – and it is the **same unit** as N C^{-1} .

Worked example – plates and an electron


Plates 5.0 mm apart carry a p.d. of 200 V. Find E , and the force on an electron between them. ($e = 1.6 \times 10^{-19} \text{ C}$)

$$E = \frac{V}{d} = \frac{200}{5.0 \times 10^{-3}} = 4.0 \times 10^4 \text{ V m}^{-1}$$

$$F = qE = (1.6 \times 10^{-19})(4.0 \times 10^4) = 6.4 \times 10^{-15} \text{ N}$$

Convert **mm to m** first – forgetting it scales E by a thousand. The force is the **same everywhere** in the gap: that is what uniform means.

A charge crossing a uniform field



Constant force $\Rightarrow$ constant acceleration

$$a = qE/m:$$

- released **at rest**: speeds up along the field, gaining KE
- entering **at right angles**: a **parabolic** 抛物线 path

Exactly a **projectile** 抛体 problem
 – steady velocity across, uniform acceleration down. This is how a **cathode-ray tube** 阴极射线管 steered its beam.

Two ways a charge is set moving

Released at rest

It accelerates along the field (positive charge) or against it (negative), gaining **kinetic energy** 动能 qV after falling through a potential difference V .

Fired across

Constant velocity along the plates, constant acceleration across them – the same parabola as a projectile.

In an atom

An **electron** 电子 orbiting a **nucleus** 原子核 of charge $+Ze$ at radius r : the Coulomb attraction supplies the **centripetal force** 向心力, $\frac{Ze^2}{4\pi\epsilon_0 r^2} = \frac{mv^2}{r}$.

Energy method or force method – both work. Use energy when the question gives a potential difference, force when it gives a field.

Worked example – acceleration in a uniform field

Two plates 2.0 cm apart have a p.d. of 500 V across them. Find the acceleration of an electron between them.

$$E = \frac{V}{d} = \frac{500 \text{ V}}{0.020 \text{ m}} = 2.5 \times 10^4 \text{ V m}^{-1}$$

$$\begin{aligned} F &= eE = 1.6 \times 10^{-19} \times 2.5 \times 10^4 \\ &= 4.0 \times 10^{-15} \text{ N} \end{aligned}$$

$$\begin{aligned} a &= \frac{F}{m_e} = \frac{4.0 \times 10^{-15} \text{ N}}{9.11 \times 10^{-31} \text{ kg}} \\ &= 4.4 \times 10^{15} \text{ m s}^{-2} \end{aligned}$$

towards the **positive** plate.

Halve d at the same V and E , F and a all **double**. Gravity on the electron ($\sim 10^{-29} \text{ N}$) is negligible.

The velocity selector

“Show that the speed u of ions that pass undeviated is $u = V/Bd$.” [2]

Undeviated: the forces balance,

$$qE = Bqu$$

$$u = \frac{E}{B} = \frac{V/d}{B} = \frac{V}{Bd}$$

q cancels, and m never appears: an alpha particle at the same speed also passes straight through.

$V = 150 \text{ V}$, $d = 5.0 \text{ mm}$, $B = 0.060 \text{ T}$.
Find u .

$$\begin{aligned} u &= \frac{150}{0.060 \times 5.0 \times 10^{-3}} \\ &= 5.0 \times 10^5 \text{ m s}^{-1} \end{aligned}$$

Faster ions?

Bqv grows with v but qE does not, so a faster ion is deflected **in the direction of the magnetic force**.

3

Coulomb's law

Sheet 18.3 – what you need to know

You must be able to:

- state **Coulomb's law** in words and use
$$F = Q_1 Q_2 / 4\pi\epsilon_0 r^2$$
- balance a Coulomb force against weight and tension
- use the Coulomb force as a centripetal force in an atom

What the exam asks

“State Coulomb's law.” [2]

“Determine the ratio electric field strength due to the other electron / electric field strength due to the nucleus.” [2]

“Determine the mass of sphere Y.” [4]

Practise on exercise sheet 18.3, then try the real exam questions on past-paper sheet 18.3.


Coulomb's law

like charges



repel — pushed apart

opposite charges



attract — pulled together

$$F = \frac{Q_1 Q_2}{4\pi\epsilon_0 r^2}$$

with the **permittivity of free space** 真空介电常数

$$\epsilon_0 = 8.85 \times 10^{-12} \text{ F m}^{-1} \text{ and } \frac{1}{4\pi\epsilon_0} \approx 9.0 \times 10^9 \text{ N m}^2 \text{ C}^{-2}.$$

Inverse square, along the joining line – **like repels, opposite attracts**. That sign choice is the one thing gravity cannot do.

Worked example – two point charges


Find the force between $+2.0\text{ nC}$ and $+3.0\text{ nC}$ set 4.0 cm apart.

$$F = \frac{(9.0 \times 10^9)(2.0 \times 10^{-9})(3.0 \times 10^{-9})}{(0.040)^2}$$

$\approx 3.4 \times 10^{-5}\text{ N}$, and **repulsive** (both +)

State the **direction** as well as the size – “repulsive” is part of the answer. And r goes in as **metres**, squared.

A charged sphere hanging in equilibrium



Y (+20 nC) hangs on a 1.2 m thread, pushed 0.080 m sideways by X (+20 nC), 0.060 m away. Find the mass of Y.

$$F = \frac{Q_1 Q_2}{4\pi\epsilon_0 r^2} = \frac{8.99 \times 10^9 \times (20 \times 10^{-9})^2}{0.060^2}$$

$$= 1.0 \times 10^{-3} \text{ N}$$

$$\tan \theta = \frac{F}{mg} \approx \frac{0.080}{1.2}$$

$$m = \frac{F}{g \tan \theta} = \frac{1.0 \times 10^{-3} \times 1.2}{9.81 \times 0.080}$$

$$= 1.5 \times 10^{-3} \text{ kg}$$

Coulomb forces in an atom

Hydrogen: the electron orbits at $r = 5.3 \times 10^{-11}$ m. Find its speed.

$$F = \frac{e^2}{4\pi\epsilon_0 r^2} = 8.2 \times 10^{-8} \text{ N}$$

$$F = \frac{mv^2}{r} \Rightarrow v = \sqrt{\frac{Fr}{m}}$$

$$v = \sqrt{\frac{8.2 \times 10^{-8} \times 5.3 \times 10^{-11}}{9.11 \times 10^{-31}}} \\ = 2.2 \times 10^6 \text{ m s}^{-1}$$

Helium: nucleus $+2e$; the two electrons are on opposite sides, each at r . Find $E_{\text{other electron}}/E_{\text{nucleus}}$.

$$\frac{e/(2r)^2}{2e/r^2} = \frac{1}{4} \times \frac{1}{2} = \frac{1}{8}$$

It points **outwards**, so it reduces the net inward force.

“State Coulomb’s law.” [2]

The force between two **point charges** is proportional to the **product of the charges** and inversely proportional to the **square of their separation**.

4

Field of a point charge

Sheet 18.4 – what you need to know

You must be able to:

- use $E = Q/4\pi\epsilon_0 r^2$, measuring r from the centre of a sphere
- sketch E against distance, with its sign
- link V , E , Q and C for a charged sphere

What the exam asks

“Explain why the electric potential near an isolated proton is positive.” [3]

“Sketch the variation of the electric field E at point P with x for values of x between $x = -3R$ and $x = 3R$.” [3]

“Determine the capacitance of the sphere.” [2]


Practise on exercise sheet 18.4, then try the real exam questions on past-paper sheet 18.4.

Spheres as point charges

A charged spherical **conductor** 导体 of total charge Q :

- **outside**: identical to a point charge Q at its centre (measure r from the **centre**)
- **inside** a hollow one: the field is **zero**

Zero field inside makes the whole conductor one **equipotential** 等势面 – and is why a car shields you from lightning.



$$E = F/q \quad (\text{NC}^{-1})$$

The field of a point charge


Coulomb's law gives the force between **two** charges. Divide by the test charge and only the **source** is left:

$$E = \frac{F}{q} = \frac{Q}{4\pi\epsilon_0 r^2}$$

Out of a positive Q , into a negative one, falling as $1/r^2$.

The same shape as $g = GM/r^2$ – but charges can **cancel**, so for several of them add the fields as a **vector** sum.

The field of a point charge, in detail




The field of a point charge Coulomb's law gives the force between two charges.

Signs: why $V > 0$ near a proton, and E either side of a sphere

“Explain why the electric potential near an isolated proton is positive.”
[3]

Potential is the work done **per unit positive charge** bringing it from **infinity**.

A positive test charge is **repelled** by the proton,
so work must be done **on** it: V is positive.



Left of the sphere the field points the **other way**, so E is **negative** there.

Worked example – charge and capacitance of a sphere

An isolated charged sphere has potential 45 kV and field strength $3.0 \times 10^5 \text{ V m}^{-1}$ at its surface. Find its radius, its charge and its capacitance.

$$V = \frac{Q}{4\pi\epsilon_0 r} \text{ and } E = \frac{Q}{4\pi\epsilon_0 r^2}, \text{ so } V = Er.$$

$$r = \frac{V}{E} = \frac{4.5 \times 10^4}{3.0 \times 10^5} = 0.15 \text{ m}$$

$$Q = 4\pi\epsilon_0 rV$$

$$= 4\pi \times 8.85 \times 10^{-12} \times 0.15 \times 4.5 \times 10^4$$

$$= 7.5 \times 10^{-7} \text{ C}$$

$$C = \frac{Q}{V} = 1.7 \times 10^{-11} \text{ F}$$

$C = 4\pi\epsilon_0 r$ for an isolated sphere: it depends only on the radius.

5

Electric potential

Sheet 18.5 – what you need to know

You must be able to:

- define **electric potential** and use $V = Q/4\pi\epsilon_0 r$
- add potentials from several charges, with signs
- sketch V along a line and link E to its gradient

What the exam asks

“Define electric potential at a point.” [2]

“Show that the electric potential V at point P when $x = 10 \text{ pm}$ is equal to 130 V .” [2]

“Sketch the variation of V with x between $x = 10 \text{ pm}$ and $x = 110 \text{ pm}$.” [3]

Practise on exercise sheet 18.5, then try the real exam questions on past-paper sheet 18.5.


Electric potential

Electric potential 电势 is the work done per unit **positive** charge bringing a test charge in from **infinity** 无穷远:

$$V = \frac{W}{q} = \frac{Q}{4\pi\epsilon_0 r}$$

Note $1/r$, **not** $1/r^2$. V is a **scalar** and takes the sign of Q – so several charges just **add**, signs and all, with no directions to resolve.

Electric potential, in detail



Worked example – potential near a charge

Find V at 4.0 cm from a point charge of +3.0 nC.

$$V = \frac{Q}{4\pi\epsilon_0 r} = \frac{(9.0 \times 10^9)(3.0 \times 10^{-9})}{0.040}$$
$$\approx 6.8 \times 10^2 \text{ V}$$


Same charge, same distance as the Coulomb example – but r appears to the **first** power here. Mixing up $1/r$ and $1/r^2$ is the standard slip.

Adding potentials: a proton and an electron

A proton is at $x = 0$ and an electron at $x = 120$ pm. Show that $V = 130$ V at $x = 10$ pm.

Potential is a **scalar**: add with signs.

$$\begin{aligned}
 V &= \frac{e}{4\pi\epsilon_0} \left(\frac{1}{r_p} - \frac{1}{r_e} \right) \\
 &= 1.44 \times 10^{-9} \left(\frac{1}{10^{-11}} - \frac{1}{1.1 \times 10^{-10}} \right) \\
 &= 130 \text{ V}
 \end{aligned}$$



- $V = 0$ at the **midpoint** (60 pm): equal and opposite terms
- positive near the proton, negative near the electron

Where $V = 0$ is not where $E = 0$

Charges $+5.0 \text{ nC}$ and -2.0 nC are 0.70 m apart. Where between them is $V = 0$?

$$\frac{5.0}{x} = \frac{2.0}{0.70 - x}$$

$$5.0(0.70 - x) = 2.0x$$


$$x = \frac{3.5}{7.0} = 0.50 \text{ m from the } +5.0 \text{ nC}$$

Is the field strength also zero there?

No. Between unlike charges both fields point **the same way** (towards the negative charge), so they add.

V adds as a **scalar**; E adds as a **vector**. $E = 0$ needs the fields to cancel, which happens **between like charges**.

Field is the potential gradient



$E = -dV/dr$ · field = negative gradient of potential

The field is the **negative potential gradient**:

$$E = -\frac{dV}{dx}$$

- between plates V falls evenly $\Rightarrow E = V/d$, constant
- point charge:

$$E = -\frac{dV}{dr} = \frac{Q}{4\pi\epsilon_0 r^2}$$

Minus sign: E points **downhill**, high V to low.

Electric potential energy


A charge q where the potential is V has $E_P = qV$; for two point charges

$$E_P = \frac{Qq}{4\pi\epsilon_0 r}$$

- **like**: $E_P > 0$ – energy released if they fly apart
- **opposite**: $E_P < 0$ – a **bound** 束缚 well

Both tend to **zero** as $r \rightarrow \infty$ – that is the choice of zero, exactly as for gravity.

Electric potential energy, in detail



A charge q sitting where the potential is V has $E_P = qV$, so for two point charges

$$E_P = \frac{Qq}{4\pi\epsilon_0 r}.$$

Like charges give $E_P > 0$, opposite charges $E_P < 0$; both tend to zero at infinity.

Gravitational vs electric

Quantity	Gravitational	Electric
Source	mass M (always +)	charge Q (either sign)
Field strength	$g = GM/r^2$	$E = Q/(4\pi\epsilon_0 r^2)$
Force	$F = mg$	$F = qE$
Gravitational potential 引力势 vs potential	$\phi = -GM/r$	$V = Q/(4\pi\epsilon_0 r)$
PE of two	$-GMm/r$	$Qq/(4\pi\epsilon_0 r)$
Nature	always attractive	attract <i>or</i> repel

Identical machinery, one difference: gravity's minus sign is **permanent** (mass only attracts), while the electric sign is inherited from the **charge**.

6

Recap

Exam tips

- **Coulomb's law** $F = Q_1 Q_2 / 4\pi\epsilon_0 r^2$ (inverse-square, but can attract or repel).
- Distinguish a **uniform field** ($E = V/d$, between plates) from a **radial field** ($E = Q/4\pi\epsilon_0 r^2$).
- Electric potential $V = Q/4\pi\epsilon_0 r$ (sign follows the charge); the field points from high to low potential.

Topic 18 – key takeaways

1 Coulomb

$F = Q_1 Q_2 / 4\pi\epsilon_0 r^2$ – attract or repel

2 Two fields

uniform $E = V/d$; radial $E = Q/4\pi\epsilon_0 r^2$

3 Potential


$V = Q/4\pi\epsilon_0 r$ – scalar, signed, $1/r$

4 The link

$E = -dV/dx$ – the field points downhill in V

Check yourself

- 1 Which way does a field line point?
- 2 Halve the plate gap at fixed V – what happens to E ?
- 3 Is potential a vector or a scalar?
- 4 What is the field inside a hollow charged sphere?



Answers 1. the force on a + charge 2. doubles 3. scalar 4. zero