

# Medical Physics

A2 Physics ■ Topic 24

A-Level Physics



## What we will cover

- 1    Ultrasound    24.1
- 2    Diagnostic information from echoes    24.1.1
- 3    X-rays    24.2
- 4    Controls, layers, sharpness, CT    24.2.1
- 5    PET scanning    24.3
- 6    Recap

1

Ultrasound

---

# Sheet 24.1 – what you need to know

## You must be able to:

- explain the piezo-electric effect both ways
- use  $Z = \rho c$  and  $\alpha = (Z_2 - Z_1)^2 / (Z_2 + Z_1)^2$
- use  $I = I_0 e^{-\mu x}$ , and solve it for  $\mu$

## What the exam asks

“Explain how ultrasound waves are detected by a piezoelectric crystal.” [2]

“Calculate the linear attenuation coefficient  $\mu$  ... Give a unit with your answer.” [2]

“Explain, without calculation, what is likely to happen when ultrasound is incident on a tissue–water boundary.” [2]

*Practise on exercise sheet 24.1, then try the real exam questions on past-paper sheet 24.1.*

## Piezo-electric – two linked effects

A **piezo-electric** 压电 crystal (quartz, PZT) changes shape when a p.d. is put across it, and makes an e.m.f. when its shape is changed.



### apply a p.d.

the crystal vibrates –  
used to **make** ultrasound

### squeeze it

an e.m.f. appears – used  
to **detect** ultrasound


## One transducer both sends and listens

An alternating p.d. (a few MHz) makes the crystal vibrate, sending **longitudinal** 纵波 waves above 20 kHz (1–10 MHz for imaging). Returning ultrasound makes it vibrate and produce an e.m.f.

One transducer switches between sending and listening. **Coupling gel** 耦合剂 pushes out the air – without it almost all the ultrasound would reflect at the skin.

## Inside the transducer

The crystal sits between a **damping block** (to kill ringing so pulses stay short) and a **matching layer** (to cut the reflection at the skin).



# Specific acoustic impedance

$$Z = \rho c$$

unit  $\text{kg m}^{-2} \text{s}^{-1}$

At a boundary the fraction **reflected** is

$$\frac{I_R}{I_0} = \left( \frac{Z_1 - Z_2}{Z_1 + Z_2} \right)^2$$

| <b>material</b> | $Z / 10^6 \text{ kg m}^{-2} \text{s}^{-1}$ |
|-----------------|--------------------------------------------|
| air             | 0.0004                                     |
| soft tissue     | 1.63                                       |
| bone            | 7.8                                        |

| <b>boundary</b> | <b>reflected</b> |
|-----------------|------------------|
| tissue / air    | 99.9%            |
| tissue / bone   | 43%              |

Gel removes the air gap, so the pulse gets **in**.

# Impedance, reflection and attenuation

|                          |                                                                                                                                                      |
|--------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------|
| acoustic impedance       | $Z = \rho c$ – <b>density</b> 密度 times the speed of sound. Soft tissue:<br>$1060 \times 1540 \approx 1.6 \times 10^6 \text{ kg m}^{-2}\text{s}^{-1}$ |
| reflection at a boundary | the fraction reflected depends on the <b>difference</b> in $Z$ between the two media                                                                 |
| best for imaging         | different <b>enough</b> to give an echo, but not so different that nothing passes on                                                                 |
| why gel is used          | air's $Z$ is so low that almost everything would reflect at the skin                                                                                 |
| attenuation              | $I = I_0 e^{-\mu x}$ ; higher-energy X-rays penetrate further, so their $\mu$ is <b>smaller</b>                                                      |
| contrast                 | bone has a much larger $\mu$ than soft tissue, which is why it shows white                                                                           |


To find a thickness, rearrange to  $x = \frac{1}{\mu} \ln \frac{I_0}{I}$ . Half-value thickness is the special case  $I = I_0/2$ .

## Attenuation – intensity falls exponentially


$$I = I_0 e^{-\mu x}$$

$\mu$  is the **attenuation coefficient** 衰减系数 (unit  $\text{m}^{-1}$ ). The same form applies to X-rays.

Each **half-value thickness**  $x_{1/2} = \ln 2 / \mu$  halves the intensity again.



# Attenuation – intensity falls exponentially, in detail



Intensity falls exponentially with thickness; a larger attenuation coefficient (bone) falls off faster.

# Impedance and attenuation: the answers that score

## “Define specific acoustic impedance.” [2]

The product of the **density** of a medium and the **speed of sound** in it:  $Z = \rho c$ . (So  $\rho = Z/c$  in a table.)

## “Explain, without calculation, what happens at a tissue–water boundary.” [2]

Their  $Z$  values are **similar**, so very little is reflected: almost all is **transmitted**.

Check:  $\alpha = \left( \frac{1.63 - 1.48}{1.63 + 1.48} \right)^2 = 0.002$ , only 0.2%.

## Ultrasound falls to 35% of its intensity through 4.0 cm of soft tissue. Find $\mu$ , with a unit.

$$\frac{I}{I_0} = e^{-\mu x} \Rightarrow \mu = \frac{1}{x} \ln \frac{I_0}{I}$$

$$\mu = \frac{1}{0.040 \text{ m}} \ln \frac{1}{0.35} = 26 \text{ m}^{-1}$$

2

Echoes

---

## Sheet 24.1.1 – what you need to know

### You must be able to:

- describe how a crystal sends and receives ultrasound
- find a depth from an echo time, and what fraction of the pulse returns
- choose a frequency; explain the gel, and why lungs cannot be imaged

### What the exam asks

“Describe how the piezoelectric crystal in a transducer generates ultrasound waves for use in medical diagnosis.” [3]

“Describe how  $\alpha$  depends on the relative values of  $Z_1$  and  $Z_2$ .” [2]

“Explain why ultrasound cannot be used to produce an image inside an air-filled cavity such as the lungs.” [2]

*Practise on exercise sheet 24.1.1, then try the real exam questions on past-paper sheet 24.1.*


## Pulse-echo: depth from the time delay

- 1 send a short pulse
- 2 at each tissue boundary, part reflects and part goes on
- 3 detect each echo
- 4 depth  $d = ct/2$  (there and back); amplitude  $\Rightarrow$  strength of the reflection

**Echo after  $60 \mu\text{s}$ ;  $c = 1500 \text{ m s}^{-1}$**

$$d = \frac{ct}{2} = \frac{1500 \times 60 \times 10^{-6}}{2} = 0.045 \text{ m (4.5 cm)}$$


## Pulse-echo: depth from the time delay, in detail



$$\frac{I_R}{I_0} = \left( \frac{Z_1 - Z_2}{Z_1 + Z_2} \right)^2 \text{ — a bigger impedance difference gives a bigger echo}$$


At a boundary, part of the pulse reflects and part transmits – a bigger impedance difference gives a bigger echo.

# Pulse-echo imaging



An A-scan shows the transmitted pulse and the echoes from each tissue boundary

# What the operator sees



A B-scan builds a 2-D image from many pulse-echo lines swept across the body.

# Making and receiving ultrasound

## “Describe how the crystal generates ultrasound.” [3]

An **alternating p.d.** is applied across the crystal;  
it **expands and contracts**, so it vibrates;  
at its **resonant** frequency the vibrations are largest, sending out ultrasound.

## “Explain how ultrasound is detected.” [2]

The returning wave makes the crystal **vibrate**, which produces an **alternating p.d.** across it.

- **pulses**: the same crystal listens between them, and the delay gives the depth
- **lungs**: air's  $Z$  is tiny, so nearly **all** is reflected at the first air boundary and nothing returns from inside
- higher  $f$ : shorter  $\lambda$ , **finer detail**, but more attenuation, so **less depth**
- non-ionising, so safe for a **fetus**

## Worked example – how much of the echo comes back

**A boundary lies 5.0 cm deep in tissue with  $\mu = 23 \text{ m}^{-1}$ ; it reflects a fraction  $\alpha = 0.010$ . What fraction of the sent intensity returns to the probe?**

The wave travels **there and back**:

$$2d = 0.10 \text{ m.}$$

$$\begin{aligned}\frac{I}{I_0} &= \alpha e^{-\mu(2d)} = 0.010 \times e^{-23 \times 0.10} \\ &= 0.010 \times 0.100 = 1.0 \times 10^{-3}\end{aligned}$$

Only a thousandth comes back, so the receiver must be very sensitive.

Using  $d$  instead of  $2d$  is the usual lost mark.

Gel removes the air between probe and skin. Without it, almost all the ultrasound reflects at the skin.

3

**X-rays**

---

## Sheet 24.2 – what you need to know

### You must be able to:

- describe the X-ray tube, with its polarity
- use  $\lambda_{\min} = hc/eV$  and photon momentum  $p = E/c$
- explain contrast; find the percentage attenuated

### What the exam asks

“Use + and – signs to label terminals X and Y to indicate the polarity of the high voltage.” [1]


“Explain why X-rays can be used to produce images of internal body structures that have good contrast.” [3]

“Calculate the percentage of the initial intensity of the wave that is attenuated by the tissue.” [2]

*Practise on exercise sheet 24.2, then try the real exam questions on past-paper sheet 24.2.*


## How an X-ray tube works

- 1 heated **cathode** 阴极 emits electrons by **thermionic emission** 热电子发射
- 2 a high p.d. accelerates them across a vacuum to a metal **target** 靶 anode (often tungsten)
- 3 they slow sharply: most KE becomes heat; a small part is X-ray photons (**Bremsstrahlung** 轫致辐射). Knocking out inner electrons gives **characteristic** 特征 lines.



# The X-ray tube

Most of the electrons' kinetic energy becomes **heat**; a small part leaves as X-ray photons, so the anode is angled and cooled.



## Minimum wavelength and imaging

The most energy one photon can have is  $eV$ :


$$hf_{\max} = eV, \quad \lambda_{\min} = \frac{hc}{eV}$$

Bone (higher density and proton number) attenuates more and looks lighter; soft tissue looks darker. A **contrast medium** 造影剂 (barium meal) makes soft tissues stand out. Half-value thickness  $x_{1/2} = \ln 2 / \mu$  halves the intensity.

$$x = 3.0 \text{ cm}, \quad \mu = 40 \text{ m}^{-1}$$

$$I/I_0 = e^{-40 \times 0.030} = e^{-1.2} \approx 0.30$$

# Minimum wavelength




X-ray: continuous bremsstrahlung + characteristic peaks

A typical X-ray spectrum – a continuous Bremsstrahlung curve cut off at  $\lambda_0$ , with sharp characteristic peaks

## Why bone shows white

Dense bone (high proton number) absorbs strongly and leaves the film unexposed; the air-filled lungs let X-rays through and darken it.



# X-ray contrast and percentage attenuated

**“Explain why X-rays can produce images with good contrast.” [3]**

Different tissues have different **attenuation coefficients**  $\mu$ ;  
 so very different fractions are transmitted;  
 so the intensity reaching the detector differs **sharply** between regions.

A 100 keV photon:  $E = 1.6 \times 10^{-14}$  J,  
 $p = E/c = 5.3 \times 10^{-23}$  N s,  
 $\lambda_{\min} = 1.2 \times 10^{-11}$  m.

**3.0 cm of tissue transmits 72%. What percentage is attenuated by 5.0 cm?**

$$\mu = \frac{1}{0.030} \ln \frac{1}{0.72} = 11.0 \text{ m}^{-1}$$

$$\frac{I}{I_0} = e^{-11.0 \times 0.050} = 0.58$$

attenuated =  $1 - 0.58 = 42\%$

The trap: the question asks for the part **removed**,  $1 - I/I_0$ , not the part transmitted.

## Ultrasound and X-rays side by side

|                 | ultrasound                          | X-rays                             |
|-----------------|-------------------------------------|------------------------------------|
| produced by     | a vibrating piezo-electric crystal  | electrons hitting a metal target   |
| image made from | waves reflected at boundaries       | waves transmitted through the body |
| ionising?       | no                                  | yes: keep the dose low             |
| best for        | soft tissue, a fetus, moving organs | bone; CT for 3-D slices            |

Ultrasound fails at air and bone boundaries (almost total reflection); X-rays show little difference between soft tissues unless a contrast medium is used.

# 4

## Controls and CT

---

## Sheet 24.2.1 – what you need to know

### You must be able to:

- say what the filament current and the p.d. each control
- attenuate through layers; use the half-value thickness
- tell sharpness from contrast; explain CT

### What the exam asks

“Calculate the percentage of the X-ray energy that is absorbed by a 2.8 cm thickness of this medium.” [3]

“Explain how X-rays are produced at the metal target.” [2]

“State the name of the charged particles that are accelerated by the applied p.d.” [1]

*Practise on exercise sheet 24.2.1, then try the real exam questions on past-paper sheet 24.2.*

# The tube's two controls, layers and sharpness

- **filament current**: more electrons, so more photons – the **intensity**
- **accelerating p.d.**: the largest photon energy – the **hardness**
- an aluminium **filter** removes soft, low-energy X-rays that only add to the skin dose
- **sharpness**: a small focal spot, a still patient, the detector close to the patient

**“Explain how X-rays are produced at the target.” [2]**

**Electrons** decelerate rapidly in the target; the kinetic energy they lose leaves as X-ray **photons**.

**Find the percentage of the X-ray energy absorbed by 2.8 cm of a medium with  $\mu = 0.20 \text{ cm}^{-1}$ .**

$$\frac{I}{I_0} = e^{-0.20 \times 2.8} = 0.57 \Rightarrow 43\% \text{ absorbed}$$


Through layers the exponents **add**:  $I = I_0 e^{-(\mu_1 x_1 + \mu_2 x_2)}$ .

# CT – many angles, then a computer

## Computed tomography 计算机断层扫描:

- 1 tube and detectors **rotate**, imaging one thin slice from many angles
- 2 a computer combines them into a 2-D cross-section
- 3 the patient moves on; the next slice is imaged
- 4 slices stack into a **3-D** image

Tissues that overlap on one X-ray are **separated**.



5

PET

---

## Sheet 24.3 – what you need to know

### You must be able to:

- say what a **tracer** is and write the  $\beta^+$  decay
- explain annihilation into two 511 keV photons
- locate the event from a coincidence and a time difference

### What the exam asks

“State what is meant by a tracer.” [2]

“State the name of the particles that are detected.” [1]

“Explain how these particles are formed inside the body.” [2]


*Practise on exercise sheet 24.3, then try the real exam questions on past-paper sheet 24.3.*

## A $\beta^+$ tracer, then annihilation

A **tracer** 示踪剂 (often FDG, fluorine-18 on a glucose analogue) is taken up more by active tissue. In PET it is a  $\beta^+$  emitter – it gives out a **positron** 正电子.

The positron meets an electron and they **annihilate** 湮灭: two photons, opposite directions, each 511 keV ( $= m_e c^2$ ).

$$2hf = 2m_e c^2$$




two photons, opposite directions — the coincidence fixes the line

## A coincidence fixes the line

The two photons fly off **back to back** and hit the detector ring. Two **simultaneous** arrivals (a coincidence) fix the line the annihilation lay on.

Many lines from many angles build a 3-D map of the tracer: tissues with high **metabolic activity**.

Arrival-time differences refine the position along the line (time-of-flight PET).




# Tracers and annihilation

|                     |                                                                                                                   |
|---------------------|-------------------------------------------------------------------------------------------------------------------|
| a tracer 示踪剂        | a substance containing radioactive nuclei, introduced into the body                                               |
| how it works        | it is taken up more by the tissue being studied, so that tissue emits more strongly                               |
| the choice          | a $\gamma$ or $\beta^+$ emitter with a short half-life, so the dose falls quickly                                 |
| an antiparticle 反粒子 | the same mass, opposite charge – the positron is the electron's                                                   |
| annihilation 湮灭     | a particle meets its antiparticle and their mass turns into electromagnetic energy                                |
| the signature       | two $\gamma$ photons of 0.51 MeV leave in opposite directions, and detecting both fixes the line the event lay on |

PET works because momentum conservation forces the two photons back to back. That geometry is what locates the tracer.

# PET: the answers that score, and locating the event



two photons detected together: the event lies on the line AB  
the time difference gives where on the line:  $d = c \Delta t / 2$  from the midpoint

**Photon B arrives 0.40 ns after A. How far from the midpoint was the annihilation?**

$$d = \frac{c \Delta t}{2} = \frac{3.00 \times 10^8 \times 0.40 \times 10^{-9}}{2} = 6.0 \text{ cm}$$


**“State what is meant by a tracer.” [2]**

A substance containing a **radioactive nuclide** that is put into the body and **absorbed** by the tissue being studied.

**“Name the particles detected, and explain how they form.” [3]**

**Gamma-ray photons.** A positron from  $\beta^+$  decay meets an electron and they **annihilate**: two 511 keV photons go off in **opposite** directions (momentum is conserved).

## A finished PET image



**Positron emission tomography** 正电子发射断层成像 (PET) maps function, not shape: warm colours mark where the tracer collected – the tissue with the fastest **metabolism** 代谢.

6

Recap

---

## Exam tips

- **Ultrasound** reflects at boundaries (acoustic impedance); a coupling gel reduces reflection at the skin.
- **X-rays** are attenuated as  $I = I_0 e^{-\mu x}$ ; contrast media (barium, iodine) absorb strongly.
- **PET** uses a positron emitter; annihilation gives two  $\gamma$ -rays in opposite directions, located by the detector ring.

## Topic 24 – key takeaways

### 1 Ultrasound

piezo;  $d = ct/2$ ; gel;  $Z = \rho c$

### 2 X-rays

$$\lambda_{\min} = hc/eV; I = I_0 e^{-\mu x}$$

### 3 CT

many angles of one slice, then stack

### 4 PET

two 511 keV photons; coincidence line

## Check yourself

- 1 Why  $d = ct/2$ , not  $ct$ ?
- 2 Why put gel between the transducer and the skin?
- 3 What sets  $\lambda_{\min}$  of an X-ray spectrum?
- 4 Why do PET photons fly in opposite directions?

**Answers** 1. there and back 2. air/skin reflects almost everything 3.  $hf_{\max} = eV$   
4. momentum conservation (pair at rest)