


Astronomy and Cosmology

A2 Physics ■ Topic 25

A-Level Physics



What we will cover

- 1 Luminosity, flux and standard candles 25.1
- 2 Stellar temperatures and radii 25.2
- 3 Redshift, Hubble, the Big Bang 25.3
- 4 Recap

1

Standard candles

Sheet 25.1 – what you need to know

You must be able to:

- define **luminosity** and use $F = L/4\pi d^2$
- explain how a standard candle gives a distance
- use the Doppler shift, including from a rotating star

What the exam asks

“Explain how standard candles are used to determine the distance to a galaxy.” [3]

“... Show that the radius of the Sun is 6.93×10^8 m.” [3]

“State and explain how the wavelength observed from Z compares with the emitted wavelength.” [2]

Practise on exercise sheet 25.1, then try the real exam questions on past-paper sheet 25.1.

Total power, then power per unit area


Luminosity 光度 L : the **total power** a star radiates (W). At distance d it has spread over a sphere $4\pi d^2$:

$$F = \frac{L}{4\pi d^2}$$

radiant flux intensity

辐射通量密度, W m^{-2}

Measure F , know $L \Rightarrow d = \sqrt{\frac{L}{4\pi F}}$.



inverse-square law

平方反比定律: double d , and F falls to a **quarter**.


Worked example – flux at the Earth

Sun: $L = 3.8 \times 10^{26} \text{ W}$, $d = 1.5 \times 10^{11} \text{ m}$

$$F = \frac{L}{4\pi d^2} = \frac{3.8 \times 10^{26}}{4\pi(1.5 \times 10^{11})^2}$$

$$\approx 1.4 \times 10^3 \text{ W m}^{-2}$$

Worked example – flux at the Earth, in detail



The same power spreads over a larger area as distance grows, so flux falls as $1/d^2$

An object whose luminosity is known from its type

Find one in a distant galaxy, measure F , get $d = \sqrt{L/(4\pi F)}$. Reaches far beyond **parallax** 视差.

Cepheids


造父变星

pulsation period is tightly
linked to luminosity
– the period gives L

Type Ia supernovae

a white dwarf at critical
mass always has about
the same peak luminosity

Standard candles




Cepheid: longer period $\Rightarrow$ brighter · standard candles

For Cepheid variables the pulsation period sets the luminosity, making them standard candles

Each rung calibrates the next

Parallax reaches the nearest stars; Cepheids reach nearby galaxies; supernovae reach the far ones. Every rung is calibrated by the one below it.



each rung is calibrated by the one below, reaching ever further

Standard candles: the answers that score

“Explain how standard candles are used to determine the distance to a galaxy.” [3]

A standard candle is an object of **known luminosity** L (a Cepheid, a type Ia supernova);
measure its **radiant flux intensity** F at the Earth;


$$\text{then } d = \sqrt{\frac{L}{4\pi F}}.$$

“Suggest two reasons why stars differ in apparent brightness.” [2]

Different **luminosities**; different **distances**.

Flux **lower** than predicted (dust absorbs some light)? Then d comes out **too large**.

Measuring the Sun's radius from its rotation



At one limb the 656.30 nm line is shifted by 0.00460 nm. The period is 2.07×10^6 s. Show that $r = 6.93 \times 10^8$ m.

$$\begin{aligned}
 v &= c \frac{\Delta\lambda}{\lambda} = 3.00 \times 10^8 \times \frac{0.00460}{656.30} \\
 &= 2.10 \times 10^3 \text{ m s}^{-1} \\
 r &= \frac{vT}{2\pi} = \frac{2.10 \times 10^3 \times 2.07 \times 10^6}{2\pi} \\
 &= 6.93 \times 10^8 \text{ m}
 \end{aligned}$$

The centre of the disc moves **across** the line of sight, so its light is **not** shifted.

2

Stellar radii

Sheet 25.2 – what you need to know

You must be able to:

- use Wien's law $\lambda_{\text{max}} T = \text{constant}$
- use $L = 4\pi r^2 \sigma T^4$, including by ratios
- find a star's radius or distance from L , F and T

What the exam asks

“Explain why the observed wavelength and the emitted wavelength have different values.”
[2]

“Explain how the temperature determined using the observed wavelength compares with the true value.” [2]

“Determine the distance of the star from the Earth.” [2]

Practise on exercise sheet 25.2, then try the real exam questions on past-paper sheet 25.2.

Wien – hotter means bluer

A blackbody spectrum peaks at λ_{max} set by temperature:


$$\lambda_{\text{max}} T = b \approx 2.90 \times 10^{-3} \text{ m K}$$

Cool red star $\sim 3000 \text{ K}$ peaks in the infrared; the Sun $\sim 5800 \text{ K}$ near 500 nm ; a hot blue-white star $\sim 20\,000 \text{ K}$ in the ultraviolet.

$$\lambda_{\text{max}} = 500 \text{ nm}$$

$$T = b/\lambda_{\text{max}} = 2.90 \times 10^{-3} / (500 \times 10^{-9}) \approx 5800 \text{ K}$$

Wien – hotter means bluer, in detail



Wien: $\lambda_{\text{max}} T = \text{const} \cdot \text{hotter} \Rightarrow \text{peak shifts left}$

A hotter black body radiates more, and its peak wavelength shifts towards the blue (Wien's law)

Stefan–Boltzmann – $L \propto r^2 T^4$

A star as a blackbody sphere: The **Stefan–Boltzmann law** 斯特藩-玻尔兹曼定律, with the **Stefan–Boltzmann constant** 斯特藩-玻尔兹曼常数 σ :

$$L = 4\pi\sigma r^2 T^4, \quad \sigma = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$$

Twice the radius $\Rightarrow$ four times L . Twice the temperature $\Rightarrow$ sixteen times L . To estimate a radius: $\lambda_{\text{max}} \rightarrow T$ (Wien); F and $d \rightarrow L$; then

$$r = \sqrt{\frac{L}{4\pi\sigma T^4}}$$

Estimating a star's radius

Wien, then Stefan–Boltzmann

- 1 $\lambda_{\max} \Rightarrow T$
($\lambda_{\max} T = 2.9 \times 10^{-3} \text{ m K}$)
- 2 $L = 4\pi d^2 F$ from flux and distance
- 3 $L = 4\pi r^2 \sigma T^4 \Rightarrow r = \sqrt{\frac{L}{4\pi\sigma T^4}}$



Two laws and a distance are enough to size a star nobody can resolve.

Luminosity, and the law behind it

Luminosity

The **luminosity** 光度 L of a star is its **total power** 总功率 of radiation – the **energy** 能量 radiated per second in all directions.

Stefan–Boltzmann law

The **Stefan–Boltzmann law**: $L = 4\pi r^2 \sigma T^4$, where $\sigma = 5.67 \times 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$ is the **Stefan–Boltzmann constant**.

Why it gives a radius

Wien's law gives T from the peak wavelength, the flux and distance give L – so the only unknown left is r .

$L \propto T^4$: doubling the surface temperature multiplies the power by sixteen, which is why hot stars are so much brighter than their size suggests.

From flux to a star's radius, and the distance ladder

Stefan–Boltzmann law 斯特藩-玻尔兹曼定律	$L = 4\pi r^2 \sigma T^4$; σ is the Stefan–Boltzmann constant 斯特藩-玻尔兹曼常量
the route	Wien's law gives T ; flux and distance give L ; rearrange for r
why it matters	radii of stars that cannot be resolved as a disc
the ladder	each method is calibrated by the one below it
the rungs	radar, parallax, standard candles, then Hubble's law
Big Bang evidence	galactic redshift, the cosmic microwave background 宇宙微波背景, the H/He abundance

Each rung reaches further and is less precise. An error low on the ladder carries all the way up.

Ratio reasoning with $L = 4\pi r^2 \sigma T^4$

Star B has the same luminosity as the Sun but twice its surface temperature. Compare its radius.

$$r^2 \propto \frac{L}{T^4}, \text{ so } r \propto \frac{1}{T^2}: \quad r_B = \frac{1}{4} r_{\odot}.$$

Hot and small: a **white dwarf**.


Same radius, half the temperature?

$$L \propto T^4: \quad L \text{ is } \left(\frac{1}{2}\right)^4 = \frac{1}{16} \text{ as large.}$$

A star of luminosity $3.8 \times 10^{28} \text{ W}$ gives a flux of $2.0 \times 10^{-9} \text{ W m}^{-2}$ at the Earth. How far away is it?

$$\begin{aligned} d &= \sqrt{\frac{L}{4\pi F}} = \sqrt{\frac{3.8 \times 10^{28}}{4\pi \times 2.0 \times 10^{-9}}} \\ &= 1.2 \times 10^{18} \text{ m} \end{aligned}$$

A redshifted spectrum gives too low a temperature



“Explain why the observed and emitted wavelengths differ.” [2]

The star (galaxy) is **moving away**, so every wavelength is **redshifted**:

$$\lambda_{\text{obs}} > \lambda_{\text{emit}}.$$

“How does the temperature from the observed wavelength compare with the true value?” [2]

$T = b/\lambda_{\text{max}}$ and the observed λ_{max} is **longer**, so the temperature found is **too low**.

The Pillars of Creation



The Pillars of Creation in the Eagle Nebula – clouds of gas and dust lit by hot, newly formed stars

3

Hubble's law

Sheet 25.3 – what you need to know

You must be able to:

- use $\Delta\lambda/\lambda \approx \Delta f/f \approx v/c$
- use Hubble's law $v = H_0 d$ and estimate the age of the Universe
- explain how redshift supports the Big Bang

What the exam asks


“Explain how redshift leads to the idea that the Universe is expanding.” [3]

“Sketch the variation of v with d .” [2]

“State the name of the quantity represented by the gradient of the line.” [1]

Practise on exercise sheet 25.3, then try the real exam questions on past-paper sheet 25.3.

Spectral lines shifted to the red




Lines from distant galaxies sit at **longer** wavelengths: **redshift** 红移 – the galaxy is **receding**. For $v \ll c$:

$$\frac{\Delta\lambda}{\lambda} \approx \frac{v}{c}$$

Almost every distant galaxy is redshifted (a few near ones are **blueshifted** 蓝移): the Universe is **expanding**.

The same lines, moved

A galaxy's lines keep their **pattern** but sit at longer wavelengths – which is how the shift is recognised as a shift, and not as different elements.




Hubble's law

$$v \approx H_0 \cdot d$$

where H_0 is the **Hubble constant** 哈勃常数, $\approx 2.3 \times 10^{-18} \text{ s}^{-1}$. Always SI units. Running the expansion backwards, distances shrink to zero at $t = -1/H_0$ – the **Big Bang** 大爆炸. For steady expansion the age is


$$T_{\text{age}} \approx \frac{1}{H_0} \approx 14 \text{ billion years}$$

Hubble's law, in detail



Hubble's law: a galaxy's recession speed is proportional to its distance, $v = H_0 d$

How far the ladder reaches



Andromeda – Cepheids put it well outside our own galaxy.



Almost every smudge here is a galaxy, most of them receding.

Each rung calibrates the next: parallax, then **Cepheid** 造父变星 variables, then redshift.

Worked example – a receding galaxy

Emitted $4.62 \times 10^{-7} \text{ m}$, **seen at** $4.91 \times 10^{-7} \text{ m}$. $H_0 = 2.3 \times 10^{-18} \text{ s}^{-1}$.

$\Delta\lambda = 0.29 \times 10^{-7} \text{ m}$, so $v \approx (\Delta\lambda/\lambda)c \approx 1.9 \times 10^7 \text{ m s}^{-1}$.


Distance $d = v/H_0 \approx 8.3 \times 10^{24} \text{ m}$.

Redshift and the expanding Universe

“Explain how redshift leads to the idea that the Universe is expanding.” [3]

Light from distant galaxies is **redshifted**, so they are **moving away**; the **further** the galaxy, the **greater** its redshift (speed); so all of space is expanding, as from a single point.

An approaching source is **blueshifted**: $\Delta\lambda$ is negative.



Sketch: a straight line **through the origin**. The gradient is the **Hubble constant**.

Worked example – two recipes for one galaxy

A Cepheid puts a galaxy at $d = 1.0 \times 10^{24}$ m. A line emitted at 656.3 nm is observed at 661.1 nm. Find H_0 and estimate the age of the Universe.

$$v = c \frac{\Delta\lambda}{\lambda} = 3.00 \times 10^8 \times \frac{4.8}{656.3}$$

$$= 2.2 \times 10^6 \text{ m s}^{-1}$$

$$H_0 = \frac{v}{d} = \frac{2.2 \times 10^6}{1.0 \times 10^{24}} = 2.2 \times 10^{-18} \text{ s}^{-1}$$

$$t \approx \frac{1}{H_0} = 4.5 \times 10^{17} \text{ s}$$

$$\approx 1.4 \times 10^{10} \text{ years}$$

assuming the galaxies have always moved apart at the **same speed**.

Distance from a **standard candle**; speed from the **redshift**. The two together give H_0 .

4

Recap

Exam tips

- $F = L/4\pi d^2$ (inverse-square): a standard candle has known luminosity L , so the distance follows from the measured flux F .
- **Wien's law** ($\lambda_{\max} \propto 1/T$) gives surface temperature; **Stefan's law** $L = 4\pi r^2 \sigma T^4$.
- **Redshift** $z = \Delta\lambda/\lambda \approx v/c$; Hubble's law $v = H_0 d$ leads to the age of the universe (evidence for the Big Bang).

Topic 25 – key takeaways

1 Flux

$$F = L / (4\pi d^2); \text{ inverse-square}$$

2 Candles

Cepheids, Type Ia; d from known L

3 Star

$$\text{Wien } \lambda_{\text{max}} T = b; L = 4\pi\sigma r^2 T^4$$

4 Hubble

$$v = H_0 d; \text{ age } \approx 1/H_0$$

Check yourself

- 1 Double the distance: what happens to flux?
- 2 What makes a Cepheid a standard candle?
- 3 A hotter star: longer or shorter λ_{max} ?
- 4 Hubble's law points to what origin of the Universe?

Answers 1. falls to a quarter 2. period gives luminosity 3. shorter 4. the Big Bang