

4 (a) State the value of absolute zero on:

(i) the Celsius temperature scale

temperature = ..... °C [1]

(ii) the thermodynamic temperature scale. Give a unit with your answer.

temperature = ..... unit ..... [1]

(b) A sample contains a fixed amount of gas. The gas has pressure  $p$ , volume  $V$  and thermodynamic temperature  $T$ .

Fig. 4.1 shows the variation of  $pV$  with  $kT$  for the sample, where  $k$  is the Boltzmann constant.




Fig. 4.1

(i) State what is indicated about the nature of the gas from the variation shown in Fig. 4.1.

..... [1]

(ii) Determine the number  $N$  of molecules of the gas in the sample.

$N =$  ..... [2]

(iii) Use your answer in (b)(ii) to determine the amount  $n$  of gas in the sample.

$n =$  ..... mol [1]

(c) The root-mean-square (r.m.s.) speed of the molecules of the gas is  $1900 \text{ m s}^{-1}$  when  $pV$  is equal to  $270 \text{ J}$ .

Determine the mass, in  $u$ , of one molecule of the gas, where  $u$  is the unified atomic mass unit.

mass = .....  $u$  [4]

[Total: 10]

2 A student investigates the cooling of a liquid in a beaker.

The temperature  $\theta_R$  of the laboratory is measured using a thermometer.

Hot water is added to an insulated beaker, as shown in Fig. 2.1.




Fig. 2.1

The thermometer measures the temperature of the water. At time  $t$  the temperature of the water is  $\theta$ .

A series of readings of  $t$  and  $\theta$  are taken.

It is suggested that  $\theta$  and  $t$  are related by the equation

$$\theta = \theta_R + (\theta_0 - \theta_R)e^{-Kt}$$

where  $\theta_0$  is the temperature at  $t = 0$  and  $K$  is a constant.

(a) A graph is plotted of  $\ln(\theta - \theta_R)$  on the  $y$ -axis against  $t$  on the  $x$ -axis.

Determine expressions for the gradient and  $y$ -intercept.

gradient = .....

$y$ -intercept = .....

(b) Values of  $t$  and  $\theta$  are given in Table 2.1.

Table 2.1

| $t/\text{min}$ | $\theta/^\circ\text{C}$ | $(\theta - \theta_R)/^\circ\text{C}$ | $\ln((\theta - \theta_R)/^\circ\text{C})$ |
|----------------|-------------------------|--------------------------------------|-------------------------------------------|
| 6.0            | $75.0 \pm 0.5$          |                                      |                                           |
| 12.0           | $64.5 \pm 0.5$          |                                      |                                           |
| 18.0           | $57.0 \pm 0.5$          |                                      |                                           |
| 24.0           | $50.0 \pm 0.5$          |                                      |                                           |
| 30.0           | $44.5 \pm 0.5$          |                                      |                                           |
| 36.0           | $41.0 \pm 0.5$          |                                      |                                           |


The value of  $\theta_R$  is  $(18.5 \pm 0.5)^\circ\text{C}$ .

Calculate and record values of  $(\theta - \theta_R)/^\circ\text{C}$  and  $\ln((\theta - \theta_R)/^\circ\text{C})$  in Table 2.1. Include the absolute uncertainties in  $(\theta - \theta_R)$  and  $\ln((\theta - \theta_R)/^\circ\text{C})$ .

[2]

- (c) (i) Plot a graph of  $\ln((\theta - \theta_R)/^\circ\text{C})$  against  $t/\text{min}$ . Include error bars for  $\ln((\theta - \theta_R)/^\circ\text{C})$ . [2]
- (ii) Draw the straight line of best fit and a worst acceptable straight line on your graph. Label both lines. [2]
- (iii) Determine the gradient of the line of best fit. Include the absolute uncertainty in your answer.

gradient = ..... [2]



- (iv) Determine the y-intercept of the line of best fit. Include the absolute uncertainty in your answer.

y-intercept = ..... [2]

- (d) (i) Using your answers to (a), (c)(iii) and (c)(iv), determine the values of  $K$  and  $\theta_0$ . Include appropriate units.

$K = \dots\dots\dots$

$\theta_0 = \dots\dots\dots$  [2]

- (ii) Determine the absolute uncertainty in your value of  $\theta_0$ .

absolute uncertainty = ..... [1]

- (e) Determine the time  $t$  for the temperature to reach  $25.0^\circ\text{C}$ .

$t = \dots\dots\dots \text{min}$  [1]

[Total: 15]