

28 Science: data representation and defensible extrapolation

Name: _____ Class: _____ Date: _____ Total: 13 marks

KEY WORDS rate of change • interpolation • extrapolation • evidence
• rate • proposal

Objective

- Read values with correct axes, units and conditions
- Translate a table into a trend and interpolate within a measured interval
- Evaluate an extrapolation against the model's assumptions

Original public practice; marks below are classroom self-check points, not official criteria.

1 Key knowledge and worked examples


Data Representation passages use tables, graphs or diagrams as primary evidence . Start with the variable labels and units, and identify which conditions are held constant. A row may report one trial or an average; a legend may distinguish groups measured under different conditions. Locate the requested value before using prior scientific knowledge.

Translate representations without changing their meaning. A rising quantity can have a falling rate of increase. A steep line is meaningful only relative to axis scales and units. When comparing curves, distinguish an absolute value, a difference and a rate. Use corresponding x-values for a fair comparison unless the stem explicitly asks for another relationship.

Interpolation estimates inside observed bounds. For a straight segment between two points, use the fraction of the input interval and apply it to the output difference. Extrapolation extends beyond those bounds and depends more strongly on the proposed model. Negative predicted quantities or physical limits supplied by the passage can reveal a model's invalid extension.

Reevaluation asks whether new evidence supports an earlier trend or requires a different explanation. One discordant measurement could reflect noise or a model limitation; use the stated uncertainty and experimental conditions. The dataset here is original and hypothetical, so the task's given model supplies the relevant relationships. It is not a claim about every real material.

A rate of change compares output change with the matching input interval. $r = \Delta T / \Delta t = (32 - 28)^\circ\text{C} / (4 - 2) \text{ min} = 2^\circ\text{C/min}$. A rising temperature can have a falling rate. Read units before calculating.



Original worked example from existing native teaching; transfer tasks use their own data.

Worked example from the native lesson: Original trial: identical containers hold a test liquid, starting at 20°C in the same room. Recorded times in minutes are 0, 2, 4, 6; measured temperatures in °C are 20, 28, 34, 38. The liquid warms, but consecutive two-minute rises are 8, 6 and 4°C: its average warming rate decreases. Linear interpolation between 2 and 4 minutes gives $T(3) = 28 + (3 - 2) / (4 - 2) \times (34 - 28) = 31^\circ\text{C}$. Extending the first interval's 4°C/min rate to 6 minutes predicts 44°C, inconsistent with the measured 38°C. New evidence therefore rejects a constant-rate model across all six minutes.

■ Complete original case context

Original hypothetical research summary – A cooling sleeve

A laboratory team tests sleeves intended to reduce a small sensor's temperature during operation. All sensors start at 22°C. Temperature is measured after ten minutes of use with the same power setting, room conditions and measuring instrument. Sleeve A costs 4 units, B costs 7 units, and an unsleeved sensor is the control. The design brief requires a cost no greater than 5 units and a final temperature no greater than 35°C. Lower temperature alone is not the only design criterion.

Experiment 1 uses identical sensors and records three independent trials per sleeve. Final temperatures in °C are: control 41,42,43; A 33,34,35; B 30,31,32. The team resets the starting temperature and checks the sensor charge before each trial. They report the means and the full observed ranges. The ranges describe these trials, not every future result. The team has not yet measured sleeve durability.

Experiment 2 investigates thickness for material A. Each sleeve has the same length and fit; thicknesses are 1,2,3 mm. Mean final temperatures are 38,34,32°C respectively. All other stated conditions match Experiment 1. Changing thickness also changes material

quantity, so a future design review must measure costs instead of assuming they stay at 4 units. A student proposes testing a 3-mm sleeve in material B against a 1-mm sleeve in material A to identify thickness’s effect. This comparison would change two factors at once.

Two original models explain the observed temperature pattern. Model P says final temperature depends only on thickness, regardless of material. Model Q says both thickness and material matter; at equal thickness it predicts that material B produces a lower final temperature than A under the same conditions. Both models predict lower final temperatures as thickness rises over the tested 1–3 mm interval. A decreasing trend alone therefore cannot separate them.

Experiment 3 tests new A and B sleeves, each 2 mm thick, in matched conditions. Mean final temperatures are A 34°C and B 31°C. The measuring instrument’s stated resolution is 0.1°C, and the team checks its calibration with a reference. Repeated trials would still be needed to characterise variability and rule out other differences in manufacture. The result conflicts with P’s material-independence prediction and is consistent with Q’s direction; it is not proof that Q is the only possible explanation.

A separate time series for one sensor with sleeve A records temperatures 22,28,32,34°C at 0,2,4,6 minutes. These observations show warming with successively smaller two-minute increases. No measurements beyond six minutes are supplied in this series. A proposal to continue the first interval’s slope to twelve minutes is a model assumption that later measurements may contradict, not a reading from the table.

2 Practice

2.1 Original trial: identical containers hold a test liquid, starting at 20°C in the same room. Recorded times in minutes are 0, 2, 4, 6; measured temperatures in °C are 20, 28, 34, 38. The liquid warms, but consecutive two-minute rises are 8, 6 and 4°C: its average warming rate decreases. Linear interpolation between 2 and 4 minutes gives $T(3)=28+(3-2)/(4-2)\times(34-28)=31^{\circ}\text{C}$. Extending the first interval’s 4°C/min rate to 6 minutes predicts 44°C, inconsistent with the measured 38°C. New evidence therefore rejects a constant-rate model across all six minutes.

What is the average rise from minute 4 to 6?

- A 4°C per minute
- B 6°C per minute
- C 38°C per minute
- D 2°C per minute

2.2 Original trial: identical containers hold a test liquid, starting at 20°C in the same room. Recorded times in minutes are 0, 2, 4, 6; measured temperatures in °C are 20, 28, 34, 38. The liquid warms, but consecutive two-minute rises are 8, 6 and 4°C: its average warming rate decreases. Linear interpolation between 2 and 4 minutes gives $T(3)=28+(3-2)/(4-2)\times(34-28)=31^{\circ}\text{C}$. Extending the first interval’s 4°C/min rate to 6 minutes predicts 44°C, inconsistent with the measured 38°C. New evidence therefore rejects a constant-rate model across all six minutes.

Which statement fits all four measurements?

- A Temperature rises while average warming rate falls
- B Temperature falls throughout
- C Temperature rises by equal increments
- D The liquid reaches 44°C at minute 6

2.3 Linear interpolation between 34°C at minute 4 and 38°C at minute 6 gives _____°C at minute 5.

2.4 An estimate outside the measured range is interpolation.

3 Exam-style questions

Use the complete original case context in section 1.

3.1 For the separate time series, calculate the three average two-minute warming rates. Describe temperature and rate separately.

3.2 Estimate temperature at minute 3 by linear interpolation. Extend the first interval's slope to minute 6 and compare with the actual measurement. [3]

3.3 Is a minute-12 value read from these data or extrapolated? Explain why a graph with a rising curve does not establish indefinite constant-rate warming. [3]
