

Past-paper walk-through

DNA and RNA Structure ■ 6.1

eXercise

Questions in this set

- 2022 Q2

Reading the mark scheme

- **B1** a mark that stands on its own – the point is either made or not.
- **M1** a *method* mark: the right approach, even if the number is wrong.
- **A1** an *answer* mark, and it usually needs its M mark first.
- **C1** a working mark on the way to the answer.
- **//** separates answers the examiner accepts equally.
- **ecf** = error carried forward: a later part is marked on your own earlier answer.
- **owtte** = “or words to that effect” – the wording need not match.

Question 2

2022 ■ Q2

During meiosis, double-strand breaks occur in chromatids. The breaks are either repaired by the exchange of genetic material between homologous nonsister chromatids, which is the process known as crossing over (Figure 1A), or they are simply repaired without any crossing over (Figure 1B). Plant breeders developing new varieties of corn are interested in determining whether, in corn, a correlation exists between the number of meiotic double-strand chromatid breaks and the number of crossovers.

[Figure 1: two diagrams of paired homologous chromosomes (each shown as two X-shaped sister-chromatid pairs, one gray and one white, joined at a chiasma near the bottom). Panel A: shows a double-strand break being repaired WITH crossing over — the chromatid pair undergoes an intermediate exchange step (crossed strands shown) and resolves into two chromosomes each with a mix of gray and white chromatid segments. Panel B: shows a double-strand break repaired WITHOUT crossing over —

Question 2

the chromatid pair resolves directly back into the original all-gray and all-white chromosomes with no exchange of segments.]

Figure 1. Double-strand breaks in chromatids are repaired with crossing over (A) or without crossing over (B).

Using specialized staining and microscopy techniques, scientists counted the number of double-strand chromatid breaks and the number of crossovers in the same number of meiotic gamete-forming cells of six inbred strains of corn (Table 1).



TABLE 1. NUMBER OF CHROMATID DOUBLE-STRAND BREAKS AND AVERAGE NUMBER OF CROSSOVERS IN INBRED STRAINS OF CORN

Strain of Corn	Number of Double-Strand Breaks	Average Number of Crossovers
($\pm 2SE_{\bar{x}}$) I 710 19.5 $\pm$ 0.5	II 650 18.0 $\pm$ 0.7	III 600 17.5 $\pm$ 1.0
IV 510 16.0 $\pm$ 1.0	V 425 14.0 $\pm$ 0.5	VI 325 11.0 $\pm$ 1.5

Question 2

(a) The double-strand breaks occur along the DNA backbone. Describe the process by which the breaks occur.

Question 2



Question 2

2022 ■ Q2

During meiosis, double-strand breaks occur in chromatids. The breaks are either repaired by the exchange of genetic material between homologous nonsister chromatids, which is the process known as crossing over (Figure 1A), or they are simply repaired without any crossing over (Figure 1B). Plant breeders developing new varieties of corn are interested in determining whether, in corn, a correlation exists between the number of meiotic double-strand chromatid breaks and the number of crossovers.

[Figure 1: two diagrams of paired homologous chromosomes (each shown as two X-shaped sister-chromatid pairs, one gray and one white, joined at a chiasma near the bottom). Panel A: shows a double-strand break being repaired WITH crossing over — the chromatid pair undergoes an intermediate exchange step (crossed strands shown) and resolves into two chromosomes each with a mix of gray and white chromatid segments. Panel B: shows a double-strand break repaired WITHOUT crossing over — the chromatid pair resolves directly back into the original all-gray and all-white chromosomes with no exchange of segments.]

Question 2

Figure 1. Double-strand breaks in chromatids are repaired with crossing over (A) or without crossing over (B).

Using specialized staining and microscopy techniques, scientists counted the number of double-strand chromatid breaks and the number of crossovers in the same number of meiotic gamete-forming cells of six inbred strains of corn (Table 1).

TABLE 1. NUMBER OF CHROMATID DOUBLE-STRAND BREAKS AND AVERAGE NUMBER OF CROSSTERS IN INBRED STRAINS OF CORN

Strain of Corn	Number of Double-Strand Breaks	Average Number of Crossovers ($\pm 2SE_{\bar{x}}$)
I	710	19.5 ± 0.5
II	650	18.0 ± 0.7
III	600	17.5 ± 1.0
IV	510	16.0 ± 1.0
V	425	14.0 ± 0.5
VI	325	11.0 ± 1.5

(b) Using the template in the space provided for your response, construct an appropriately labeled graph that represents the data in Table 1 and allows examination of a possible correlation between double-strand breaks and crossovers. Based on the

Question 2

data, determine whether corn strains I, II, and III differ in their average number of crossovers.

Question 2

2022 ■ Q2

During meiosis, double-strand breaks occur in chromatids. The breaks are either repaired by the exchange of genetic material between homologous nonsister chromatids, which is the process known as crossing over (Figure 1A), or they are simply repaired without any crossing over (Figure 1B). Plant breeders developing new varieties of corn are interested in determining whether, in corn, a correlation exists between the number of meiotic double-strand chromatid breaks and the number of crossovers.

[Figure 1: two diagrams of paired homologous chromosomes (each shown as two X-shaped sister-chromatid pairs, one gray and one white, joined at a chiasma near the bottom). Panel A: shows a double-strand break being repaired WITH crossing over — the chromatid pair undergoes an intermediate exchange step (crossed strands shown) and resolves into two chromosomes each with a mix of gray and white chromatid segments. Panel B: shows a double-strand break repaired WITHOUT crossing over — the chromatid pair resolves directly back into the original all-gray and all-white chromosomes with no exchange of segments.]

Question 2

Figure 1. Double-strand breaks in chromatids are repaired with crossing over (A) or without crossing over (B).

Using specialized staining and microscopy techniques, scientists counted the number of double-strand chromatid breaks and the number of crossovers in the same number of meiotic gamete-forming cells of six inbred strains of corn (Table 1).

TABLE 1. NUMBER OF CHROMATID DOUBLE-STRAND BREAKS AND AVERAGE NUMBER OF CROSSOVERS IN INBRED STRAINS OF CORN

Strain of Corn	Number of Double-Strand Breaks	Average Number of Crossovers ($\pm 2SE_{\bar{x}}$)
I	710	19.5 ± 0.5
II	650	18.0 ± 0.7
III	600	17.5 ± 1.0
IV	510	16.0 ± 1.0
V	425	14.0 ± 0.5
VI	325	11.0 ± 1.5

(c) Based on the data, describe the relationship between the average number of double-strand breaks and the average number of crossovers in the strains of corn analyzed in the experiment.

Question 2

2022 ■ Q2

During meiosis, double-strand breaks occur in chromatids. The breaks are either repaired by the exchange of genetic material between homologous nonsister chromatids, which is the process known as crossing over (Figure 1A), or they are simply repaired without any crossing over (Figure 1B). Plant breeders developing new varieties of corn are interested in determining whether, in corn, a correlation exists between the number of meiotic double-strand chromatid breaks and the number of crossovers.

[Figure 1: two diagrams of paired homologous chromosomes (each shown as two X-shaped sister-chromatid pairs, one gray and one white, joined at a chiasma near the bottom). Panel A: shows a double-strand break being repaired WITH crossing over — the chromatid pair undergoes an intermediate exchange step (crossed strands shown) and resolves into two chromosomes each with a mix of gray and white chromatid segments. Panel B: shows a double-strand break repaired WITHOUT crossing over — the chromatid pair resolves directly back into the original all-gray and all-white chromosomes with no exchange of segments.]

Question 2

Figure 1. Double-strand breaks in chromatids are repaired with crossing over (A) or without crossing over (B).

Using specialized staining and microscopy techniques, scientists counted the number of double-strand chromatid breaks and the number of crossovers in the same number of meiotic gamete-forming cells of six inbred strains of corn (Table 1).

TABLE 1. NUMBER OF CHROMATID DOUBLE-STRAND BREAKS AND AVERAGE NUMBER OF CROSSOVERS IN INBRED STRAINS OF CORN

Strain of Corn	Number of Double-Strand Breaks	Average Number of Crossovers ($\pm 2SE_{\bar{x}}$)
I	710	19.5 ± 0.5
II	650	18.0 ± 0.7
III	600	17.5 ± 1.0
IV	510	16.0 ± 1.0
V	425	14.0 ± 0.5
VI	325	11.0 ± 1.5

(d) Crossing over (Figure 1A) creates physical connections that are required for proper separation of homologous chromosomes during meiosis. A diploid cell with four pairs of homologous chromosomes undergoes meiosis to produce four haploid cells. Crossing over occurs between only three of the pairs. Predict the number of chromosomes most

Question 2

likely present in each of the four haploid cells. Provide reasoning to justify your prediction. Explain how plant breeders can use the information in Table 1 to help develop new varieties of corn.

Solution 2

(a)

Mark scheme

- The double-strand breaks occur through (enzymatic) hydrolysis of the covalent bonds between the sugars and phosphates (nucleotides) of the DNA backbone.

Solution 2

(b) Mark scheme

- 1 pt** Appropriate axis scaling on the graph (x-axis: number of double-strand breaks, roughly 0-800; y-axis: average number of crossovers, roughly 0-20+, both with regular, appropriate intervals). [1 pt] Accurately plotted X,Y points for each of the six corn strains (I-VI): (710,19.5) I, (650,18.0) II, (600,17.5) III, (510,16.0) IV, (425,14.0) V, (325,11.0) VI. [1 pt] Accurate error bars ($\pm 2SE$) plotted on each point per Table 1, e.g. Strain I: 19.5 ± 0.5 ; Strain VI: 11.0 ± 1.5 . [1 pt] Determination: there is no statistical difference in the average number of crossovers between Strains II and III (their error bars overlap); Strain I is higher/different from Strains II and III.

Solution 2

(c)

Mark scheme

- There is, in general, a direct correlation / positive relationship between the average number of double-strand breaks and the average number of crossovers across the corn strains analyzed - as double-strand breaks increase, crossovers increase.

Solution 2

(d) Mark scheme

- 1 pt Prediction: two of the four haploid cells will have three chromosomes ($n-1$) and two will have five chromosomes ($n+1$). [1 pt] Justification: during Meiosis I, three of the four homologous pairs (the ones with crossing over) separate normally, but the one pair without crossing over does not separate properly (experiences nondisjunction); in Meiosis II the sister chromatids separate normally, producing the $n-1/n+1$ combination in the four resulting cells. [1 pt] Application for plant breeders: because crossing over increases genetic diversity, breeders can breed/select corn strains with high double-strand-break numbers (which correlate with high crossover numbers), or increase the number of double-strand breaks, to increase genetic variation for developing new varieties.