

AP Biology

Scoring Guidelines

2018 SCORING GUIDELINES

Question 1




Figure 1. Phylogenetic tree representing the evolutionary relatedness among bear populations based on mitochondrial DNA sequence comparisons

Polar bears are highly adapted for life in cold climates around the North Pole. Brown bears, black bears, and pandas are found in warmer environments. Researchers collected complete mitochondrial DNA sequences from several populations of bears and constructed a phylogenetic tree to represent their evolutionary relatedness (Figure 1).

A researcher studying adaptation in bears sequenced the nuclear gene encoding a lysosomal trafficking protein (LYST) in polar bears, brown bears, black bears, and panda bears. There are seven inferred amino acid substitutions that are found only in polar bears. Mutations that cause similar substitutions in the human LYST protein are associated with Chediak-Higashi syndrome, an autosomal recessive condition in which pigment is absent from the hair and eyes. The researcher used the inferred amino acid sequences to build the distance matrix shown in Table 1.

TABLE 1. AMINO ACID DIFFERENCES IN THE LYST PROTEIN AMONG BEAR SPECIES

	Panda	Black	Brown	Polar
Panda	–			
Black	33	–		
Brown	34	1	–	
Polar	40	7	8	–

2018 SCORING GUIDELINES

Question 1 (continued)

- (a) Use the phylogenetic tree in Figure 1 to **estimate** the age in hundreds of thousands of years of the most recent common ancestor of all brown bears. **Identify** the population of brown bears to which polar bears are most closely related based on the mitochondrial DNA sequence comparison. **Identify** two populations whose positions could be switched without affecting the relationships illustrated in the phylogenetic tree.

Estimate (1 point)

- First two digits of the answer must be between 30 and 35.


Identification (1 point)

- European

Identification (1 point)

- European/Polar OR Asian/Western (North American)

- (b) **Construct** a cladogram on the template to represent a model of the evolutionary relatedness among the bear species based on the differences in LYST protein sequences (Table 1). **Circle** the position on the cladogram that represents the out-group.

**Construction (1 point)**

- Correctly illustrated evolutionary relationship among the four species

Circling (1 point)

- Correctly circled out-group based on orientation of cladogram

- (c) A student claims that mitochondrial DNA sequence comparisons provide a more accurate phylogeny of bear species than do LYST protein sequence comparisons. **Provide ONE piece of reasoning** to support the student's claim.

Reasoning (1 point)

- Genes show more variability (in nucleotide sequence) than proteins do (in amino acid sequences).
- mtDNA genome contains multiple genes vs. one *lyst* gene.
- The phenotype associated with the *lyst* gene is under strong selection.

2018 SCORING GUIDELINES

Question 1 (continued)

- (d) A researcher genetically engineers a mouse strain by deleting the mouse *lyst* gene and replacing it with the polar bear *lyst* gene. **Predict** the most likely difference in phenotype of the transgenic mouse strain compared to the wild-type mouse strain. **Justify** your prediction.

Prediction (1 point)

- Mouse fur and/or eyes will not have pigment/will have reduced pigment.
- Mouse (fur) will be white/lighter.

Justification (1 point)

- Polar bear *lyst* gene/LYST protein is associated with a lack of pigment/white hair.
- Mutated human *lyst* gene/ LYST protein is associated with a lack of pigment in hair and eyes.

- (e) **Describe** how the mutation in the *lyst* gene became common in the polar bear population. If the *lyst* gene were the only determinant of fur color, **predict** the percent of white offspring produced by a mating between a polar bear and a brown bear.

Description (1 point)

- Natural selection for the white fur phenotype

Prediction (1 point)

- 0%

2018 SCORING GUIDELINES

Question 2




Figure 1. Cellular response to infection by pathogenic bacteria

Some pathogenic bacteria enter cells, replicate, and spread to other cells, causing illness in the host organism. Host cells respond to these infections in a number of ways, one of which involves activating particular enzymatic pathways (Figure 1). Cells normally produce a steady supply of inactive caspase-1 protein. In response to intracellular pathogens, the inactive caspase-1 is cleaved and forms an active caspase-1 (step 1). Active caspase-1 can cleave two other proteins. When caspase-1 cleaves an inactive interleukin (step 2), the active portion of the interleukin is released from the cell. An interleukin is a signaling molecule that can activate the immune response. When caspase-1 cleaves gasdermin (step 3), the N-terminal portions of several gasdermin proteins associate in the cell membrane to form large, nonspecific pores.

Researchers created the model in Figure 1 using data from cell fractionation studies. In the experiments, various parts of the cell were separated into fractions by mechanical and chemical methods. Specific proteins known to be located in different parts of the cell were used as markers to determine the location of other proteins. The table below shows the presence of known proteins in specific cellular fractions.

CELL FRACTIONS CONTAINING DIFFERENT CELLULAR PROTEINS

	Aconitase (Krebs cycle protein)	DNA polymerase	GAPDH (glycolytic protein)	Sodium- potassium pump	NF- κ B (Immune response protein)
Whole cell sample	+	+	+	+	+
Fraction 1	+				
Fraction 2		+			+
Fraction 3			+		+
Fraction 4				+	

+ = presence of protein

2018 SCORING GUIDELINES

Question 2 (continued)

- (a) **Describe** the effect of inhibiting step 3 on the formation of pores AND on the release of interleukin from the cell.

Description (2 points)

- Pores will not form.
- Interleukin release will not be affected/interleukin release continues.

- (b) **Make a claim** about how cleaving inactive caspase-1 results in activation of caspase-1. A student claims that preinfection production of inactive precursors shortens the response time of a cell to a bacterial infection. **Provide ONE reason** to support the student's claim.

Claim (1 point)

- Removes inhibitor/repressor/inhibitory domain of protein
- Changes the shape/protein structure

Reasoning (1 point)

- Cleaving a precursor/protein/molecule is faster than making one upon infection.
- Cells do not have to wait for transcription and translation/protein synthesis.

- (c) A student claims that the NF- κ B protein is located in the cytoplasm until the protein is needed for transcription. **Justify** the student's claim with evidence. **Identify TWO** fractions where N-terminal gasdermin would be found in cells infected with pathogenic bacteria.

Justification (1 point)

- NF- κ B and glycolytic enzymes/GAPDH are found together (in the cytoplasm).

Identification (2 points)

- Fraction 3
- Fraction 4

- (d) **Describe** the most likely effect of gasdermin pore formation on water balance in the cell in a hypotonic environment.

Description (1 point)

- Water enters the cell.

- (e) **Explain** how gasdermin pore formation AND interleukin release contribute to an organism's defense against a bacterial pathogen.

Explanation (2 points)

- Cell lysis destroys infected cells OR cell lysis prevents bacteria from replicating.
- Interleukin signaling will stimulate immune cells/components of the immune system (to destroy the infected cells or bacteria).

2018 SCORING GUIDELINES



Question 3

Seagrasses are aquatic plants that reproduce sexually. Male seagrass flowers produce sticky pollen that is carried by circulating water to female flowers, resulting in fertilization. A researcher claims that mobile aquatic invertebrates can also transfer pollen from male to female flowers in the absence of circulating water. To investigate this claim, the researcher set up aquariums to model the possible interactions between the invertebrates and seagrasses.

- (a) Use the symbols below and the template aquariums to demonstrate the experimental design for testing the researcher's claim that mobile aquatic invertebrates can pollinate seagrass in the absence of circulating water. **Draw** the appropriate symbols in the negative control aquarium AND the experimental aquarium. Do not use any symbol more than once in the same aquarium.

Male Flower	Female Flower	Invertebrates
		

Drawing (2 points)

Negative Control Aquarium (1 point)	Experimental Aquarium (1 point)
	

- (b) **Identify** the dependent variable in the experiment. **Predict** the experimental results that would support the researcher's claim that mobile aquatic invertebrates can also transfer pollen from male to female flowers in the absence of circulating water.

Identification (1 point maximum)	Prediction (1 point maximum)
Number/presence of pollen grains on female flowers OR pollination	More pollen grains transferred/pollination seen in experimental aquarium
Number/presence of fertilized plants/flowers OR fertilization	More fertilized plants/flowers/fertilization seen in experimental aquarium
Number/presence of seed/fruit/offspring produced OR reproduction	More seeds/fruits/offspring produced/reproduction in experimental aquarium

2018 SCORING GUIDELINES

Question 4




Figure 1. Percent survival of five strains of bedbugs treated with beta-cyfluthrin. A (+) indicates the gene is present; a (-) indicates the gene is deleted. Error bars represent the 95% confidence interval.

The common bedbug (*Cimex lectularius*) is a species of insect that is becoming increasingly resistant to insecticides. Bedbugs possess several genes suspected of contributing to the resistance, including *P450*, *Abc8*, and *Cps*. To investigate the role of these genes in insecticide resistance, researchers deleted one or more of these genes in different strains of bedbugs, as indicated in Figure 1, and treated the strains with the insecticide beta-cyfluthrin. Each strain was genetically identical except for the deleted gene(s) and was equally fit in the absence of beta-cyfluthrin. The percent survival of each strain following beta-cyfluthrin treatment is shown in Figure 1.

- (a) **Identify** the control strain in the experiment. Use the means and confidence intervals in Figure 1 to **justify** the claim that *Abc8* is effective at providing resistance to beta-cyfluthrin.

Identification (1 point)

- Strain I

Justification (1 point)

- Error bars/CIs from strain I/control/WT do not overlap with strain III/*Abc8* deleted strain.
- Mean % survival of strain III/*Abc8* deletion falls outside the 95% confidence interval of strain I/control/WT.
- Strain III/*Abc8* deletion shows a statistically significant difference from strain I/control.

2018 SCORING GUIDELINES

Question 4 (continued)

- (b) *P450* encodes an enzyme that detoxifies insecticides. *Abc8* encodes a transporter protein that pumps insecticides out of cells. *Cps* encodes an external structural protein located in the exoskeleton that greatly reduces the absorption of insecticides. Based on this information and the data in Figure 1, **explain** how a deletion of both *P450* and *Abc8* results in lower survival in bedbugs compared with a deletion of *Cps* only.

Explanation (1 point per row; 2 points maximum)

Strain	<i>P450</i> and <i>Abc8</i>	<i>Cps</i> only	Explanation
V	Deleted	Present	Bedbugs can neither detoxify nor pump out insecticide, which results in a lower chance of bedbug survival.
IV	Present	Deleted	Bedbugs can detoxify and pump out insecticide, which results in a higher chance of bedbug survival.

2018 SCORING GUIDELINES

Question 5




Figure 1. Probability of nest success in an environment with predators

Some birds, including great spotted cuckoos, lay their eggs in the nests of other birds, such as reed warblers. The warbler parents raise the unrelated chicks and provide them with food that would otherwise be given to their biological offspring. A researcher conducted an investigation to determine the type of relationship between warblers and cuckoos in an environment without predators. The researcher found that nests containing only warblers were more likely to be successful than nests containing warblers and cuckoos (data not shown). A successful nest is defined as a nest where at least one chick becomes an adult warbler.

In some geographic areas, several species of nest predators are present. Researchers have found that cuckoo chicks, while in the nest, produce a smelly substance that deters nest predators. The substance does not remain in the nest if cuckoo chicks are removed. Figure 1 shows the probability that nests containing only warblers or containing both warblers and cuckoos will be successful in an environment with predators. In a follow-up experiment, the researchers added cuckoos to a nest that contained only warblers (group 1) and removed cuckoos from a nest containing warblers and cuckoos (group 2).

- (a) **Describe** the symbiotic relationship that exists between the cuckoo and warbler in an environment without predators.

Description (1 point)


- Cuckoos are parasites (of the warbler).
- The cuckoo benefits from the relationship, and the warbler is harmed by the relationship.

2018 SCORING GUIDELINES

Question 5 (continued)

(b) On the template provided, **draw** bars in the appropriate locations to predict the relative probability of success for the nest in the presence of predators where:

- the cuckoos were added to the nest containing only warblers (group 1)
- the cuckoos were removed from the nest containing warblers and cuckoos (group 2)



Graph (2 points)

- Cuckoo added (group 1): Bar must be HIGHER than the “Warblers Only” bar.
- Cuckoo removed (group 2): Bar must be LOWER than the “Warblers and Cuckoos” bar.

(c) **Identify** the symbiotic relationship that exists between the cuckoo and the warbler in the presence of predators.

Identification (1 point)

- Mutualism
- Both organisms benefit


2018 SCORING GUIDELINES

Question 6

Cystic fibrosis is a genetic condition that is associated with defects in the CFTR protein. The CFTR protein is a gated ion channel that requires ATP binding in order to allow chloride ions (Cl^-) to diffuse across the membrane.

(a) In the provided model of a cell, **draw** arrows to describe the pathway for production of a normal CFTR protein from gene expression to final cellular location.

Drawing (1 point)



The response must follow this pathway: nucleus/nuclear envelope → endoplasmic reticulum → Golgi apparatus → plasma membrane.

The response may be in the form of a continuous arrow or multiple discontinuous arrows.

(b) **Identify** the most likely cellular location of the ribosomes that synthesize CFTR protein.

Identification (1 point)

- (Rough) Endoplasmic Reticulum/ER

(c) **Identify** the most likely cellular location of a mutant CFTR protein that has an amino acid substitution in the ATP-binding site.

Identification (1 point)

- In the (cellular/plasma) membrane

2018 SCORING GUIDELINES

Question 7

In the tongue sole fish (*Cynoglossus semilaevis*), sex is determined by a combination of genetics and environmental temperature. Genetically male fish have two Z chromosomes (ZZ), and genetically female fish have one Z chromosome and one W chromosome (ZW). When fish are raised at 22°C, ZZ fish develop into phenotypic males and ZW fish develop into phenotypic females. However, when fish are raised at 28°C, the Z chromosome is modified (denoted as Z*). Z*W individuals develop as phenotypic males that are fertile and can pass on the Z* chromosome to their offspring even when the offspring are raised at 22°C. A cross between a ZW female and a Z*Z male is shown in the Punnett square below.

	Z	W
Z*	Z* Z	Z* W
Z	ZZ	ZW

- (a) **Predict** the percent of phenotypic males among the F₁ offspring of the cross shown in the Punnett square if the offspring are raised at 22°C.

Prediction (1 point)

75%

- (b) At least one Z or Z* chromosome is necessary for survival of the fish. A researcher crossed two fish and observed a 2:1 ratio of males to females among the offspring. Based on the information, **identify** the genotype of the male parent in the cross. **Describe** ONE fitness cost to the female of mating with this particular male.

Identification (1 point)

Z* W

Description (1 point)

- Fewer offspring will develop/survive.
- 1/4 of the offspring are predicted to die.
- Some of her offspring will have the Z* chromosome/all of her male offspring will have a Z* chromosome.

2018 SCORING GUIDELINES

Question 8

Acetylcholine receptor (AChR) proteins are found at the synapse between neurons and skeletal muscle cells. Acetylcholine released from neurons binds to a specific site on the receptor proteins, which causes an ion channel in the receptors to open and allow sodium ions (Na⁺) to enter muscle cells. The resulting depolarization of muscle cells initiates muscle contractions. Another molecule, nicotine, can also bind to certain types of AChR proteins and activate the receptors.

A researcher is investigating two different types of AChR proteins: type 1 and type 2. To determine which stimuli activate the receptors, the researcher exposes muscle cells expressing the different types of receptor proteins to stimuli and observes the results indicated in Table 1.

TABLE 1. RESPONSE OF AChR PROTEINS TO DIFFERENT STIMULI

AChR Protein Type	Acetylcholine	Nicotine
Type 1	+	+
Type 2	+	–

+ indicates activation
– indicates no activation

- (a) **Describe** the difference in the structure AND function between AChR type 1 and AChR type 2.

Description (2 points)

Points may be earned from only one row.

Structure (1 point maximum)	Function (1 point maximum)
Binding sites differ in shape/specificity/number	- Differential binding of molecules to type 1 and type 2 receptors - Activated by one (ACh) molecule or both (ACh and nicotine) molecules - No difference in response (both open channels OR both result in depolarization OR both cause muscle contraction)
Differential binding of molecules to type 1 and type 2 receptors	- Activated by one (ACh) or both (ACh and nicotine) molecules - No difference in response (both open channels OR both result in depolarization OR both cause muscle contraction)
Receptors activated by one (ACh) or both (ACh and nicotine) molecules	No difference in response (both open channels OR both result in depolarization OR both cause muscle contraction)

- (b) Acetylcholinesterase is an enzyme that breaks down acetylcholine in the synapse. **Describe** the effect of inhibiting acetylcholinesterase on the muscle cells with AChR type 2.

2018 SCORING GUIDELINES

Question 8 (continued)

Description (1 point)

- Continued activation
- Repeated opening of sodium channels OR repeated depolarization OR muscle spasms

2017

AP[®]

CollegeBoard

AP Biology

Scoring Guidelines

2017 SCORING GUIDELINES

Question

TABLE 1. EFFECT OF 0.1 mM CAFFEINE ON MEMORY IN BEES

Treatment	Memory (average probability of revisiting a nectar source $\pm 2SE_{\bar{x}}$)	
	10 Minutes	24 Hours
Control	0.72 ± 0.09	0.41 ± 0.07
Caffeine	0.83 ± 0.07	0.78 ± 0.08

In flowering plants pollination is a process that leads to the fertilization of an egg and the production of seeds. Some flowers attract pollinators, such as bees, using visual and chemical cues. When a bee visits a flower, in addition to transferring pollen, the bee can take nectar from the flower and use it to make honey for the colony.

Nectar contains sugar, but certain plants also produce caffeine in the nectar. Caffeine is a bitter-tasting compound that can be toxic to insects at high concentrations. To investigate the role of caffeine in nectar, a group of researchers studied the effect of 0.1 mM caffeine on bee behavior. The results of an experiment to test the effect of caffeine on bees' memory of a nectar source are shown in Table 1.

On the axes provided, **construct** an appropriately labeled graph to illustrate the effect of caffeine on probability of bees revisiting a nectar source (memory). **(3 points)**

Construct graph (3 points)

- Correctly plotted means on a bar graph/modified bar graph
- Appropriate labels, units, and scaling
- Correctly plotted error bars

Based on the results, **describe** the effect of caffeine on each of the following: **(2 points)**

- Short-term (10 minute) memory of a nectar source
- Long-term (24 hour) memory of a nectar source

Description (2 points)

Short-term	Caffeine does not affect short-term memory/memory at 10 minutes.
Long-term	Caffeine improves/increases the long-term memory/memory at 24 hours.

Design an experiment using artificial flowers to investigate potential negative effects of increasing caffeine concentrations in nectar on the number of floral visits by bees. **Identify** the null hypothesis, an appropriate control treatment, and the predicted results that could be used to reject the null hypothesis. **(3 points)**

2017 SCORING GUIDELINES

Question continued)

Identification (3 points; 1 point per row)

Null hypothesis	Increasing caffeine concentration has no effect (on the number of floral visits by bees).
Control	(Nectar/flowers with) no caffeine
Predicted results	- The number of floral visits by bees is different at increasing caffeine concentrations. - The number of floral visits by bees is different than the control.

searchers found that nectar with caffeine tends to have a lower sugar content than nectar without caffeine. Plants use less energy to produce the caffeine in nectar than they do to produce the sugar in nectar. **Propose ONE benefit** to plants that produce nectar with caffeine and a lower sugar content. **Propose ONE cost** to bees that visit the flowers of plants that produce nectar with caffeine and a lower sugar content. (2 points)

Proposed plant benefit (1 point)

- More pollen is transferred/more visits by pollinators.
- Plants store energy/have more energy available for other uses.

Proposed bee cost (1 point)

- (Individual) bees visit more flowers.
- (Individual) bees use more energy.
- The colony/bees may produce less honey
- The colony/bees may produce lower quality honey/honey that provides less energy.

2017 SCORING GUIDELINES

Question 2




Figure 1. The effect of karrikins (KAR) and trimethylbutenolides (TMB) on seed germination in *Lactuca* plants. Error bars represent $\pm 2SE_{\bar{x}}$.

Fires frequently occur in some ecosystems and can destroy all above-ground vegetation. Many species of plants in these ecosystems respond to compounds in smoke that regulate seed germination after a major fire. Karrikins (KAR) and trimethylbutenolides (TMB) are water-soluble compounds found in smoke that are deposited in the soil as a result of a fire. KAR and TMB bind to receptor proteins in a seed. In a study on the effects of smoke on seeds, researchers recorded the timing and percent of seed germination in the presence of various combinations of KAR and TMB. The results are shown in Figure 1.

In a second investigation into the effect of available water on seed germination after a fire, researchers treated seeds with KAR or TMB. The treated seeds were then divided into two treatment groups. One group received a water rinse and the other group received no water rinse. The seeds were then incubated along with a group of control seeds that were not treated. The results are shown in the table.

EFFECT OF CHEMICAL TREATMENT AND WATER RINSE ON GERMINATION

Treatment Group	Chemical Treatment		Water Rinse	Germination Result
	KAR	TMB		
1 (control)	–	–	–	Control result
2	+	–	–	Different from control
3	–	+	–	Different from control
4 (control)	–	–	+	Control result
5	+	–	+	Different from control
6	–	+	+	Same as control

2017 SCORING GUIDELINES

Question 2 (continued)

(a) The researchers made the following claims about the effect of KAR and the effect of TMB on seed germination relative to the control treatment.

- KAR alone affects the timing of seed germination.
- KAR alone affects the percentage of seeds that germinate.
- TMB alone affects the timing of seed germination.
- TMB alone affects the percentage of seeds that germinate.

Provide support using data from Figure 1 for each of the researchers' claims. **(4 points)**

Claim	Support (1 point each row; 4 points maximum)
KAR affects timing	• germination starts earlier/sooner/faster/quicker
KAR affects percentage	• higher percentage of seeds germinate in the presence of only KAR
TMB affects timing	• germination starts later/slower
TMB affects percentage	• lower percentage of seeds germinate in presence of only TMB

(b) **Make a claim** about the effect of rinsing on the binding of KAR to the receptor in the seed and about the effect of rinsing on the binding of TMB to the receptor in the seed. Identify the appropriate treatment groups and results from the table that, when compared with the controls, **provide support** for each claim. **(4 points)**

Claim (2 points maximum; 1 point for KAR; 1 for TMB)	Support (2 points maximum; 1 point for KAR; 1 for TMB)				
• KAR remains (bound after rinsing) • Rinsing does not affect KAR (binding)	KAR with no rinse	KAR with rinse	different than	Controls	
	Group 2	Group 5		Group 1	Group 4
• TMB does not remain (bound)	TMB with no rinse		different than	Control	
	Group 3			Group 1	
• Rinsing affects TMB (binding)	TMB with rinse		same as	Control	
	Group 6			Group 4	

(c) There is intense competition by plants to successfully colonize areas that have been recently cleared by a fire. **Describe** ONE advantage of KAR regulation and ONE advantage of TMB regulation to plants that live in an ecosystem with regular fires. **(2 points)**

Description (1 point per row; 2 points maximum)	
Advantage of KAR regulation	• Germination occurs at times of increased resources availability. • Plants with KAR regulation can outcompete other plants (without KAR regulation). • Germination occurs when fewer competitors are present/land is barren.
Advantage of TMB regulation	• Plants with TMB regulation do not germinate/can maintain seed dormancy until (enough) water is available. • Plants with TMB regulation do not germinate in a dry environment.

2017 SCORING GUIDELINES

Question 3

Gibberellin is the primary plant hormone that promotes stem elongation. GA 3-beta-hydroxylase (GA3H) is the enzyme that catalyzes the reaction that converts a precursor of gibberellin to the active form of gibberellin. A mutation in the *GA3H* gene results in a short plant phenotype. When a pure-breeding tall plant is crossed with a pure-breeding short plant, all offspring in the F₁ generation are tall. When the F₁ plants are crossed with each other, 75 percent of the plants in the F₂ generation are tall and 25 percent of the plants are short.

		Second Base in Codon					
		U	C	A	G		
First Base in Codon	U	UUU } Phe UUC } UUA } Leu UUG }	UCU } UCC } Ser UCA } UCG }	UAU } Tyr UAC } UAA } Stop UAG } Stop	UGU } Cys UGC } UGA } Stop UGG } Trp	U C A G	Third Base in Codon
	C	CUU } CUC } CUA } Leu CUG }	CCU } CCC } Pro CCA } CCG }	CAU } His CAC } CAA } Gln CAG }	CGU } CGC } CGA } Arg CGG }	U C A G	
	A	AUU } AUC } Ile AUA } AUG } Met or Start	ACU } ACC } Thr ACA } ACG }	AAU } Asn AAC } AAA } Lys AAG }	AGU } AGC } Ser AGA } AGG } Arg	U C A G	
	G	GUU } GUC } GUA } Val GUG }	GCU } GCC } GCA } Ala GCG }	GAU } Asp GAC } GAA } Glu GAG }	GGU } GGC } GGA } Gly GGG }	U C A G	

Figure 1. The universal genetic code

(a) The wild-type allele encodes a GA3H enzyme with alanine (Ala), a nonpolar amino acid, at position 229. The mutant allele encodes a GA3H enzyme with a threonine (Thr), a polar amino acid, at position 229.

Describe the effect of the mutation on the enzyme and **provide reasoning** to support how this mutation results in a short plant phenotype in homozygous recessive plants. **(2 points)**

Description (1 point)	Reasoning (1 point)
The amino acid substitution changes the shape/structure/function of the protein.	The mutation decreases/eliminates gibberellin production.

(b) Using the codon chart provided, **predict** the change in the codon sequence that resulted in the substitution of alanine for threonine at amino acid position 229. **(1 point)**

Prediction (1 point maximum)

- G ↔ A in the first position (of the codon)
- 5'-GCN-3' ↔ 5'-ACN-3'
- 5'-NGC-3' ↔ 5'-NGT-3' in the template strand of DNA

(c) **Describe** how individuals with one (heterozygous) or two (homozygous) copies of the wild-type *GA3H* allele can have the same phenotype. **(1 point)**

Description (1 point)

- Enough active enzyme is produced from one wild-type/dominant allele.
- Enough gibberellin is produced in the presence of one wild-type/dominant allele.

2017 SCORING GUIDELINES

Question 4

DIETARY COMPOSITION OF ORGANISMS IN AN AQUATIC ECOSYSTEM


Organism	Food Source (% of diet)				
	Algae	Stoneflies	Midges	Hellgrammites	Caddisflies
Algae					
Stoneflies			90		10
Midges	100				
Hellgrammites		20	10		70
Caddisflies	70		30		

The table above shows how much each organism in an aquatic ecosystem relies on various food sources. The rows represent the organisms in the ecosystem, and the columns represent the food source. The percentages indicate the proportional dietary composition of each organism. High percentages indicate strong dependence of an organism on a food source.

(a) Based on the food sources indicated in the data table, **construct** a food web in the template below. Write the organism names on the appropriate lines AND draw the arrows necessary to indicate the energy flow between organisms in the ecosystem. (2 points)

Construction of food web (2 points maximum)

- All four organisms placed on the appropriate lines
- All four arrows correctly drawn between organisms



2017 SCORING GUIDELINES

Question 4 (continued)

(b) In an effort to control the number of midges, an area within the ecosystem was sprayed with the fungus *Metarhizium anisopliae*, which significantly decreased the midge population. Based on the data in the table, **predict** whether the spraying of the fungus will have the greatest short-term impact on the population of the stoneflies, the caddisflies, or the hellgrammites. **Justify** your prediction. (2 points)

Prediction (1 point)

- Stoneflies

Justification (1 point)

- Stoneflies have a higher dependence on the midges than do the hellgrammites and caddisflies.
- Midges are 90 percent of the stonefly diet, while 30 percent of the caddisfly and 10 percent of the hellgrammite diet are midges.

2017 SCORING GUIDELINES

Question 5




Figure 1. Characteristics of a pond community over time

Microcystis aeruginosa is a freshwater photosynthetic cyanobacterium. When temperatures increase and nutrients are readily available in its pond habitat, *M. aeruginosa* undergoes rapid cell division and forms an extremely large, visible mass of cells called an algal bloom. *M. aeruginosa* has a short life span and is decomposed by aerobic bacteria and fungi. **Identify** the metabolic pathway and the organism that is primarily responsible for the change in oxygen level in the pond between times I and II AND between times III and IV.

Identification (2 points per row; 4 points maximum)

Time Period	Metabolic pathway (1 point per box)	Organism (1 point per box)
I – II	Photosynthesis	Cyanobacteria (<i>M. aeruginosa</i>)
III – IV	Cellular respiration	Decomposers/fungi/bacteria

2017 SCORING GUIDELINES

Question 6




Figure 1. Comet assay to detect double-stranded breaks in DNA

A comet assay is a technique used to determine the amount of double-strand breaks in DNA (DNA damage) in cells. The nucleus of an individual cell is placed on a microscope slide coated with an agarose gel. An electric current is applied to the gel that causes DNA to move (electrophoresis), and the DNA is stained with a fluorescent dye. When viewed using a microscope, undamaged DNA from the nucleus appears as a round shape (the head), and the fragments of damaged DNA extend out from the head (the tail). The length of the tail corresponds to the amount of the damage in the DNA (see Figure 1).

- (a) To explain the movement of DNA fragments in the comet assay, **identify** one property of DNA and **provide reasoning** to support how the property contributes to the movement during the comet assay technique. (2 points; both points must be earned from the same row.)

Identification (1 point)	Reasoning (1 point)
DNA has a (negative) charge.	DNA moves toward the positive/oppositely charged pole.
DNA can be different sizes.	(Different size DNA fragments) move at different rates.

- (b) In a different experiment, cells are treated with a chemical mutagen that causes only nucleotide substitutions in DNA. **Predict** the likely results of a comet assay for this treatment. (1 point)

Prediction (1 point)

- Head (only) OR (head with) no tail.
- Tail will be shorter than a cell with double-stranded breaks in DNA.

2017 SCORING GUIDELINES

Question 7

Many species of bacteria grow in the mouths of animals and can form biofilms on teeth (plaque). Within plaque, the outer layers contain high levels of oxygen and the layers closest to the tooth contain low levels of oxygen. The surface of the tooth is covered in a hard layer of enamel, which can be dissolved under acidic conditions. When the enamel breaks down, the bacteria in plaque can extract nutrients from the tooth and cause cavities.

Certain types of bacteria (e.g., *Streptococcus mutans*) thrive in the innermost anaerobic layers of the plaque and are associated with cavities. Other types of bacteria (*Streptococcus sanguinis*) compete with *S. mutans* but are unable to thrive in acidic environments.

- (a) **Identify** the biochemical pathway *S. mutans* uses for metabolizing sugar and **describe** how the pathway contributes to the low pH in the inner layers of plaque. (2 points; both points must be earned from the same row.)

Identification	Description
fermentation	(lactic) acid/lactate
anaerobic respiration	acid
glycolysis	(pyruvic) acid/pyruvate

- (b) Normal tooth brushing effectively removes much of the plaque from the flat surfaces of teeth but cannot reach the surfaces between teeth. Many commercial toothpastes contain alkaline components, which raise the pH of the mouth. **Predict** how the population sizes of *S. mutans* AND *S. sanguinis* in the bacterial community in the plaque between the teeth are likely to change when these toothpastes are used. (1 point)

Prediction (1 point)

- S. mutans* decreases AND *S. sanguinis* increases

2017 SCORING GUIDELINES

Question 8

Estrogens are small hydrophobic lipid hormones that promote cell division and the development of reproductive structures in mammals. Estrogens passively diffuse across the plasma membrane and bind to their receptor proteins in the cytoplasm of target cells.

- (a) **Describe** ONE characteristic of the plasma membrane that allows estrogens to passively cross the membrane. (1 point)

Description (1 point)

- Hydrophobic/nonpolar
- Space between phospholipids

- (b) In a laboratory experiment, a researcher generates antibodies that bind to purified estrogen receptors extracted from cells. The researcher uses the antibodies in an attempt to treat estrogen-dependent cancers but finds that the treatment is ineffective. **Explain** the ineffectiveness of the antibodies for treating estrogen-dependent cancers. (2 points)

Explanation (2 points)

- Antibodies are unable to enter the cell.
- (Extracellular) antibodies will not bind to (intracellular) estrogen receptors.

AP[®] Biology

2016 Scoring Guidelines

© 2016 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2016 SCORING GUIDELINES

Question 1




Figure 1. Sampling sites of marine mussels at various locations (1-8) in Long Island Sound and the Atlantic Ocean

TABLE 1. PERCENT OF INDIVIDUALS POSSESSING *lap*⁹⁴ ALLELE

	Long Island Sound					Atlantic Ocean		
Site	1	2	3	4	5	6	7	8
<i>lap</i> ⁹⁴ frequency (%)	13	16	25	37	55	59	59	59
Salinity	Low $\longrightarrow$ High					High		

Leucine aminopeptidases (LAPs) are found in all living organisms and have been associated with the response of the marine mussel, *Mytilus edulis*, to changes in salinity. LAPs are enzymes that remove N-terminal amino acids from proteins and release the free amino acids into the cytosol. To investigate the evolution of LAPs in wild populations of *M. edulis*, researchers sampled adult mussels from several different locations along a part of the northeast coast of the United States, as shown in Figure 1. The researchers then determined the percent of individuals possessing a particular *lap* allele, *lap*⁹⁴, in mussels from each sample site (table 1).

2016 SCORING GUIDELINES

Question 1 (continued)

- (a) On the axes provided, **construct** an appropriately labeled bar graph to illustrate the observed frequencies of the *lap⁹⁴* allele in the study populations. **(3 points)**

Construct graph (3 points)

- Correctly plotted bar graph that accurately represents the trend
- Correct axis labeling
- Correct scale and units

- (b) Based on the data, **describe** the most likely effect of salinity on the frequency of the *lap⁹⁴* allele in the marine mussel populations in Long Island Sound. **Predict** the likely *lap⁹⁴* allele frequency at a sampling site between site 1 and site 2 in Long Island Sound. **(2 points)**

Description (1 point)	Prediction (1 point)
• As salinity increases <i>lap⁹⁴</i> frequency increases • As salinity decreases <i>lap⁹⁴</i> frequency decreases	Between 13 and 16 percent (or a selected value between 13 and 16 percent)

- (c) **Describe** the most likely effect of LAP⁹⁴ activity on the osmolarity of the cytosol. **Describe** the function of LAP⁹⁴ in maintaining water balance in the mussels living in the Atlantic Ocean. **(2 points)**

Describe effect of LAP ⁹⁴ activity (1 point)	Describe function of LAP ⁹⁴ in maintaining water balance (1 point)
• LAP⁹⁴ increases osmolarity/solute concentration of the cytosol • LAP⁹⁴ decreases water potential of the cytosol	Prevents water loss to the environment

- (d) Marine mussel larvae are evenly dispersed throughout the study area by water movement. As larvae mature, they attach to the rocks in the water. **Explain** the differences in *lap⁹⁴* allele frequency among adult mussel populations at the sample sites despite the dispersal of larvae throughout the entire study area. **Predict** the likely effect on distribution of mussels in Long Island Sound if the *lap⁹⁴* allele was found in all of the mussels in the population. **Justify** your prediction. **(3 points)**

Explanation (1 point)	Prediction (1 point)	Justification (1 point)
• Mussels with <i>lap⁹⁴</i> allele are more likely to survive in high salinity/less likely to survive in low salinity. • Mussels without <i>lap⁹⁴</i> allele are less likely to survive in high salinity/more likely to survive in low salinity.	• Mussel population will increase in high salinity. • Mussel population will decline in low salinity.	• Mussels in high salinity with <i>lap⁹⁴</i> allele will osmoregulate. • Mussels in low salinity with <i>lap⁹⁴</i> allele will not osmoregulate.

2016 SCORING GUIDELINES

Question 2




Figure 1. Bacterial population growth in the presence of two nutrients (nutrient I and nutrient II)

Bacteria can be cultured in media with carefully controlled nutrient composition. The graph above shows the growth of a bacterial population in a medium with limiting amounts of two nutrients, I and II.

- (a) **Estimate** the maximum population density in $\frac{\text{cells}}{\text{mL}}$ for the culture. Using the data, **describe** what prevents further growth of the bacterial population in the culture. **(2 points)**

Estimate (1 point)

- 10^8

Description (1 point)

- When both nutrients are depleted

- (b) Using the data, **calculate** the growth rate in $\frac{\text{cells}}{\text{mL} \times \text{hour}}$ of the bacterial population between hours 2 and 4. **(1 point).**

Calculation (1 point)

- 4,995

2016 SCORING GUIDELINES

Question 2 (continued)

- (c) **Identify** the preferred nutrient source of the bacteria in the culture over the course of the experiment. Use the graph to **justify** your response. **Propose** ONE advantage of the nutrient preference for an individual bacterium. **(3 points)**

Identification (1 point)

- Nutrient I is the preferred nutrient.

Justification (1 point)

- When both nutrients are present in the growth medium, only nutrient I is used.
- Nutrient II is only used after nutrient I is depleted.

Proposed advantage (1 point)

- Do not spend energy making enzymes/proteins that the cell doesn't need.
- Do not have to express all metabolic genes at once.
- The preferred nutrient provides more energy.

- (d) **Describe** how nutrient I most likely regulates the genes for metabolism of nutrient I and the genes for metabolism of nutrient II. **Provide** TWO reasons that the population does not grow between hours 5 and 6. **(4 points)**

Description (2 points)

- Nutrient I promotes expression of genes required for metabolism of nutrient I.
- Nutrient I represses expression of genes required for metabolism of nutrient II.

Reasoning (2 points)

- Nutrient I is depleted from the growth medium OR neither nutrient is being consumed.
- Takes time to produce proteins/enzymes required to metabolize nutrient II.

2016 SCORING GUIDELINES

Question 3




Figure 1. Percent dry weight of different plant structures during the growing season for an annual plant

The graph above illustrates the percent dry weight of different parts of a particular annual plant (plants that live less than one year) from early May to late August. The percent dry weight can be used to estimate the amount of energy a plant uses to produce its leaves, vegetative buds, stems, roots, and reproductive parts (seeds, receptacles, and flowers).

- (a) **Identify** the direct source of the energy used for plant growth during the first week of May, and **identify** the part of the plant that grew the most during the same period. **(2 points)**

Identify direct source of energy (1 point)

- Seed
- Stored organic nutrients/carbohydrates

Identify plant part (1 point)

- Roots

- (b) Based on the data on the graph, **estimate** the percent of the total energy that the plant has allocated to the growth of leaves on the first day of July. **(1 point)**

Identification (1 point)

- Any value between 45-55 percent


- (c) Compared with perennials (plants that live more than two years), annual plants often allocate a much greater percentage of their total energy to growth of their reproductive parts in any given year. **Propose** ONE evolutionary advantage of the energy allocation strategy in annual plants compared with that in perennial plants. **(1 point)**

Proposed advantage (1 point)

- Increased chance of reproduction before the plants die.
- If the plants do not use the strategy, they decrease the likelihood they will ever reproduce.

2016 SCORING GUIDELINES

Question 4



The figure represents the process of expression of gene X in a eukaryotic cell.

- (a) The primary transcript in the figure is 15 kilobases (kb) long, but the mature mRNA is 7 kb in length. **Describe** the modification that most likely resulted in the 8 kb difference in length of the mature mRNA molecule. **Identify** in your response the location in the cell where the change occurs. (2 points)

Describe process (1 point)

- Removal of introns
- RNA processing

Identification (1 point)

- Nucleus

- (b) **Predict** the length of the mature gene X mRNA if the full-length gene is introduced and expressed in prokaryotic cells. **Justify** your prediction. (2 points)

Prediction (1 point)


- 15 kb
- Longer than the mature mRNA in the eukaryote

Justification (1 point)

- mRNA processing typically does not occur in prokaryotes

2016 SCORING GUIDELINES


Question 5




The graph above shows the mass of plants from two different species over time. The plants grew while attached to each other. The plants were separated at the time indicated by the vertical line in the graph.

Using template 1, **graph** the predicted shape of the plant-mass lines after separation of the two plants if the plants were in an obligate mutualistic relationship. On template 2, **graph** the predicted shape of the plant-mass lines if the species 2 plant was a parasite of the species 1 plant. **Justify** each of your predictions. (4 points)

TEMPLATE 1: OBLIGATE MUTUALISM



TEMPLATE 2: PARASITISM



	Graph characteristics (1 point each graph; 2 points maximum)	Justification (1 point each box; 2 points maximum)
Obligate Mutualism	Both of the growth curves level off or decline.	• Each species depends on the other for survival. • Without the relationship, both species are harmed.
Parasitism	Species 1 continues to increase while species 2 levels off or declines.	• The parasite requires an association with the host to survive but harms the host. • Without the relationship, the parasite cannot survive while the host continues to grow.

2016 SCORING GUIDELINES

Question 6




Figure 1. Detectability of eDNA fragments of varying lengths

Living and dead organisms continuously shed DNA fragments, known as eDNA, into the environment. To detect eDNA fragments in the environment, the polymerase chain reaction (PCR) can be used to amplify specific eDNA fragments. eDNA fragments of different lengths persist in the environment for varying amounts of time before becoming undetectable (Figure 1).

To investigate whether silver carp, an invasive fish, have moved from a nearby river system into Lake Michigan, researchers tested water samples for the presence of eDNA specific to silver carp (Figure 2).

- (a) **Justify** the use of eDNA sampling as an appropriate technique for detecting the presence of silver carp in an environment where many different species of fish are found. **Propose** ONE advantage of identifying long eDNA fragments as opposed to short fragments for detecting silver carp. **(2 points)**

Justify (1 point)

- eDNA allows detection of the fish without visual identification/catching the fish.

Proposed advantage (1 point)

- Longer fragments indicate more recent presence of fish.
- Longer fragments are more likely to contain a sequence that is specific to silver carp.
- Longer sequences/more base pairs may increase accuracy/specificity/confidence that the eDNA is from a silver carp and not a related species.

- (b) The researchers tested a large number of water samples from Lake Michigan and found eDNA specific to silver carp in a single sample in the lake, as indicated in Figure 2. The researchers concluded that the single positive sample was a false positive and that no silver carp had entered Lake Michigan.

Provide reasoning other than human error to support the researchers' claim. **(1 point)**

Reasoning (1 point)

- eDNA entered the lake by means other than the fish (e.g., river flow, boats, waste from predators).





Figure 2. Map of the waterways that connect a nearby river system to Lake Michigan

2016 SCORING GUIDELINES

Question 7




In a certain species of plant, the diploid number of chromosomes is 4 ($2n = 4$). Flower color is controlled by a single gene in which the green allele (G) is dominant to the purple allele (g). Plant height is controlled by a different gene in which the dwarf allele (D) is dominant to the tall allele (d). Individuals of the parental (P) generation with the genotypes $GGDD$ and $ggdd$ were crossed to produce F_1 progeny.

- (a) **Construct** a diagram below to depict the four possible normal products of meiosis that would be produced by the F_1 progeny. Show the chromosomes and the allele(s) they carry. Assume the genes are located on different chromosomes and the gene for flower color is on chromosome 1. **(1 point)**

Construct diagram (1 point)

- Diagram must include all of the following:
 - Each cell has one unduplicated chromosome 1 (with G or g).
 - Each cell has one unduplicated chromosome 2 (with D or d).
 - Genotype combinations should be: GD , Gd , gD , gd .



- (b) **Predict** the possible phenotypes and their ratios in the offspring of a testcross between an F_1 individual and a $ggdd$ individual. **(1 point)**

Prediction (1 point)

- 1 green dwarf: 1 green tall: 1 purple dwarf: 1 purple tall

- (c) If the two genes were genetically linked, **describe** how the proportions of phenotypes of the resulting offspring would most likely differ from those of the testcross between an F_1 individual and a $ggdd$ individual. **(1 point)**

Identify difference (1 point)

- The majority/greater than 50 percent would have the parental plant phenotypes
- Greater than 25 percent would be green dwarf plants and greater than 25 percent would be purple tall plants
- Less than 25 percent would be green tall plants and less than 25 percent would be purple dwarf plants

2016 SCORING GUIDELINES

Question 8




Figure 1. Effect of exercise on blood prolactin levels in adult males. The data represent the mean $\pm 2SE_{\bar{x}}$.

Researchers conducted a study to investigate the effect of exercise on the release of prolactin into the blood. The researchers measured the concentration of prolactin in the blood of eight adult males before (T = 0 hour) and after one hour (T = 1 hour) of vigorous exercise. As a control, the researchers measured the concentration of blood prolactin in the same group of individuals at the same times of day one week later, but without having them exercise. The results are shown in Figure 1.

- (a) **Justify** the use of the without-exercise treatment as the control in the study design. **(1 point)**

Justification (1 point)

- Attribute changes in the concentration of blood prolactin to exercise only
- Rule out normal fluctuations in prolactin release/levels

- (b) Using evidence from the specific treatments, **determine** whether prolactin release changes after exercise. **Justify** your answer. **(2 points)**

Determination (1 point)

- Exercise does not affect prolactin release

Justification (1 point)

- The T=1 hour with-exercise mean and the T=1 hour without-exercise mean are within $\pm 2SE_{\bar{x}}$.
- The $\pm 2SE_{\bar{x}}$ error bars for the T=1 hour with-exercise time point and the T=1 hour time without-exercise point overlap.
- The $\pm 2SE_{\bar{x}}$ error bars for the T=0 and T=1 hour with-exercise time points overlap.
- The T=0 hour with-exercise mean and the T=1 hour with exercise-mean are within $\pm 2SE_{\bar{x}}$.