

Question 6: Analyze Data**4 points**

Housekeeping genes encode proteins involved in universally important processes such as transcription, translation, and glycolysis. Because these genes appear to be expressed in all cells at constant levels, the expression of housekeeping genes is often used as a control when comparing how the expression of other genes varies under different conditions.

Researchers studying the effect of pesticides on declining bee populations wanted to determine whether the expression of four housekeeping genes (*GAPDH*, *RPL32*, *RPS5*, and *TBP-AF*) was in fact constant in bees across different variables. The researchers collected samples of mRNA for each of the four genes and compared how their expression varied across the developmental stage of the bee, the sex of the bee, and the cell type from which the sample was taken. The mRNA from the samples was reverse transcribed to produce DNA copies of each gene. PCR was then used to amplify the DNA, and the Cq value was determined. The Cq value is the number of PCR cycles needed to produce a specified number of DNA copies (Figure 1). A high Cq value for a sample indicates the gene was expressed at a low level.

To analyze whether any of the examined variables affected expression of the housekeeping genes, researchers examined the range of Cq values for each gene in response to each variable. Genes with a wide range of Cq values were determined to be affected by the variable, while genes with a narrow range of Cq values were determined to be unaffected by the variable.




Figure 1. The effect of developmental stage, sex, and cell type on the Cq value of four housekeeping genes

- | | | |
|-----|--|----------------|
| (a) | Based on the data in <u>Figure 1</u> , identify the gene that had the lowest median Cq value when bees of different developmental stages were compared. | 1 point |
| | <ul style="list-style-type: none"> • <i>RPS5</i> | |
| (b) | The Cq value is inversely proportional to the amount of mRNA from that gene in the starting sample. Based on the data in <u>Figure 1</u> , identify the gene that has the lowest level of gene expression regardless of variable. | 1 point |
| | <ul style="list-style-type: none"> • <i>TBP-AF</i> | |
| (c) | The scientists investigated the effect of pesticides on the expression of other genes in one cell type of a group of bees containing males and females of the same developmental stage. They hypothesized that <i>TBP-AF</i> would serve as the best control gene for this experiment. Use the data to evaluate their hypothesis. | 1 point |
| | <ul style="list-style-type: none"> • Their hypothesis is supported because <i>TBP-AF</i> has the <u>smallest Cq range/most constant expression</u> (when comparing sexes). | |
| (d) | Explain how expression of a gene such as <i>GAPDH</i> can vary from one cell type to another within the same bee. | 1 point |
| | <ul style="list-style-type: none"> • Different cell types contain <u>different levels of/different</u> transcription factors, and therefore regulate the expression of genes in different ways. | |

Total for question 6 4 points

2019

AP[®]

CollegeBoard

2019 SCORING GUIDELINES

Question 1




Figure 1. Model of two-step enzymatic plant pathway for synthesis of IAA from tryptophan

Auxins are plant hormones that coordinate several aspects of root growth and development. Indole-3-acetic acid (IAA) is an auxin that is usually synthesized from the amino acid tryptophan (Figure 1). Gene *Trp-T* encodes an enzyme that converts tryptophan to indole-3-pyruvic acid (I3PA), which is then converted to IAA by an enzyme encoded by the gene *YUC*.

- (a) **Circle ONE** arrow that represents transcription on the template pathway. **Identify** the molecule that would be absent if enzyme YUC is nonfunctional.

Circle (1 point)

- Circle around either arrow pointing from a gene (*Trp-T* or *YUC*) to mRNA

Identification (1 point)

- IAA

- (b) **Predict** how the deletion of one base pair in the fourth codon of the coding region of gene *Trp-T* would most likely affect the production of IAA. **Justify** your prediction.

Prediction (1 point)

- Reduction in IAA production OR No production of IAA

Justification (1 point)

- The mutation will result in the translation of an inactive/nonfunctional Trp-T enzyme.
- The mutation will result in no translation of the Trp-T enzyme.
- The mutation will result in no/reduced production of I3PA.

- (c) **Explain** one feedback mechanism by which a cell could prevent production of too much IAA without limiting I3PA production.

Explanation (2 points)

- Negative feedback/feedback inhibition/increasing amounts of IAA inhibits the pathway.
- Production of YUC enzyme is inhibited OR YUC enzyme activity is inhibited.

2019 SCORING GUIDELINES**Question 1 (continued)**

- (d) Rhizobacteria are a group of bacteria that live in nodules on plant roots. Rhizobacteria can produce IAA and convert atmospheric nitrogen into forms that can be used by plants. Plants release carbon-containing molecules into the nodules. Based on this information, **identify** the most likely ecological relationship between plants and rhizobacteria. **Describe** ONE advantage to the bacteria of producing IAA.

Identification (1 point)

- Mutualism

Description (1 point)

- Increases habitat/number of nodules for the rhizobacteria.
- The bacteria receive carbon/carbon-containing molecules (as a result of increased plant growth).

- (e) A researcher removed a plant nodule and identified several “cheater” rhizobacteria that do not produce IAA or fix nitrogen. **Describe** the evolutionary advantage of being a bacterial cheater in a population composed predominantly of noncheater bacteria. Plants can adjust the amount of carbon-containing molecules released into nodules in response to the amount of nitrogen fixed in the nodule. **Predict** the change in the bacterial population that would cause the plant to reduce the amount of carbon-containing molecules provided to the nodule.

Description (1 point)

- Cheaters/bacteria that benefit without producing IAA/fixing nitrogen have more energy for reproduction.

Prediction (1 point)

- Decrease in the nitrogen-fixing/noncheater bacteria
- Decrease in the amount of nitrogen fixed (by bacteria)

2019 SCORING GUIDELINES

Question 2

A student studying two different aquatic, plant-eating, unicellular protist species (species A and B) designed an experiment to investigate the ecological relationship between the two species (Table 1).

TABLE 1. EXPERIMENTAL TREATMENT GROUPS

Group I.	Species A and B are each grown in separate containers.
Group II.	Species A and B are grown together in the same container.

In treatment group I, the student placed 10 individuals of species A into a container with liquid growth medium and 10 individuals of species B into a separate container with an equal amount of the same liquid growth medium. In treatment group II, the student placed 5 individuals of each species into a single container with the liquid growth medium. The student then maintained the containers under the same environmental conditions and recorded the number of individuals in each population at various time points. The results are shown in Table 2.

TABLE 2. NUMBER OF INDIVIDUALS IN EACH PROTIST POPULATION IN BOTH TREATMENT GROUPS

Time (h)	Group I. Grown Separately		Group II. Grown Together	
	Species A	Species B	Species A	Species B
0	10	10	5	5
10	100	50	45	20
20	400	200	100	50
30	1100	500	250	25
40	1400	650	525	20
50	1500	700	900	10
60	1500	700	1250	0
70	1500	700	1400	0

(a) The growth curves for species B in group I and for species A in group II (shaded columns) have been plotted on the template. Use the template to **complete** an appropriately labeled line graph to illustrate the growth of species A in treatment group I and species B in treatment group II (unshaded columns).

Completion (3 points)

- Correctly plotted lines for remaining two treatments
- Correctly labeled axes including units
- Correctly labeled data lines

2019 SCORING GUIDELINES**Question 2 (continued)**

(b) As shown in the table, the researcher established treatment group II with 5 individuals of each species. **Provide reasoning** for the reduced initial population sizes.

Reasoning (1 point)

- Reduced initial population sizes keep the total number of organisms the same in all containers.
- Reduced initial population sizes serve as a control for population density.

(c) The student claims that species A and B compete for the same food source. **Provide TWO pieces of evidence** from the data that support the student's claim.

Evidence (1 point per row; 2 points max.)

Comparison of Groups	Evidence
I-A to II-A	• Growth rate is faster in I/slower in II
I-A to II-A	• Grows to a higher population density in I/lower population density in II
I-B to II-B	• Growth rate is faster in I/slower in II
I-B to II-B	• Grows to a higher population density in I/lower population density in II/ II dies out/II goes to zero

(d) **Predict** TWO factors that most likely limit the population growth of species A in treatment group I.

Prediction (2 points)

Acceptable factors include:

- Food
- Space
- Metabolic waste
- Dissolved oxygen

(e) Many protists contain an organelle called a contractile vacuole that pumps water out of the cell. The student repeated the experiment using a growth medium with a lower solute concentration. **Predict** how the activity of the contractile vacuole will change under the new experimental conditions. **Justify** your prediction.

Prediction (1 point)

- The contractile vacuole will be more active.

Justification (1 point)

- The environment is hypotonic with respect to the cell.
- The cell is hypertonic with respect to environment.
- Water has entered the cell (which could cause lysis).
- The cell has lower water potential than the environment/the environment has higher water potential than the cell.

2019 SCORING GUIDELINES**Question 3**

The pyruvate dehydrogenase complex (PDC) catalyzes the conversion of pyruvate to acetyl-CoA, a substrate for the Krebs (citric acid) cycle. The rate of pyruvate conversion is greatly reduced in individuals with PDC deficiency, a rare disorder.

(a) **Identify** the cellular location where PDC is most active.

Identification (1 point)

- Mitochondria
- Mitochondrial matrix

(b) **Make a claim** about how PDC deficiency affects the amount of NADH produced by glycolysis AND the amount of NADH produced by the Krebs (citric acid) cycle in a cell. **Provide reasoning** to support your claims based on the position of the PDC-catalyzed reaction in the sequence of the cellular respiration pathway.

(1 point per row; 2 points max.)

	Claim	Reasoning
Glycolysis	No change	• Glycolysis continues; PDC is not needed. • Glycolysis occurs before conversion of pyruvate to acetyl-CoA.
Krebs cycle	Decrease	• The Krebs cycle is greatly reduced/slowed down if there is no/less acetyl-CoA. • The Krebs cycle occurs after conversion of pyruvate to acetyl-CoA.

(c) PDC deficiency is caused by mutations in the *PDHA1* gene, which is located on the X chromosome. A male with PDC deficiency and a homozygous female with no family history of PDC deficiency have a male offspring. **Calculate** the probability that the male offspring will have PDC deficiency.

Calculation (1 point)

- The probability of inheritance is 0.
- The offspring cannot/will not have PDC deficiency.

2019 SCORING GUIDELINES

Question 4




Figure 1. Release of neurotransmitters into the synapse in response to an action potential




Figure 2. Model of a typical action potential in a neuron

Acetylcholine is a neurotransmitter that can activate an action potential in a postsynaptic neuron (Figures 1 and 2). A researcher is investigating the effect of a particular neurotoxin that causes the amount of acetylcholine released from presynaptic neurons to increase.

(a) **Describe** the immediate effect of the neurotoxin on the number of action potentials in a postsynaptic neuron. **Predict** whether the maximum membrane potential of the postsynaptic neuron will increase, decrease, or stay the same.

Description (1 point)

- It will increase the number of action potentials.

Prediction (1 point)

- It will stay the same.

(b) The researcher proposes two models, A and B, for using acetylcholinesterase (AChE), an enzyme that degrades acetylcholine, to prevent the effect of the neurotoxin. In model A, AChE is added to the synapse. In model B, AChE is added to the cytoplasm of the postsynaptic cell. **Predict** the effectiveness of EACH proposed model. **Provide reasoning** to support your predictions.

(1 point per row; 2 points max.)

	Prediction	Reasoning
Model A	Effective	Acetylcholine is in the synapse.
Model B	Not effective	Acetylcholine is not in the cytoplasm of the postsynaptic cell.

2019 SCORING GUIDELINES

Question 5

TABLE 1. DIVERGENCE (IN PERCENT) OF MITOCHONDRIAL DNA SEQUENCES AMONG FIVE PRIMATE SPECIES

	Human	Gorilla	Orangutan	Gibbon	Chimpanzee
Human	-	10.3	16.1	18.1	8.8
Gorilla		-	16.7	18.9	10.6
Orangutan			-	18.9	17.2
Gibbon				-	18.9
Chimpanzee					-


A researcher studying the evolutionary relationship among five primate species obtained data from a sequence of mitochondrial DNA (mtDNA) from a representative individual of each species. The researcher then calculated the percent divergence in the sequences between each pair of primate species (Table 1).

(a) Based on fossil data, the researcher estimates that humans and their most closely related species in the data set diverged approximately seven million years ago. Using these data, **calculate** the rate of mtDNA percent divergence per million years between humans and their most closely related species in the data set. Round your answer to two decimal places.

Calculation (1 point)

- 1.25 OR 1.26

(b) Using the data in the table, **construct** a cladogram on the template provided. **Provide reasoning** for the placement of gibbons as the outgroup on the cladogram.

**Construction (1 point)**

- From top to bottom: Human/Chimpanzee (interchangeable), Gorilla, Orangutan

Reasoning (1 point)

- Gibbon mtDNA is the least similar (to all of the other species)/most different (from all of the other species).
- Gibbon mtDNA is the most divergent (from all of the other species).

2019 SCORING GUIDELINES**Question 5 (continued)**

(c) On the cladogram, **draw** a circle around all of the species that are descended from the species indicated by the node within the square.

Circle (1 point)

- Circle species 1, 2, and 3, as numbered from the top.

2019 SCORING GUIDELINES

Question 6

	MEDIUM	STRAINS		
		Wild Type	Mutant 1	Mutant 2
Treatment I	All amino acids present	+	+	+
Treatment II	No amino acids present	+	–	–
Treatment III	All amino acids present EXCEPT methionine	+	–	+
Treatment IV	All amino acids present EXCEPT leucine	+	+	–

Table 1. The data show the growth of haploid *Saccharomyces cerevisiae* yeast strains on media that differ in amino acid content. A plus sign (+) indicates that the yeast strains grow, and a minus sign (–) indicates that the strains do not grow.

The yeast *Saccharomyces cerevisiae* is a single-celled organism. Amino acid synthesis in yeast cells occurs through metabolic pathways, and enzymes in the synthesis pathways are encoded by different genes. The synthesis of a particular amino acid can be prevented by mutation of a gene encoding an enzyme in the required pathway.

A researcher conducted an experiment to determine the ability of yeast to grow on media that differ in amino acid content. Yeast can grow as both haploid and diploid cells. The researcher tested two different haploid yeast strains (Mutant 1 and Mutant 2), each of which has a single recessive mutation, and a haploid wild-type strain. The resulting data are shown in Table 1.

(a) **Identify** the role of treatment I in the experiment.

Identification (1 point)

- (Positive) control (for yeast growth).
- To test the viability of all yeast strains.
- Treatment I allows the researcher to be confident that changes in experimental outcome are due to differences in treatments.

(b) **Provide reasoning** to explain how Mutant 1 can grow on treatment I medium but cannot grow on treatment III medium.

Reasoning (1 point)

- Mutant 1 can use methionine when it is present in the medium, but Mutant 1 cannot synthesize methionine.

(c) Yeast mate by fusing two haploid cells to make a diploid cell. In a second experiment, the researcher mates the Mutant 1 and Mutant 2 haploid strains to produce diploid cells. Using the table provided, **predict** whether the diploid cells will grow on each of the four media. Use a plus sign (+) to indicate growth and a minus sign (–) to indicate no growth.

Prediction (1 Point)

- There will be growth (+) in all four cells of the fourth column.

2019 SCORING GUIDELINES

Question 7

A researcher is studying patterns of gene expression in mice. The researcher collected samples from six different tissues in a healthy mouse and measured the amount of mRNA from six genes. The data are shown in Figure 1.




Figure 1. mRNA expression levels of six genes

(a) Based on the data provided, **identify** the gene that is most likely to encode a protein that is an essential component of glycolysis. **Provide reasoning** to support your identification.

Identification (1 point)

- *Gene G*

Reasoning (1 point)

- (*Gene G*) is the only gene expressed in all (six) tissues, AND glycolysis occurs in all (six) tissues.
- (*Gene G*) mRNA is the only mRNA present in all (six) tissues, AND glycolysis occurs in all (six) tissues.

(b) The researcher observed that tissues with a high level of *gene H* mRNA did not always have *gene H* protein. **Provide reasoning** to explain how tissues with high *gene H* mRNA levels can have no *gene H* protein.



Reasoning (1 point)

- The mRNA is not exported from the nucleus.
- *Gene H* mRNA is not translated/RNA interference prevent(s) translation.
- Post-transcriptional modifications.

2019 SCORING GUIDELINES

Question 8

TABLE 1. CHANGES IN MORNING GLORY PETAL CELLS DURING FLOWER OPENING

	BUD	OPEN FLOWER
		
Vacuole pH	6.6	7.7
Flower Color	Red	Blue
Cell Volume	Small	Large

The petal color of the Mexican morning glory (*Ipomoea tricolor*) changes from red to blue, and the petal cells swell during flower opening. The pigment heavenly blue anthocyanin is found in the vacuole of petal cells. Petal color is determined by the pH of the vacuole. A model of a morning glory petal cell before and after flower opening is shown in Table 1.

(a) **Identify** the cellular component in the model that is responsible for the increase in the pH of the vacuole during flower opening AND **describe** the component's role in changing the pH of the vacuole.

Identification (1 point)

- (K^+ / H^+) transport protein

Description (1 point)

- It moves H^+ out of the vacuole.

(b) A researcher claims that the activation of the K^+ / H^+ transport protein causes the vacuole to swell with water. **Provide reasoning** to support the researcher's claim.

Reasoning (1 point)

- The concentration of solute (K^+) is increasing inside the vacuole.
- The solute (K^+) is moving into the vacuole, making it hypertonic/hyperosmotic/lowering water potential.

AP[®] Biology

2015 Scoring Guidelines

© 2015 The College Board. College Board, Advanced Placement Program, AP, AP Central, and the acorn logo are registered trademarks of the College Board.

Visit the College Board on the Web: www.collegeboard.org.

AP Central is the official online home for the AP Program: apcentral.collegeboard.org.

2015 SCORING GUIDELINES

Question 1

Many species have circadian rhythms that exhibit an approximately 24-hour cycle. Circadian rhythms are controlled by both genetics and environmental conditions, including light.

Researchers investigated the effect of light on mouse behavior by using a running wheel with a motion sensor to record activity on actograms, as shown in Figure 1.




Figure 1. Strategy for recording mouse activity data. When a mouse is active on the running wheel, the activity is recorded as a dark horizontal line on an actogram. When the mouse is inactive, no dark line is recorded.

For the investigation, adult male mice were individually housed in cages in a soundproof room at 25°C. Each mouse was provided with adequate food, water, bedding material, and a running wheel. The mice were exposed to daily periods of 12 hours of light (L) and 12 hours of dark (D) (L12:D12) for 14 days, and their activity was continuously monitored. The activity data are shown in Figure 2.

2015 SCORING GUIDELINES

Question 1 (continued)




Figure 2. Actogram of mouse activity under L12:D12 conditions. Each row represents a 24-hour period, and the dark horizontal lines represent activity on the running wheel.

After 14 days in L12:D12, the mice were placed in continuous darkness (DD), and their activity on the running wheel was recorded as before. The activity data under DD conditions are shown in Figure 3.




Figure 3. Actogram of mouse activity under DD conditions. Each row represents a 24-hour period, and the dark horizontal lines represent activity on the running wheel.

2015 SCORING GUIDELINES

Question 1 (continued)

- (a) The nervous system plays a role in coordinating the observed activity pattern of the mice in response to light-dark stimuli. **Describe** ONE role of each of the following anatomical structures in responding to light-dark stimuli.

- A photoreceptor in the retina of the eye
- The brain
- A motor neuron

	Descriptions (1 point per box; 3 points maximum)
Photoreceptor	Detects light/dark stimulus and initiates/transmits signal
Brain	Integrates/processes/coordinates information
Motor neuron	Transmits signal from brain to an effector

- (b) Based on an analysis of the data in Figure 2, **describe** the activity pattern of the mice during the light and dark periods of the L12:D12 cycle.

Description (1 point)

- Active during dark phase AND inactive during light phase
- Active ONLY during the dark period
- Inactive ONLY during the light period

- (c) The researchers claim that the genetically controlled circadian rhythm in the mice does not follow a 24-hour cycle. **Describe** ONE difference between the daily pattern of activity under L12:D12 conditions (Figure 2) and under DD conditions (Figure 3), and use the data to **support** the researchers' claim.

Description (1 point)

- Active period begins a little earlier each day
- Active/inactive period is shorter than 12 hours each day
- Daily circadian rhythm is less than 24 hours
- Pattern of activity shifts each day

Support (1 point)

- Without light, active/inactive periods are determined only by the genetically controlled circadian rhythm.
- If it were a 24-hour circadian rhythm, the pattern of activity in DD would be the same as the pattern of activity in L12:D12.

- (d) To investigate the claim that exposure to light overrides the genetically controlled circadian rhythm, the researchers plan to repeat the experiment with mutant mice lacking a gene that controls the circadian rhythm. **Predict** the observed activity pattern of the mutant mice under L12:D12 conditions and under DD conditions that would support the claim that light overrides the genetically controlled circadian rhythm.

2015 SCORING GUIDELINES

Question 1 (continued)

Conditions	Predicted Activity Pattern (1 point per box; 2 points maximum)
Mutant under L12:D12	Normal rhythm/rhythm similar to wild-type mouse under L12:D12 (Figure 2)
Mutant under DD	• Random activity throughout the 24 hour period • No pattern/rhythm • Constantly active/constantly inactive

- (e) In nature, mice are potential prey for some predatory birds that hunt during the day. **Describe** TWO features of a model that represents how the predator-prey relationship between the birds and the mice may have resulted in the evolution of the observed activity pattern of the mice.

	Description (1 point per box; 2 points maximum)
Selective Advantage	• Selection for individuals active at night • Selection against individuals active during the day • Day-active variants susceptible to predation • Night-active variants able to avoid predation
Reproductive Success	• Mice selected for produce more offspring • Mice selected against produce fewer offspring

2015 SCORING GUIDELINES

Question 2




Figure 1. Glycolysis and pyruvate oxidation




Figure 2. Krebs cycle




Figure 3. Electron transport chain

Cellular respiration includes the metabolic pathways of glycolysis, the Krebs cycle, and the electron transport chain, as represented in the figures. In cellular respiration, carbohydrates and other metabolites are oxidized, and the resulting energy-transfer reactions support the synthesis of ATP.

(a) Using the information above, **describe** ONE contribution of each of the following in ATP synthesis.

- Catabolism of glucose in glycolysis and pyruvate oxidation
- Oxidation of intermediates in the Krebs cycle
- Formation of a proton gradient by the electron transport chain

Process	Description (1 point each box; 3 points maximum)
Catabolism of glucose in glycolysis and pyruvate oxidation	• Produces NADH for use in ETC • Produces acetyl-CoA for entry into Krebs cycle • Provides energy for (substrate level) phosphorylation of ADP
Oxidation of intermediates in the Krebs cycle	• Produces NADH or FADH₂ for use in ETC • Releases high energy electrons for use in ETC • Provides energy to pump protons against their concentration gradient • Produces GTP for (substrate level) phosphorylation of ADP
Formation of a proton gradient by the electron transport chain	• The flow of protons through membrane-bound ATP synthase generates ATP • Provides energy for (oxidative) phosphorylation of ADP

2015 SCORING GUIDELINES

Question 2 (continued)

(b) Use each of the following observations to **justify** the claim that glycolysis first occurred in a common ancestor of all living organisms.

- Nearly all existing organisms perform glycolysis.
- Glycolysis occurs under anaerobic conditions.
- Glycolysis occurs only in the cytosol.

Observation	Justification (1 point each box; 3 points maximum)
Nearly all existing organisms perform glycolysis	• Trait/gene/process originated early and was inherited/passed down/highly conserved • Glycolysis provided a selective advantage that was passed on to descendants
Glycolysis occurs under anaerobic conditions	Origin of glycolysis pre-dates free atmospheric oxygen/photosynthesis
Glycolysis occurs only in the cytosol	Origin of glycolysis pre-dates cell types with membrane-bound organelles/eukaryotes/endosymbiosis

(c) A researcher estimates that, in a certain organism, the complete metabolism of glucose produces 30 molecules of ATP for each molecule of glucose. The energy released from the total oxidation of glucose under standard conditions is 686 kcal/mol. The energy released from the hydrolysis of ATP to ADP and inorganic phosphate under standard conditions is 7.3 kcal/mol. **Calculate** the amount of energy available from the hydrolysis of 30 moles of ATP. **Calculate** the efficiency of total ATP production from 1 mole of glucose in the organism. **Describe** what happens to the excess energy that is released from the metabolism of glucose.

	Calculation/description (1 point each box; 3 points maximum)
Calculate available energy in ATP	219 kcal
Calculate efficiency	0.31 - 0.32 or 31 - 32%
Describe fate of excess energy	Released as heat/increases entropy

(d) The enzymes of the Krebs cycle function in the cytosol of bacteria, but among eukaryotes the enzymes function mostly in the mitochondria. **Pose** a scientific question that connects the subcellular location of the enzymes in the Krebs cycle to the evolution of eukaryotes.

Question (1 point)

- A valid scientific question related to evolution of eukaryotes (e.g., Since the Krebs cycle occurs in the “cytoplasm” of the mitochondria (matrix), does it suggest that mitochondria were once prokaryotes?)

2015 SCORING GUIDELINES

Question 3

The amino-acid sequence of cytochrome *c* was determined for five different species of vertebrates. The table below shows the number of differences in the sequences between each pair of species.


THE NUMBER OF AMINO ACID DIFFERENCES
IN CYTOCHROME *c* AMONG FIVE SPECIES

	<i>E. ferus</i>	<i>D. polylepis</i>	<i>G. gallus</i>	<i>A. forsteri</i>	<i>E. africanus</i>
<i>E. ferus</i>	0	21	11	13	1
<i>D. polylepis</i>		0	18	17	20
<i>G. gallus</i>			0	3	10
<i>A. forsteri</i>				0	12
<i>E. africanus</i>					0

- (a) Using the data in the table, **create** a phylogenetic tree on the template provided to reflect the evolutionary relationships of the organisms. **Provide reasoning** for the placement on the tree of the species that is least related to the others.

Phylogenetic tree (1 point)

NOTE: There can be free rotation around the nodes in the tree.



Reasoning (1 point)

- *D. polylepis* has the most differences in amino acids (or changes in DNA or proteins as they relate to amino acids).
- (b) **Identify** whether morphological data or amino acid sequence data are more likely to accurately represent the true evolutionary relationships among the species, and **provide reasoning** for your answer.

2015 SCORING GUIDELINES**Question 3 (continued)****Identification (1 point)**

- Amino acid/molecular data

Reasoning (1 point)

- Morphology may be similar (due to convergent evolution/analogous structures) even if there are differences in amino acid/DNA sequences.
- Molecular data (e.g. amino acid changes, DNA changes) directly show genetic make-up/ reveal evolution.

OR

Identification (1 point)

- Morphological data

Reasoning (1 point)

- Similar molecular sequences may result in different morphologies.
- An example of species with similar proteins but different morphology (e.g., chimps and humans).

2015 SCORING GUIDELINES

Question 4

Both mitosis and meiosis are forms of cell division that produce daughter cells containing genetic information from the parent cell.

- (a) **Describe** TWO events that are common to both mitosis and meiosis that ensure the resulting daughter cells inherit the appropriate number of chromosomes.

Description (1 point each; 2 points maximum)


- Spindle elements (microtubules) form/attach to chromosomes
- Chromatin condenses
- Alignment of chromosomes across center of cell prior to chromosome separation
- Separation of chromatids/centromeres to daughter cells
- G2/M checkpoint occurs in both processes
- Replication or synthesis of DNA precedes mitosis/meiosis
- Cytokinesis separates daughter cells after mitosis/meiosis

- (b) The genetic composition of daughter cells produced by mitosis differs from that of the daughter cells produced by meiosis. **Describe** TWO features of the cell division processes that lead to these differences.

Feature	Description (1 point each row; 2 points maximum)	
	Mitosis	Meiosis
Number of divisions/ number of resulting cells	1 division/ 2 cells result	2 divisions/ 4 cells result
Ploidy of daughter cells	• Same as parent cell • Diploid • ($2n \rightarrow 2n$ or $n \rightarrow n$)	• Half of parent cell • Haploid • ($4n \rightarrow 2n$; $2n \rightarrow n$)
Chromatids separate	Occurs	Not in meiosis I/only in meiosis II
Crossing over	Does not occur	Occurs
Homologous chromosomes separate/independently assort	Does not occur	Occurs

2015 SCORING GUIDELINES

Question 5



Phototropism in plants is a response in which a plant shoot grows toward a light source. The results of five different experimental treatments from classic investigations of phototropism are shown above.

- (a) **Give support** for the claim that the cells located in the tip of the plant shoot detect the light by comparing the results from treatment group I with the results from treatment group II and treatment group III.

Support (2 points maximum)

- In treatment II the tip is removed and the plant no longer bends toward light.
- In treatment III the cap blocks the light to the tip and the plant no longer bends toward light.

- (b) In treatment groups IV and V, the tips of the plants are removed and placed back onto the shoot on either a permeable or impermeable barrier. Using the results from treatment groups IV and V, **describe** TWO additional characteristics of the phototropism response.

Description (2 points maximum)


- Tip produces a substance/signal/hormone (auxin) in response to light causing the plants to bend
- Substance must diffuse from the tip causing the plants to bend

2015 SCORING GUIDELINES

Question 6

In an attempt to rescue a small isolated population of snakes from decline, a few male snakes from several larger populations of the same species were introduced into the population in 1992. The snakes reproduce sexually, and there are abundant resources in the environment.

The figure below shows the results of a study of the snake population, both before and after the introduction of the outside males. In the study, the numbers of captured snakes is an indication of the overall population size.



- (a) **Describe** ONE characteristic of the original population that may have led to the population's decline in size between 1989 and 1993.

Description (1 point)

- Lacked genetic diversity/variation
- Was an aged/post-reproductive population/not enough young snakes
- Had unfavorable sex ratio/too few males
- Possessed a harmful mutation/disease

- (b) **Propose** ONE reason that the introduction of the outside males rescued the snake population from decline.

Proposal (1 point)

- Increased genetic diversity in the population
- Increased reproductive success
- Established beneficial sex ratio/sufficient proportion of males for reproduction
- Introduced resistance to disease that was affecting the original population

- (c) **Describe** how the data support the statement that there are abundant resources in the environment.

Description (1 point)

- Population can/does grow
- If resources are limited population would not grow

2015 SCORING GUIDELINES

Question 7

Smell perception in mammals involves the interactions of airborne odorant molecules from the environment with receptor proteins on the olfactory neurons in the nasal cavity. The binding of odorant molecules to the receptor proteins triggers action potentials in the olfactory neurons and results in transmission of information to the brain. Mammalian genomes typically have approximately 1,000 functional odorant-receptor genes, each encoding a unique odorant receptor.

- (a) **Describe** how the signal is transmitted across the synapse from an olfactory sensory neuron to the interneuron that transmits the information to the brain.

Description (1 point)

- Neurotransmitters are released from the olfactory neuron and bind to receptors in the postsynaptic neuron.

- (b) **Explain** how the expression of a limited number of odorant receptor genes can lead to the perception of thousands of odors. Use the evidence about the number of odorant receptor genes to **support** your explanation.

Explanation and support (2 points maximum: points may be earned from only one row)

	Explanation (1 point)	Support (1 point)
Molecular	• One odorant molecule can be recognized by more than one odorant receptor • One odorant receptor can bind to more than one odorant molecule	Mathematical combinations expand possible odors detected
CNS Control	Signals integrated in the brain	Multiple interactions among neurons in the brain
Genetic	Alternate processing/splicing (of pre-mRNA/primary transcript)	Multiple receptors can be produced from a gene

2015 SCORING GUIDELINES**Question 8**

An individual has lost the ability to activate B-cells and mount a humoral immune response.

- (a) **Propose** ONE direct consequence of the loss of B-cell activity on the individual's humoral immune response to an initial exposure to a bacterial pathogen.

Proposal (1 point)

- Does not produce antibodies
- Does not produce memory B cells

- (b) **Propose** ONE direct consequence of the loss of B-cell activity on the speed of the individual's humoral immune response to a second exposure to the bacterial pathogen.

Proposal (1 point)

- Does not mount a faster response than the first response
- Mounts a second response at the same speed as the first response
- Mounts a second response more slowly than would a normal individual

- (c) **Describe** ONE characteristic of the individual's immune response to the bacterial pathogen that is not affected by the loss of B cells.

Description (1 point)

- Cell-mediated immunity still active
- Components of cell mediated immunity (e.g., Natural Killer/ Cytotoxic T cells) still active
- Nonspecific immune response remains active
- Components of non-specific immunity (e.g., macrophage/ phagocyte, epidermis) still active