

3. Buffelgrass, an invasive grass species in southwestern desert ecosystems, is threatening the saguaro cactus, a keystone species in these ecosystems. Buffelgrass is drought-tolerant and can survive wildfires. However, the dry buffelgrass also acts as fuel for wildfires, causing the fires to be more severe. Older saguaro cacti can survive wildfires; however, many of the young cacti cannot.

Scientists conducted an experiment to determine whether they could control the abundance of the buffelgrass population. The scientists identified several native grass species that, when grown with buffelgrass, might reduce the abundance of buffelgrass. They grew buffelgrass in the presence of several different native grass species in greenhouses, in either nondrought (watered every 3 days) or drought (watered every 9 days) conditions. After twelve weeks, they measured the height and dry weight of the buffelgrass in each treatment group.

- A. **Describe** the effect that removing a keystone species will have on an ecosystem.
- B. **Identify** a control group the scientists should include in their experiment.
- C. **State** the null hypothesis of the experiment in which buffelgrass is grown in the presence of native grass species.
- D. Scientists have found that the population growth rates of native grasses are much slower than the population growth rate of buffelgrass following a wildfire. The scientists claim that wildfires will therefore increase the abundance of buffelgrass plants in the ecosystem. Based on the information given, **justify** the scientists' claim.

5. Annual plants complete their life cycle, including germination, seed production, and death, within one year. *Ambrosia trifida* (giant ragweed) is an annual plant that readily colonizes any land that has had a disturbance such as plowing. The plant is considered an invasive species in regions outside of its native range. In a particular region, the seeds of *A. trifida* germinate from early March through the end of the summer, while the seeds of other annual plants require warmer soil temperatures and thus germinate from late April through the end of the summer.

Researchers studied the influence of *A. trifida* on the biodiversity of other annual plant species that grow in the same field. In early spring, the researchers marked off identical plots of land in a field that had been plowed the previous fall and not replanted with new crops. All plants that grew on one half of the plots were left untouched (Figure 1A), while all germinating *A. trifida* seedlings were removed from the other half of the plots throughout the spring and summer (Figure 1B). In late summer, the researchers counted and identified all plants that grew in the plots. The distribution of plants is represented by the symbols in Figures 1A and 1B.




Figure 1. Representations of plant identity and distribution in experimental plots in late summer. Each box represents one typical experimental plot, and each symbol represents 10 individual plants.

- (a) **Describe** a cause of logistic growth of the ragweed population.
- (b) Based on the representation in Figure 1, **explain** why the scientists claim that plot B would be more resilient than plot A in response to a sudden environmental change.
- (c) In a third group of plots, the researchers removed all seedlings of all plants that germinated before June 1. All plants that germinated after June 1 were left untouched. Using the template in the space provided for your response and the symbols shown in Figure 1, **represent** the expected plant species that would be found in this third group of plots three months later. Draw no more than 12 symbols. Assume all other environmental conditions are the same as for the initial study described.
- (d) **Explain** how an invasive species such as ragweed affects ecosystem biodiversity, as illustrated in Figure 1.

Write your responses to this question only on the designated pages in the separate Free Response booklet.

2. A student studying two different aquatic, plant-eating, unicellular protist species (species A and B) designed an experiment to investigate the ecological relationship between the two species (Table 1).

TABLE 1. EXPERIMENTAL TREATMENT GROUPS


Group I.	Species A and B are each grown in separate containers.
Group II.	Species A and B are grown together in the same container.

In treatment group I, the student placed 10 individuals of species A into a container with liquid growth medium and 10 individuals of species B into a separate container with an equal amount of the same liquid growth medium. In treatment group II, the student placed 5 individuals of each species into a single container with the liquid growth medium. The student then maintained the containers under the same environmental conditions and recorded the number of individuals in each population at various time points. The results are shown in Table 2.

TABLE 2. NUMBER OF INDIVIDUALS IN EACH PROTIST POPULATION
IN BOTH TREATMENT GROUPS

Time (h)	Group I. Grown Separately		Group II. Grown Together	
	Species A	Species B	Species A	Species B
0	10	10	5	5
10	100	50	45	20
20	400	200	100	50
30	1100	500	250	25
40	1400	650	525	20
50	1500	700	900	10
60	1500	700	1250	0
70	1500	700	1400	0

- (a) The growth curves for species B in group I and for species A in group II (shaded columns) have been plotted on the template. Use the template to **complete** an appropriately labeled line graph to illustrate the growth of species A in treatment group I and species B in treatment group II (unshaded columns).
- (b) As shown in the table, the student established treatment group II with 5 individuals of each species. **Provide reasoning** for the reduced initial population sizes.
- (c) The student claims that species A and B compete for the same food source. **Provide TWO pieces of evidence** from the data that support the student's claim.
- (d) **Predict** TWO factors that will most likely limit the population growth of species A in treatment group I.
- (e) Many protists contain an organelle called a contractile vacuole that pumps water out of the cell. The student repeated the experiment using a growth medium with a lower solute concentration. **Predict** how the activity of the contractile vacuole will change under the new experimental conditions. **Justify** your prediction.







Figure 1. Bacterial population growth in the presence of two nutrients (nutrient I and nutrient II)

2. Bacteria can be cultured in media with a carefully controlled nutrient composition. The graph above shows the growth of a bacterial population in a medium with limiting amounts of two nutrients, I and II.
- Estimate** the maximum population density in $\frac{\text{cells}}{\text{mL}}$ for the culture. Using the data, **describe** what prevents further growth of the bacterial population in the culture.
 - Using the data, **calculate** the growth rate in $\frac{\text{cells}}{\text{mL} \times \text{hour}}$ of the bacterial population between hours 2 and 4.
 - Identify** the preferred nutrient source of the bacteria in the culture over the course of the experiment. Use the graph to **justify** your response. **Propose** ONE advantage of the nutrient preference for an individual bacterium.
 - Describe** how nutrient I most likely regulates the genes for metabolism of nutrient I and the genes for metabolism of nutrient II. **Provide** TWO reasons that the population does not grow between hours 5 and 6.

6. In an attempt to rescue a small isolated population of snakes from decline, a few male snakes from several larger populations of the same species were introduced into the population in 1992. The snakes reproduce sexually, and there are abundant resources in the environment.

The figure below shows the results of a study of the snake population both before and after the introduction of the outside males. In the study, the numbers of captured snakes indicate the overall population size.



- (a) **Describe** ONE characteristic of the original population that may have led to the population's decline in size between 1989 and 1993.
- (b) **Propose** ONE reason that the introduction of the outside males rescued the snake population from decline.
- (c) **Describe** how the data support the statement that there are abundant resources in the environment.

BIOLOGY**Section II****8 Questions****Planning Time—10 minutes****Writing Time—80 minutes**

Directions: Questions 1 and 2 are long free-response questions that require about 22 minutes each to answer and are worth 10 points each. Questions 3–8 are short free-response questions that require about 6 minutes each to answer. Questions 3–5 are worth 4 points each and questions 6–8 are worth 3 points each.

Read each question carefully and completely. Write your response in the space provided for each question. Only material written in the space provided will be scored. Answers must be written out in paragraph form. Outlines, bulleted lists, or diagrams alone are not acceptable.

1. Trichomes are hairlike outgrowths of the epidermis of plants that are thought to provide protection against being eaten by herbivores (herbivory). In a certain plant species, stem trichome density is genetically determined.

To investigate variation in stem trichome density within the plant species, a student counted the number of trichomes on the stems of six plants in each of three different populations. The student used the data to calculate the mean trichome density (numbers of hairs per square centimeter) for each population. The results are provided in the table below.

TRICHOME DENSITY IN THREE PLANT POPULATIONS (number of trichomes/cm²)

Population	Plant 1	Plant 2	Plant 3	Plant 4	Plant 5	Plant 6	Mean	Standard Error of the Mean (SEM)
I	8	11	9	10	8	6	9	1
II	12	6	15	9	13	8	11	1
III	13	17	9	14	12	16	14	1

- (a) On the axes provided, **create** an appropriately labeled graph to illustrate the sample means of the three populations to within 95% confidence (i.e., sample mean $\pm$ 2 SEM).
- (b) Based on the sample means and standard errors of the means, **identify** the two populations that are most likely to have statistically significant differences in the mean stem trichome densities. **Justify** your response.
- (c) **Describe** the independent and dependent variables and a control treatment for an experiment to test the hypothesis that higher trichome density in plants is selected for in the presence of herbivores. **Identify** an appropriate duration of the experiment to ensure that natural selection is measured, and **predict** the experimental results that would support the hypothesis.

