


The following information applies to all parts.

6. Many raptors in Africa are top predators and keystone species in their ecosystems. Raptor population sizes have been declining in recent years.

Scientists studied the relationship between the conversion of the raptors' natural habitat into farmland and the population sizes of raptors in those regions. They compiled raptor population size data from three different regions of Africa: Region 1, Region 2, and Region 3. In each of the regions, the scientists compared the annual percent change in the raptor population size in areas that were protected from farming (protected) with the annual percent change in areas where farming occurred (unprotected). Their results are shown in the box and whisker plot.

type of bird

Median Annual Percent Change in the Population Size of Raptors in Protected and Unprotected Areas



Part A

Based on the box and whisker plot, **identify** the median annual percent change in population size in the unprotected area of Region 2.

Part B

Based on the box and whisker plot, **identify** the region and area that experienced the greatest median annual percent change in population size.

Part C

Researchers hypothesize that if farming is eliminated in Region 2, there will no longer be a decline in the raptor population size. Based on the data in the box and whisker plot, **evaluate** the researchers' hypothesis.

Part D

Based on the information provided, **explain** why ecosystem resilience in unprotected areas may decline over time.

STOP
END OF EXAM

3. Buffelgrass, an invasive grass species in southwestern desert ecosystems, is threatening the saguaro cactus, a keystone species in these ecosystems. Buffelgrass is drought-tolerant and can survive wildfires. However, the dry buffelgrass also acts as fuel for wildfires, causing the fires to be more severe. Older saguaro cacti can survive wildfires; however, many of the young cacti cannot.

Scientists conducted an experiment to determine whether they could control the abundance of the buffelgrass population. The scientists identified several native grass species that, when grown with buffelgrass, might reduce the abundance of buffelgrass. They grew buffelgrass in the presence of several different native grass species in greenhouses, in either nondrought (watered every 3 days) or drought (watered every 9 days) conditions. After twelve weeks, they measured the height and dry weight of the buffelgrass in each treatment group.

- A. **Describe** the effect that removing a keystone species will have on an ecosystem.
- B. **Identify** a control group the scientists should include in their experiment.
- C. **State** the null hypothesis of the experiment in which buffelgrass is grown in the presence of native grass species.
- D. Scientists have found that the population growth rates of native grasses are much slower than the population growth rate of buffelgrass following a wildfire. The scientists claim that wildfires will therefore increase the abundance of buffelgrass plants in the ecosystem. Based on the information given, **justify** the scientists' claim.

4. The common wild oat is native to regions of Europe and Asia but is an invasive species in central California grasslands. In California, the common wild oat has almost completely replaced some species of native bunchgrass. Researchers found that aphids, a type of small insect that often carries plant viruses, have a much higher reproductive rate in grasslands that include the common wild oat than in grasslands composed of only native bunchgrass species. Additionally, the viruses carried by the aphids appear to affect only the native bunchgrasses and not the common wild oat. Native bunchgrasses infected by the virus have much higher death rates than do native bunchgrasses that are not infected.
- (a) **Describe** the change in the resilience of an ecosystem when there is a decrease in the number of species.
- (b) **Explain** how the addition of the common wild oat affects the number of native bunchgrass plants that can be supported by the California grasslands ecosystem.
- (c) Researchers suggest adding ladybugs, predators of aphids, to the California grasslands. **Predict** the effect of adding ladybugs on the abundance of the native bunchgrass population.
- (d) **Justify** your prediction in part (c).
-

Write your responses to this question only on the designated pages in the separate Free Response booklet.

If there are multiple parts to this question, write the part letter with your response.

5. The following models represent all the interacting species in two different communities with some of the same species and feeding relationships. These models assume that both communities have the same initial biomass. The models can be used to understand the effects of human activities on the communities.




Figure 1. Models of two different communities with some of the same species

- (a) **Describe** a characteristic of a community that makes a species invasive in that community but not invasive in a different community.
- (b) **Explain** why removing species PP1 will have a greater effect on community B than on community A.
- (c) An invasive species (INV) that eats individuals of species SC2 is introduced into community B. Using the template in the space provided for your response, for community B, indicate the feeding relationship for this invasive species by correctly placing **INV** to represent the invasive species and an **arrow** to represent the feeding relationship within community B.
- (d) **Explain** how human activities that add toxins to the soil could change a community with many species at each trophic level, such as community A, into a community with few species at each trophic level, such as community B.

Write your responses to this question only on the designated pages in the separate Free Response booklet.




Figure 1. The effect of karrikins (KAR) and trimethylbutenolides (TMB) on seed germination in *Lactuca* plants. Error bars represent $\pm 2SE_{\bar{x}}$.

2. Fires frequently occur in some ecosystems and can destroy all above-ground vegetation. Many species of plants in these ecosystems respond to compounds in smoke that regulate seed germination after a major fire. Karrikins (KAR) and trimethylbutenolides (TMB) are water-soluble compounds found in smoke that are deposited in the soil as a result of a fire. KAR and TMB bind to receptor proteins in a seed. In a study on the effects of smoke on seeds, researchers recorded the timing and percent of seed germination in the presence of various combinations of KAR and TMB. The results are shown in Figure 1.

In a second investigation into the effect of available water on seed germination after a fire, researchers treated seeds with KAR or TMB. The treated seeds were then divided into two treatment groups. One group received a water rinse and the other group received no water rinse. The seeds were then incubated along with a group of control seeds that were not treated. The results are shown in the table.

EFFECT OF CHEMICAL TREATMENT AND WATER RINSE ON GERMINATION

Treatment Group	Chemical Treatment		Water	Germination Result
	KAR	TMB		
1 (control)	–	–	–	Control result
2	+	–	–	Different from control
3	–	+	–	Different from control
4 (control)	–	–	+	Control result
5	+	–	+	Different from control
6	–	+	+	Same as control

(a) The researchers made the following claims about the effect of KAR and the effect of TMB on seed germination relative to the control treatment.

- KAR alone affects the timing of seed germination.
- KAR alone affects the percentage of seeds that germinate.
- TMB alone affects the timing of seed germination.
- TMB alone affects the percentage of seeds that germinate.

Provide support using data from Figure 1 for each of the researchers' claims.

- (b) **Make a claim** about the effect of rinsing on the binding of KAR to the receptor in the seed and about the effect of rinsing on the binding of TMB to the receptor in the seed. Identify the appropriate treatment groups and results from the table that, when compared with the controls, **provide support** for each claim.
- (c) There is intense competition by plants to successfully colonize areas that have been recently cleared by a fire. **Describe** ONE advantage of KAR regulation and ONE advantage of TMB regulation to plants that live in an ecosystem with regular fires.




Figure 1. Detectability of eDNA fragments of varying lengths




Figure 2. Map of the waterways that connect a nearby river system to Lake Michigan


6. Living and dead organisms continuously shed DNA fragments, known as eDNA, into the environment. To detect eDNA fragments in the environment, the polymerase chain reaction (PCR) can be used to amplify specific eDNA fragments. eDNA fragments of different lengths persist in the environment for varying amounts of time before becoming undetectable (Figure 1).

To investigate whether silver carp, an invasive fish, have moved from a nearby river system into Lake Michigan, researchers tested water samples for the presence of eDNA specific to silver carp (Figure 2).

- (a) **Justify** the use of eDNA sampling as an appropriate technique for detecting the presence of silver carp in an environment where many different species of fish are found. **Propose** ONE advantage of identifying long eDNA fragments as opposed to short fragments for detecting silver carp.
- (b) The researchers tested a large number of water samples from Lake Michigan and found eDNA specific to silver carp in a single sample in the lake, as indicated in Figure 2. The researchers concluded that the single positive sample was a false positive and that no silver carp had entered Lake Michigan. **Provide reasoning** other than human error to support the researchers' claim.

3. As part of a new suburban development, a sports complex consisting of athletic fields and buildings is constructed in a formerly wooded area.
- Predict** ONE ecological consequence on the local plant community that is likely to result during the site preparation and construction of the sports complex. **Justify** your prediction.
 - To maintain the playing fields, large quantities of water and chemicals are applied regularly to the grass-covered areas. **Predict** ONE effect on the local animal community that might result from regular use and maintenance of the playing fields. **Justify** your prediction.
4. Adult male guppies (*Poecilia reticulata*) exhibit genetically determined spots, while juvenile and adult female guppies lack spots. In a study of selection, male and female guppies from genetically diverse populations were collected from different mountain streams and placed together in an isolated environment containing no predators.

The study population was maintained for several generations in the isolated area before being separated into two groups. One group was moved to an artificial pond containing a fish predator, while a second group was moved to an artificial pond containing no predators. The two groups went through several generations in their new environments. At different times during the experiment, the mean number of spots per adult male guppy was determined as shown in the figure below. Vertical bars in the figure represent two standard errors of the mean (SEM).



- Describe** the change in genetic variation in the population between 0 and 6 months and **provide** reasoning for your description based on the means and SEM.
 - Propose** ONE type of mating behavior that could have resulted in the observed change in the number of spots per adult male guppy between 6 and 20 months in the absence of the predator.
 - Propose** an evolutionary mechanism that explains the change in average number of spots between 6 and 20 months in the presence of the predator.
-
5. Genetically modified crops have been developed that produce a protein that makes the plants resistant to insect pests. Other genetic modifications make the crops more resistant to chemicals that kill plants (herbicides).
- Describe** TWO potential biological risks of large-scale cultivation and use of such genetically modified plants.
 - For each of the risks you described in part (a), **propose** a practical approach for reducing the risk.